

# Lecture PowerPoints

## Chapter 14 *Physics: Principles with Applications, 7<sup>th</sup> edition* Giancoli

© 2014 Pearson Education, Inc.

This work is protected by United States copyright laws and is provided solely for the use of instructors in teaching their courses and assessing student learning. Dissemination or sale of any part of this work (including on the World Wide Web) will destroy the integrity of the work and is not permitted. The work and materials from it should never be made available to students except by instructors using the accompanying text in their classes. All recipients of this work are expected to abide by these restrictions and to honor the intended pedagogical purposes and the needs of other instructors who rely on these materials.

# Chapter 14

## Heat



# Contents of Chapter 14

- Heat As Energy Transfer
- Internal Energy
- Specific Heat
- Calorimetry—Solving Problems
- Latent Heat
- Heat Transfer: Conduction
- Heat Transfer: Convection
- Heat Transfer: Radiation

# 14-1 Heat As Energy Transfer

We often speak of heat as though it were a material that flows from one object to another; it is not. Rather, it is a form of energy.


Unit of heat: calorie (cal)

1 cal is the amount of heat necessary to raise the temperature of 1 g of water by 1 Celsius degree.

Don't be fooled—the calories on our food labels are really kilocalories (kcal or Calories), the heat necessary to raise 1 kg of water by 1 Celsius degree.

# 14-1 Heat As Energy Transfer

If heat is a form of energy, it ought to be possible to equate it to other forms. The experiment below found the mechanical equivalent of heat by using the falling weight to heat the water:



$$4.186 \text{ J} = 1 \text{ cal}$$

$$4.186 \text{ kJ} = 1 \text{ kcal}$$

# 14-1 Heat As Energy Transfer

Definition of heat:

Heat is energy transferred from one object to another because of a difference in temperature.

- Remember that the temperature of a gas is a measure of the kinetic energy of its molecules.

## 14-2 Internal Energy

The sum total of all the energy of all the molecules in a substance is its internal (or thermal) energy.

Temperature: measures molecules' average kinetic energy

Internal energy: total energy of all molecules

Heat: transfer of energy due to difference in temperature

## 14-2 Internal Energy

Internal energy of an ideal (atomic) gas is equal to the average kinetic energy per molecule multiplied by the number of molecules.

But since we know the average kinetic energy in terms of the temperature, we can write:



$$U = \frac{3}{2}nRT,$$

$$\left[ \begin{array}{l} \text{internal energy of} \\ \text{ideal monatomic gas} \end{array} \right] \quad (14-1)$$



# 14-2 Internal Energy

If the gas is molecular rather than atomic, rotational and vibrational kinetic energy needs to be taken into account as well.



# 14-3 Specific Heat

**TABLE 14-1 Specific Heats**  
(at 1 atm constant pressure and 20°C  
unless otherwise stated)

Substance	Specific Heat, $c$	
	J/kg · C°	kcal/kg · C° (= cal/g · C°)
Aluminum	900	0.22
Alcohol (ethyl)	2400	0.58
Copper	390	0.093
Glass	840	0.20
Iron or steel	450	0.11
Lead	130	0.031
Marble	860	0.21
Mercury	140	0.033
Silver	230	0.056
Wood	1700	0.4
Water		
Ice (−5°C)	2100	0.50
Liquid (15°C)	4186	1.00
Steam (110°C)	2010	0.48
Human body (average)	3470	0.83
Protein	1700	0.4

The amount of heat required to change the temperature of a material is proportional to the mass and to the temperature change:

$$Q = mc \Delta T, \quad (14-2)$$

The specific heat,  $c$ , is characteristic of the material. Some values are listed at left.

# 14-3 Specific Heat

Specific heats of gases are more complicated, and are generally measured at constant pressure ( $c_p$ ) or constant volume ( $c_v$ ).

Some sample values:

**TABLE 14–2**  
**Specific Heats of Gases**  
(kcal/kg · C°)

<b>Gas</b>	<b><math>c_p</math> (constant pressure)</b>	<b><math>c_v</math> (constant volume)</b>
Steam (100°C)	0.482	0.350
Oxygen	0.218	0.155
Helium	1.15	0.75
Carbon dioxide	0.199	0.153
Nitrogen	0.248	0.177

# 14-4 Calorimetry—Solving Problems

Closed system: no mass enters or leaves, but energy may be exchanged

Open system: mass may transfer as well


Isolated system: closed system where no energy in any form is transferred

For an isolated system,

Energy out of one part = energy into another part

Or:  $\text{heat lost} = \text{heat gained}$

# 14-4 Calorimetry – Solving Problems



The instrument to the left is a calorimeter, which makes quantitative measurements of heat exchange. A sample is heated to a well-measured high temperature, plunged into the water, and the equilibrium temperature measured. This gives the specific heat of the sample.


## 14-4 Calorimetry—Solving Problems

Another type of calorimeter is called a bomb calorimeter; it measures the thermal energy released when a substance burns.

This is the way the caloric content of foods is measured.

# 14-5 Latent Heat

Energy is required for a material to change phase, even though its temperature is not changing.



# 14-5 Latent Heat

Heat of fusion,  $L_F$ : heat required to change 1.0 kg of material from solid to liquid

Heat of vaporization,  $L_V$ : heat required to change 1.0 kg of material from liquid to vapor

Substance	Melting Point (°C)	Heat of Fusion		Boiling Point (°C)	Heat of Vaporization	
		kJ/kg	kcal/kg <sup>†</sup>		kJ/kg	kcal/kg <sup>†</sup>
Oxygen	−218.8	14	3.3	−183	210	51
Nitrogen	−210.0	26	6.1	−195.8	200	48
Ethyl alcohol	−114	104	25	78	850	204
Ammonia	−77.8	33	8.0	−33.4	137	33
Water	0	333	79.7	100	2260	539
Lead	327	25	5.9	1750	870	208
Silver	961	88	21	2193	2300	558
Iron	1538	289	69.1	3023	6340	1520
Tungsten	3410	184	44	5900	4800	1150

<sup>†</sup>Numerical values in kcal/kg are the same in cal/g.



# 14-5 Latent Heat

The total heat required for a phase change depends on the total mass and the latent heat:

$$Q = mL, \quad (14-4)$$

## Problem Solving: Calorimetry

1. Is the system isolated? Are all significant sources of energy transfer known or calculable?
2. Apply conservation of energy.
3. If no phase changes occur, the heat transferred will depend on the mass, specific heat, and temperature change.

## 14-5 Latent Heat

4. If there are, or may be, phase changes, terms that depend on the mass and the latent heat may also be present. Determine or estimate what phase the final system will be in.
5. Make sure that each term is in the right place and that all the temperature changes are positive.
6. There is only one final temperature when the system reaches equilibrium.
7. Solve.

## 14-5 Latent Heat

The latent heat of vaporization is relevant for evaporation as well as boiling. The heat of vaporization of water rises slightly as the temperature decreases.


On a molecular level, the heat added during a change of state does not go to increasing the kinetic energy of individual molecules, but rather to break the close bonds between them so the next phase can occur.

# 14-6 Heat Transfer: Conduction

Heat conduction can be visualized as occurring through molecular collisions.

The heat flow per unit time is given by:

$$\frac{Q}{t} = kA \frac{T_1 - T_2}{\ell} \quad (14-5)$$



# 14-6 Heat Transfer: Conduction

**TABLE 14-4**  
**Thermal Conductivities**

Substance	Thermal Conductivity, $k$	
	J ( $\text{s} \cdot \text{m} \cdot \text{C}^\circ$ )	kcal ( $\text{s} \cdot \text{m} \cdot \text{C}^\circ$ )
Silver	420	$10 \times 10^{-2}$
Copper	380	$9.2 \times 10^{-2}$
Aluminum	200	$5.0 \times 10^{-2}$
Steel	40	$1.1 \times 10^{-2}$
Ice	2	$5 \times 10^{-4}$
Glass	0.84	$2.0 \times 10^{-4}$
Brick	0.84	$2.0 \times 10^{-4}$
Concrete	0.84	$2.0 \times 10^{-4}$
Water	0.56	$1.4 \times 10^{-4}$
Human tissue	0.2	$0.5 \times 10^{-4}$
Wood	0.1	$0.3 \times 10^{-4}$
Fiberglass	0.048	$0.12 \times 10^{-4}$
Cork	0.042	$0.10 \times 10^{-4}$
Wool	0.040	$0.10 \times 10^{-4}$
Goose down	0.025	$0.060 \times 10^{-4}$
Polyurethane	0.024	$0.057 \times 10^{-4}$
Air	0.023	$0.055 \times 10^{-4}$

The constant  $k$  is called the thermal conductivity.

Materials with large  $k$  are called conductors; those with small  $k$  are called insulators.

# 14-6 Heat Transfer: Conduction

Building materials are measured using  $R$ -values rather than thermal conductivity:

$$R = \frac{l}{k}.$$



Here,  $l$  is the thickness of the material.

**TABLE 14-5  $R$ -values**

<b>Material</b>	<b>Thickness</b>	<b><math>R</math>-value (ft<sup>2</sup>·h·F°/Btu)</b>
Glass	$\frac{1}{8}$ inch	1
Brick	$3\frac{1}{2}$ inches	0.6–1
Plywood	$\frac{1}{2}$ inch	0.6
Fiberglass insulation	4 inches	12

# 14-7 Heat Transfer: Convection

Convection occurs when heat flows by the mass movement of molecules from one place to another. It may be natural or forced; both these examples are natural convection.



# 14-7 Heat Transfer: Convection

Many home heating systems are forced hot-air systems; these have a fan that blows the air out of registers, rather than relying completely on natural convection.

Our body temperature is regulated by the blood; it runs close to the surface of the skin and transfers heat. Once it reaches the surface of the skin, the heat is released through convection, evaporation, and radiation.



# 14-8 Heat Transfer: Radiation



The most familiar example of radiation is our own Sun, which radiates at a temperature of almost 6000 K.

# 14-8 Heat Transfer: Radiation

The energy radiated has been found to be proportional to the fourth power of the temperature:

$$\frac{Q}{t} = \epsilon \sigma AT^4. \quad (14-6)$$

The constant  $\sigma$  is called the Stefan-Boltzmann constant:

$$\sigma = 5.67 \times 10^{-8} \text{ W/m}^2 \cdot \text{K}^4$$

The emissivity  $e$  is a number between zero and one characterizing the surface; black objects have an emissivity near one, while shiny ones have an emissivity near zero.


## 14-8 Heat Transfer: Radiation

If you are sitting in a place that is too cold, your body radiates more heat than it can produce. You will start shivering and your metabolic rate will increase unless you put on warmer clothing.

# 14-8 Heat Transfer: Radiation


If you are in the sunlight, the Sun's radiation will warm you. In general, you will not be perfectly perpendicular to the Sun's rays, and will absorb energy at the rate:

$$\frac{Q}{t} = (1000 \text{ W/m}^2)\epsilon A \cos \theta, \quad (14-8)$$



# 14-8 Heat Transfer: Radiation

This  $\cos \theta$  effect is also responsible for the seasons.



# 14-8 Heat Transfer: Radiation

Thermography—the detailed measurement of radiation from the body—can be used in medical imaging.

Warmer areas may be a sign of tumors or infection; cooler areas on the skin may be a sign of poor circulation.



(a)



(b)

# Summary of Chapter 14

- Internal energy  $U$  refers to the total energy of all molecules in an object. For an ideal monatomic gas,

$$U = \frac{3}{2}nRT, \quad \left[ \begin{array}{l} \text{internal energy of} \\ \text{ideal monatomic gas} \end{array} \right] \quad (14-1)$$

- Heat is the transfer of energy from one object to another due to a temperature difference. Heat can be measured in joules or in calories.
- Specific heat of a substance is the energy required to change the temperature of a fixed amount of matter by  $1^\circ \text{C}$ .

# Summary of Chapter 14

- In an isolated system, heat gained by one part of the system must be lost by another.
- Calorimetry measures heat exchange quantitatively.
- Phase changes require energy even though the temperature does not change.
- Heat of fusion: amount of energy required to melt 1 kg of material.
- Heat of vaporization: amount of energy required to change 1 kg of material from liquid to vapor.



# Summary of Chapter 14

- Heat transfer takes place by conduction, convection, and radiation.
- In conduction, energy is transferred through the collisions of molecules in the substance.
- In convection, bulk quantities of the substance flow to areas of different temperature.
- Radiation is the transfer of energy by electromagnetic waves.