

MACROBOLIC NUTRITION

**PRIMING YOUR BODY TO
BUILD MUSCLE & BURN FAT**

Gerard Dente
with Kevin J. Hopkins



The information contained in this book is based upon the research and personal and professional experiences of the authors. It is not intended as a substitute for consulting with your physician or other healthcare provider. Any attempt to diagnose and treat an illness should be done under the direction of a healthcare professional.

The publisher does not advocate the use of any particular healthcare protocol but believes the information in this book should be available to the public. The publisher and authors are not responsible for any adverse effects or consequences resulting from the use of the suggestions, preparations, or procedures discussed in this book. Should the reader have any questions concerning the appropriateness of any procedures or preparation mentioned, the authors and the publisher strongly suggest consulting a professional healthcare advisor.

The author(s) of this book have a financial interest in some of the products recommended herein. This does not constitute an endorsement by Basic Health Publications, Inc.

Basic Health Publications, Inc.

8200 Boulevard East • North Bergen, NJ 07047 • 1-201-868-8336

Library of Congress Cataloging-in-Publication Data

Dente, Gerard, 1967–

Macrobiotic nutrition : priming your body to build muscle and burn body fat / Gerard Dente with Kevin J. Hopkins.
p. cm.

Includes bibliographical references and index.

ISBN 1-59120-131-4

1. Bodybuilders—Nutrition. 2. Macrobiotic diet. I. Hopkins, Kevin J., 1972– II. Title.

TX361.B64D46 2004
613.2'024'79641—dc22

2004003217

Copyright © 2004 by Gerard Dente

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording, or otherwise, without the prior written consent of the copyright owner.

Editor: Carol Rosenberg

Typesetter/Book design: Gary A. Rosenberg

Cover photo: Per Bernal

Kurt Angle photo on page 143 owned by World Wrestling Entertainment, Inc. and is reproduced with permission.

Printed in the United States of America

10 9 8 7 6 5 4 3 2 1

Contents

Acknowledgments, v

Preface, vii

Introduction You're a Bodybuilder, So Eat Like One!, 1

1. Understanding Macrobiotic Nutrition, 3
2. Carbohydrates—Don't Count Them Out, 12
3. Protein—Choosing the Right Building Blocks, 28
4. Fat—It Is Essential!, 61
5. Water—The Nutritional Paradox, 68
6. Macrobiotic Meals to Muscle Mass, 71
7. Guide to Estimating Caloric Requirements, 83
8. The Important Role of Micronutrients, 92
9. Enhancing the Macrobiotic Nutrition Effect, 119
10. Macrobiotic Meals, 150

Conclusion Progress Is the Great Motivator, 175

Appendix A Macrobiotic Caloric Requirements Tables, 176

Appendix B Macrobiotic Food Guide Pyramids, 189

Appendix C Macrobiotic Exchange Lists, 191

References, 195

Index, 207

About the Authors, 214




Photo by Irvin J. Geib.

Acknowledgments

THERE ARE MANY PEOPLE I WOULD LIKE TO THANK for their help, support, and friendship, but to name them all would take pages. I have been extremely fortunate to have so many special people in my life.

I want to give a special thanks to my mom, dad, sisters, brother, and my wife, Linda, for all of their continued support and for always allowing me to pursue my ambitions and dreams.

I want to thank all of the athletes I work with, particularly those mentioned in this book. Working with and knowing each of you is an honor and a pleasure. Each one of you is a champion. I especially want to thank Dave Hawk for our friendship and Kurt Angle, a great athlete and role model, and an even greater human being.

I am fortunate to be able to work in an industry I truly love. So, along with the people I just mentioned, I want to thank all the people I work with at MHP and all those I work with on a regular basis in the nutrition industry who have given me so much support and so many opportunities. You know who you are and I want to say . . . Thank you! Without your support, I would not be able to continually pursue my passion to research and develop new products and educate myself and others on better ways to improve health and performance.

—Gerard Dente



Photo by Irvin J. Geib.

Preface

I HAVE DEDICATED MOST OF MY LIFE to finding ways to increase muscle mass and improve physical performance. It started with my own personal quest to excel in high school football and then as a national-level competitive bodybuilder. I was fortunate enough to have fairly good genetics for building muscle. However, I knew that in order to compete at the top-level ranks against people with equal or better genetics, I had to compensate through a superior training and nutrition program. I know the extra knowledge of nutrition I gathered and applied toward my program helped me excel further than either I or anyone else had expected. During the time I was competing, I was able to maintain an off-season body weight of 285 to 295 pounds at only 10 percent body fat and a competition weight of 250 pounds at 4 to 5 percent body fat. This was back in 1995, an era when competing at 250 pounds was way above the norm.

For most bodybuilders, the goal is usually to win in competition, but the real motivation is to achieve the kind of personal progress that involves continually taking your body to the next level. That is what is so great about weightlifting and bodybuilding. If your focus is on your body and your external goals—the complete concept of what you want to achieve—bodybuilding can be very fulfilling and rewarding.

My personal quest to build my physique to its maximum potential through proper training and nutrition led me to an exploration of the research and development of sports-enhancing supplementation. Because I relied on nutrition to gain a competitive advantage over my competition and to help me reach my full genetic potential, I dedicated a lot of time to researching nutrition and supplementation and their effects on muscle building and performance. As I learned more, I realized that there was a lot of room for improvement and advancement in the sports supplements being offered to athletes, and this

ultimately led to the creation of Maximum Human Performance, Inc., in 1997. MHP's company mission is to provide athletes with the latest advancements in sports supplementation.

In addition to deriving inspiration from making progress and improvements in my own physique, I am inspired by public speaking and by helping others to achieve their fitness goals. I have given many lectures and seminars on training and nutrition over the years. Also, when you are a bodybuilder, your body serves as your uniform and it's obvious that you work out. You become a target for questions; so, no matter where you are, someone is almost sure to ask you a question about training, diet, or supplementation. I must have fielded tens of thousands of questions over the years.

It became obvious to me, by the number and kinds of questions that most people ask, that they are completely uninformed about nutrition, and it is disheartening to see how frustrated, confused, and misled they consequently become. I've seen too many people working hard at the gym and not getting results. I've answered questions like "How do I get big?" "What should I eat?" and "What supplements should I take?" too many times. It is for this reason that I felt compelled to write *Macrobiotic Nutrition*.

If you've ever asked or wondered about any of these questions . . . if you train hard and are unhappy with the results you've gotten, this book is for you! The answers and results lie within *Macrobiotic Nutrition's* 45/35/20 lean-mass equation.

I don't proclaim to be a "diet guru," nor am I going to try to take credit for any great diet revolution. In *Macrobiotic Nutrition*, I present scientific truths about nutrition in a simple, logical fashion. This book will help you understand the enormous impact food has on your body. Macrobiotic Nutrition can be applied to anyone who is working out and who wants to build muscle and burn body fat. Those of you who are hardcore will see that Macrobiotic Nutrition is the only way to make serious gains in "lean mass" and reach your full genetic potential. Macrobiotic Nutrition is not a quick fix: it is a long-term permanent solution! There are other diets and nutrition programs that may help you lose body fat, but these diets typically compromise your ability to pack on muscle. Macrobiotic Nutrition will turn your body into an efficient muscle-building, fat-burning machine! I am confident that Macrobiotic Nutrition will work for you as it has for me and for many others.

—Gerard Dente



WHEN GERARD DENTE APPROACHED ME to help write this book, I was skeptical. I understood that there was something missing within this industry, and basic nutrition was it. Gerard and I are on the same page when talking nutrition, but I was just a little concerned that we would be trying to appeal to the masses and not deliver the facts we know actually work. I soon found out that Gerard had in mind exactly what I felt was needed to guarantee bodybuilding success.

My background is in pharmacy and biochemistry, so naturally, the hormonal environment created in the body by foods and supplements was a vital interest of mine. For a few years, I worked with one of the largest compounding pharmacies in the country. I was exposed to hormonal replacement therapy practiced by some of the top physicians in the world during that time. The missing link for most patients was addressing their nutritional protocol. This slight oversight actually decreased the effectiveness of their therapy. As time went on, my love for bodybuilding and interest in nutrition drove me to open a health food store in 1995. The business grew and satisfied my desire to help educate people on the correct usage of vitamins, sports supplements, and herbs. Finally, I became a partner in a large fitness center and moved the entire health food store into the facility in 2003. I now had the ultimate approach for serving the bodybuilding community: a large state-of-the-art fitness center complete with all the cutting-edge supplements to enhance performance. I also began competing on the state level in the NPC as an open middleweight-class bodybuilder. My conditioning at contest time and my ability to stay ripped year-round drew a lot of attention from fellow competitors and patrons of our gym. Constantly barraged with questions, I felt it necessary to put it all in writing so the explanation would be easily followed.

I met Gerard quite some time ago, through my work with different physicians. One of his close friends was a client of the pharmacy where I worked, and I was very familiar with Gerard's lengthy list of bodybuilding accomplishments. Aware of each other, we finally met by chance at a New Jersey NPC bodybuilding competition. Gerard was there to launch a new supplement. We immediately struck up a conversation, and much to my surprise, he was very knowledgeable. My background in pharmacy and biochemistry intrigued him, and a friendship was formed instantly. I was very impressed with the fact that Gerard had a level of nutritional knowledge few bodybuilders could hope to possess. Coupled with his freaky genetics and 250-pound competition weight, this insightful look into nutrition turned him into an all-out superfreak!

Not long after our chance meeting, Gerard asked me to assist with the production of some new supplements to propel his company to the top of the industry. I agreed, and over the next few years TRAC, T-BOMB, TakeOFF, and their latest creation “The Up Your MASS” line were born. My background in pharmacy and drug-delivery systems proved to be invaluable for the invention of the enteric-coated delivery system of T-BOMB, which quickly became the number-one-selling prohormone in the country.

Now our chance meeting has evolved into my coauthoring *Macrobiolic Nutrition*. I strongly encourage any and all athletes who are serious about lean muscle growth to try the Macrobiolic Nutrition food plan. We address everything from protein, carbohydrates, and fats to the effects these macronutrients have on the hormones in the body. This book will be the most valuable tool in your muscle-building arsenal—just as important as the weights you lift and the recovery time you require. Macrobiolic Nutrition will put you at the top of the food chain for the maximal results from your training efforts.

—Kevin J. Hopkins

INTRODUCTION

You're a Bodybuilder, So Eat Like One!

YOU'RE DEDICATED AND COMMITTED to building an impressive, sculptured, muscular physique. You train four, five, or maybe even six times per week and stick to a strict diet to achieve the same look as those impressive professional bodybuilders you see in the magazines. Some of you may have been working hard at it for years, while some of you may be just getting started. For those of you who have been working at it for years, I say, "I'm sorry that you have been so dedicated and worked so hard, and still haven't figured out why you're not making serious gains and continual progress." For those of you just starting out, I say, "Congratulations—you are about to save yourself from years of hard work, trial and error, and frustration and misery." So, what is the secret of top bodybuilders? What do they know that you don't know? Why does the modern-day bodybuilder keep getting bigger and bigger and freaky ripped?

Come on; take a guess. I'll give you some clues. It's not their training, and it's something you consume every day. If you didn't figure it out, I'm going to tell you: **FOOD!** Yes, something as simple as food makes the whole difference. The foods you eat can have a tremendous impact on building a lean, muscular physique—even greater than your training. We all train hard yet we all can't achieve the degree of hard muscularity we desire. And don't blame your genetics, because unless you fuel your body with the proper nutrition, you will never reach your full genetic potential. Granted, we can't all be Mr. Olympia, but we can certainly improve our physiques. I'm not talking about little changes. I'm talking about changing your biochemistry and priming your body to become a muscle-building/fat-burning machine. Your body will efficiently build muscle and burn off body fat twenty-four hours a day, seven days a week, and 365 days a year. Your muscle mass will continue to increase, your body fat will go down to your desired level, and your strength and energy will go through the roof. All of these changes will occur simply by changing the way you eat. Oh,

I almost forgot to mention, you won't be doing cardio anymore! (You'll have to keep reading to find out how to get ripped without cardio.)

No doubt you've tried all kinds of diets and none of them worked—that's my point! Macrobiotic Nutrition isn't like any other diet. Macrobiotic Nutrition was designed for the hardcore, serious bodybuilder/athlete aiming to pack on lean muscle mass, increase strength, and improve performance. The nutritional requirements of bodybuilders and athletes far exceed those of the average person, because the physical demands a bodybuilder places on his or her body greatly increase the need for nutrition to help the body build and repair itself. When you add the goal of building huge amounts of muscle while decreasing body fat into the mix, it's easy to see why most people never reach their full growth potential.

The purpose of this book is to give you an understanding of food's impact on the many biochemical processes in the body that influence muscle growth and fat burning. The science behind Macrobiotic Nutrition is very advanced, but I have simplified it to make it easier to understand, and most important, so that you can apply it to achieving your goal to pack on lean mass. You may feel the information is too detailed or scientific at times, but keep reading and it will all come together. Up until now, how to get big and "shredded" has remained a mystery to you. *Macrobiotic Nutrition* is like a good mystery novel. You will pick up key pieces of information, "clues," throughout the book on how to get massive and ripped, and by its conclusion, you will have finally figured out the mystery of how to pack on lean mass!

Here are some highlights of what you are about to learn: First and foremost, not all calories are created equal. *Macrobiotic Nutrition* teaches you how to choose the right sources of carbohydrates, proteins, and fats. Macrobiotic Nutrition's 45/35/20 lean-mass equation shows you how to eat these sources in the proper ratio of macronutrients so they are optimized for your ability to build muscle, burn fat, and increase performance. You'll also learn about the importance of water, vitamins, and minerals.

Macrobiotic Nutrition's "Guide to Estimating Calories" will help you calculate the exact amount of carbohydrates, proteins, and fats that you need to achieve your goals. This book even provides a Macrobiotic meal menu, so you can eat a variety of delicious foods and make Macrobiotic Nutrition an enjoyable and easy-to-follow program. If you want to be at your best and make continual gains in your bodybuilding efforts, Macrobiotic Nutrition will get you there.

CHAPTER 1

Understanding Macrobiotic Nutrition

EVER WONDER WHY ALL OF THE BIGGEST and most ripped guys in the gym are on the weight room floor moving heavy steel, while all of the overweight or skinny guys are on the cardio machines?

How would you like to be able to pack on muscle and lose body fat without ever having to do cardio again? Yes, you read it right. Pack on muscle and lose body fat without *ever* doing cardio! You are about to learn how the top bodybuilders in the world pack on muscle and stay hard year-round, without cardio and without starving themselves. It all comes down to food, and what professional bodybuilders call Macrobiotic Nutrition's lean-mass equation. Macrobiotic Nutrition heralds a new era in bodybuilding and performance enhancement. This nutrition program is designed to create the ideal metabolic and hormonal environment for incredible gains in muscle size, strength, and endurance, while burning fat quickly and efficiently. In fact, your body will become so metabolically efficient, you won't have to do cardio to burn body fat. You won't have to starve yourself either. You'll be eating more food than you could imagine every two and a half to three hours. But before I get into the specifics and science of Macrobiotic Nutrition's lean-mass equation, let's clear the air right now about the biggest mistake amateur bodybuilders make in their quest for rock-hard mass—low-carb diets. The huge amount of media attention on low-carb diets may have led you to believe that was how to get a lean, muscular physique. Unfortunately, you probably found out the hard way that a low-carb diet isn't giving you the results you want. Rather than becoming big and hard with dense, full muscles, you feel small, flat, and weak! If you don't believe me, keep reading, because science doesn't lie. Not only does science prove it, the impressive mass achieved by the world's top bodybuilders proves it as well.



Photo by Reg Bradford.

Testimonial

“Early in my bodybuilding career, I made the mistake of going on a low-carb diet the last twelve weeks before a show. I always ended up losing a ton of muscle mass and always showed up looking flat, small, and smooth. Macrobiolic Nutrition has made all the difference in the world. Now, instead of losing mass pre-contest, I actually grow right into the show. This allows me to hit the stage looking hard, dense, and full.”

—Chris “**Big Guns**” Bennett,
NPC Top National Bodybuilder

MACROBOLIC NUTRITION'S 45/35/20 LEAN-MASS EQUATION

The 45/35/20 lean-mass equation is the foundation of Macrobiolic Nutrition. This will be the equation for your success in achieving your muscle-building and performance goals. 45/35/20 represents the ratio of carbohydrates, proteins, and fats you need to consume in your diet. I didn't pick these numbers randomly. They are based on proven science and my own personal experience of what your body needs to stimulate muscle growth, burn fat, and support recovery. If you are putting your body through rigorous workouts to build muscle and burn body fat, you better be taking in the proper nutrition. Otherwise, all your training efforts are going to be for naught, and you'll never make the gains you're looking for. The reason why most people never reach their full growth potential is not because of their lack of training, but because of their lack of nutrition.

Macrobiolic Nutrition heralds a new era in performance enhancement with its design to create the ideal metabolic and hormonal environment for increased muscle size, strength, and endurance, while reducing body fat. Macrobiolic's fundamental principle that “all calories are not created equal” is sure to raise controversy. Many other diets are based on the simple premise that caloric intake minus caloric expenditure determines weight gain or weight loss. This is true to an extent, but it is oversimplified when your goal is to improve body composition by adding lean mass and decreasing body fat. I am not disputing that every gram of protein yields four calories, every gram of carbohydrate



Photo by Garry Bartlett.

Testimonial

“When I turned pro in 1997, I competed at a body weight of only 211 pounds. I knew that in order to compete as a pro, I needed to be much bigger. I went to Venice, California, to find out what the secret was on how these guys were getting so huge. Believe it or not, I found out all the top pros were following the same basic principles of Macrobiotic Nutrition. Making these simple changes is all it took and now I compete at over 230 pounds. And the only thing I changed was my diet.”

—**Mike Morris**, IFBB Professional Bodybuilder

yields four calories, and every gram of fat yields nine calories. However, these oversimplified diet programs, written by so-called “diet experts,” recommend that you consume less food (calories) than you burn (expend) if you want to lose weight, and more calories than you expend if you want to gain weight. Wow—Isn’t that brilliant?! How many years of school did they have to go through to come up with such an earth-shattering revelation?

Well, it’s not that simple. Whether you are trying to gain weight or lose weight, these primitive diet plans will lead anyone to fail in obtaining his or her goals, especially bodybuilders and athletes. If your goal is to gain weight, you’ll gain weight all right, but it will be all fat! I know you’re not busting your ass in the gym lifting the heavy iron to put on fat. When a bodybuilder says he wants to get big, he’s talking about packing on rock-hard slabs of muscle, not about becoming a “big fatty.” Conversely, if you follow this kind of diet plan for weight loss, half of the weight you lose in the first month will be fat, but the other half will be hard-earned lean muscle. After the first month or so, your metabolism will slow down so much and you’ll be eating so little that most additional weight loss will be muscle. All that hard work you put in at the gym trying to build muscle will be wasted. Eventually, you’ll be left with a thin, soft, mushy body—what I call a “thin fat” body. Worse yet, you’ll get so disgusted with these diminished results that you’ll start eating poorly again out of frustration, and will blow up into a “big fatty.”

I’m not trying to sound obnoxious, but these diets, which so many people

follow, set them up for failure. And it annoys me to see people bust their asses in the gym, while they follow these diets, only to be disappointed.

I can't take full credit for many of the key scientific approaches used to develop Macrobiotic Nutrition. I must commend both Dr. Robert Atkins and Dr. Barry Sears for paving the way and opening the eyes of the medical community and dieticians, who are often blind to new science. I'm sure most of you are familiar with Dr. Atkins's high-protein/very low-carb "Atkins diet" and Barry Sears's "Zone diet." Both of these diets have considerable scientific merit and have been effectively used for weight reduction by millions of people. But what effect do they have on muscle mass and performance?

The fundamental theory behind the Atkins diet involves ingesting calories primarily from fat and protein. Your body resorts to using stored body fat as an energy source when carbohydrates are restricted from the diet. This process is known as ketosis. However, your body may switch to protein as another source of energy, instead of entering ketosis. When this happens, the liver converts protein into blood sugar via a process called gluconeogenesis. This is where the controversy for athletes begins to arise. How will your body discriminate between the protein in muscle or the body fat stored around it? While there is no doubt that the Atkins diet leads to weight loss in inactive people, it can lead to disaster for bodybuilders and athletes. A no-carbohydrate diet such as the Atkins diet leads to low energy levels and hinders performance. It also will lead to loss of muscle and ineffective recovery from exercise. Studies performed by the U.S. Olympic Committee have shown that an intake of anything less than 42 percent carbohydrates will hinder performance and energy. Carbohydrates are stored in the muscle as glycogen. Glycogen is what muscle uses for energy. In the absence of carbohydrates, precious protein is converted to glycogen instead of being utilized as a building block for muscle growth. This is the last thing a bodybuilder wants to have happen. To make matters even worse, gluconeogenesis, the process for converting protein to glycogen, is not nearly as efficient as the process of converting carbohydrates to glycogen, so less glycogen will be present in the muscle if carbohydrates are not available. Plus, glycogen makes muscles look full and hard, so a low carbohydrate intake is going to result in flat, soft-looking muscles.

The bottom line is that your nutritional requirements are different from your mom's, so get off her diet. A no-carb diet will hinder your ability to pack on muscle, and will leave you feeling weak and looking flat and small. You bust your ass in the gym to look good, so why waste your efforts by eating the wrong foods? Your body needs the right nutrition to get *big* and *ripped*!

Barry Sears's Zone diet is a little closer to the mark. Sears's approach closely examines blood sugar stabilization and insulin levels, much the way that Macrobiotic Nutrition does. In the Zone diet, 40 percent of your calories come

About Glycogen

Glycogen is a long chain, or polymer, of glucose molecules that is stored in the brain and liver, but mostly in muscle. Our bodies store carbohydrates as glycogen to be used for energy and to stabilize blood sugar during activity. As a rule of thumb, our bodies will store about 60 grams of carbohydrates in the liver and about 300 grams in the muscle, all of this as glycogen. Of course, the bigger you are and the more muscle you have, the more glycogen you can store. This is another benefit of having more muscle tissue than body fat. This is one of the reasons why a 250-pound athlete requires more carbohydrates and calories than a 160-pound athlete. Muscle glycogen is important to support your energy demands during workouts and to prevent the catabolic process gluconeogenesis.

from carbohydrates, 30 percent from fat, and 30 percent from protein. Sears's ratios are fine for a nonathlete and close to ideal for a regular athlete, but the 30 percent fat proportion tends to compromise the quality of the calories consumed. In order to eat that amount of fat, I feel you will inevitably raise your intake of saturated fat. Macrobiotic Nutrition puts more emphasis on the quality of the calories you eat. When it comes to outrageous muscle growth, the Zone just does not pack the punch you need to pack on the mass. It proves effective for endurance athletes, but falls short in protein demands for bodybuilders and strength athletes—for them, too little protein will hinder performance, recovery, and muscle growth. Macrobiotic's shift in caloric ratios is just what your body needs to keep it in an anabolic state, the state in which new muscle is built.

Success with Macrobiotic Nutrition relies not only on using the 45/35/20 lean-mass equation for your diet, but also in selecting the right sources and combinations of macronutrients (carbohydrates, protein, and fats) to maximize lean mass, strength, and overall performance. Macrobiotic Nutrition's key points include optimizing and regulating hormones, meal frequency, the thermogenic effect of food, and its *net effect* on metabolic efficiency. These scientifically advanced concepts will be discussed in easy-to-understand detail so you can apply them to your training program.

Because every athlete is at a different level of conditioning, Macrobiotic Nutrition outlines three different formulas for calculating your total Macrobiotic caloric needs. (See Chapter 7.) Use these formulas to calculate your own

calorie requirements based on your current condition and goal. A Macrobiotic caloric requirements table for each goal is included in Appendix A to make following the program even easier. These tables give the amount of carbohydrates, proteins, fats, and total calories athletes should consume depending on their level of conditioning. You will be able to customize a Macrobiotic Nutrition diet specifically for you, taking into account your current body weight and body composition, occupation, lifestyle, workout regimen, and most important, your *goal!*

GOALS

- **Gain Lean Muscle/Lose Body Fat**
- **Gain Lean Muscle/Maintain Current Body Fat**
- **Gain Lean Muscle/Gain Body Fat**

The key to any long-term nutrition program is consistency, but don't worry about this being a boring diet. Chapter 10 presents some really tasty Macrobiotic meals for you. From quick meals to exotic specialty meals, they are all delicious and they are all Macrobiotic! The longer you follow Macrobiotic Nutrition, the bigger, leaner, stronger, and healthier you will become. You won't want to stop, once you see the changes and incredible results. That is why Macrobiotic Nutrition is called the long-term, permanent solution.



Photo by Frank DeJianne.

Testimonial

"I'm known by Strongman competitors as an overachiever. I am not the most gifted athlete, so I have to rely on a superior training and nutrition regimen to compensate. My training sessions are so intense that I burn through training partners every few months. Macrobiotic Nutrition fuels my body with the nutrients I need for muscle growth, strength, and recovery."

—**Steve Kirit**, USA Strongman Champion
in 2002 and 2003

Testimonial

"I've been involved in power-lifting, Strongman, and strength coaching for over fifteen years. And I have never seen a program that compares to Macrobiotic Nutrition for strength training athletes. An athlete's nutrition requirements are far greater than the average person's and to optimize lean mass requires even more precise nutrition. Macrobiotic Nutrition really provides the best sources and balance of nutrients for athletes looking to gain size, strength, and improve performance."



Photo by Frank DeJianne.

optimize lean mass requires even more precise nutrition. Macrobiotic Nutrition really provides the best sources and balance of nutrients for athletes looking to gain size, strength, and improve performance."

—**Mark Philippi**, Former USA Strongman Champion and UNLV Strength Coach

A DIET FOR BODYBUILDERS AND POWER ATHLETES: BUILD MUSCLE, BURN FAT, AND INCREASE STRENGTH AND PERFORMANCE

The primary goal of every bodybuilder and athlete is to reach his or her full genetic potential. This can be achieved only through proper diet and training. Diet is even more critical than training, because diet influences your training capacity. A compromised diet means compromised training, and ultimately compromised "poor" results. If you want to be the best at your sport, you need to eat for your sport. Macrobiotic Nutrition is the bodybuilder's and power athlete's diet; it is designed to fuel muscles with the necessary nutrients to supply energy for heavy weight training, increased muscle building, and optimal fat burning. The goals here are simple: build muscle, increase strength, and control body fat.

LEAN-MASS EQUATION

Let's start with a simple example to illustrate Macrobiotic Nutrition's fundamental principle, "All calories are *not* created equal," before we even get to the science to prove it. Which breakfast do you think your body will respond to best for the goal of building muscle and controlling body fat?

Dieticians would argue that all four breakfasts yield approximately the

BREAKFAST #1 (MACROBOLIC)
Eggs and Oatmeal with Strawberries

Food Item	Quantity	Energy (kcal)	Carbohydrate (g)	Protein (g)	Fat (g)
Oatmeal	1½ cup cooked	240	45	6	2
Egg whites	8 large eggs	85	—	27	—
Strawberries, whole	1¼ cup	60	15	—	—
Whole eggs	2 large	150	—	14	10
	TOTALS	535	60	47	12
			45%	35%	20%

BREAKFAST #2 (BLUE COLLAR)
Bacon, Egg, and Cheese Sandwich

Food Item	Quantity	Energy (kcal)	Carbohydrate (g)	Protein (g)	Fat (g)
Bagel, plain	1	150	30	6	—
Whole egg	1 large	75	—	7	5
American cheese	1 ounce	100	—	7	8
Bacon	3 slices	100	—	7	8
Orange juice	1 cup	120	30	—	—
	TOTALS	545	60	27	21
			45%	20%	35%

BREAKFAST #3 (LOW CARB)
Bacon, Egg, and Cheese Omelet

Food Item	Quantity	Energy (kcal)	Carbohydrates (g)	Protein (g)	Fat (g)
Butter	1 tsp	45	—	—	5
Whole eggs	4 large	300	2.44	21	15
Cheddar cheese	1 ounce	100	0.36	7	8
Bacon	3 slices	100	—	7	8
	TOTALS	545	2.80	35	36
			2%	30%	68%

BREAKFAST #4 (WALL STREET)					
Corn Muffin and Orange Juice					
Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)
Corn muffin	1 medium	335	57.52	6.67	9.49
Butter	2 tsp	90	—	—	10.00
Orange juice	1 cup	120	30.00	—	—
TOTALS		545	87.52	6.67	19.49
			63%	5%	32%

same amount of calories, so they would all have the same impact. The low-carb community would argue that breakfast #3 is going to facilitate fat burning best. Common sense and knowledge of Macrobiotic Nutrition tell you that the balanced breakfast #1 would be best, and this book is going to prove it to you.

The cornerstone of Macrobiotic Nutrition is the 45/35/20 lean-mass equation. Breakfast #1 is a good example of a Macrobiotic meal. Macrobiotic Nutrition derives its calories from select “nutrient-dense” sources, in a ratio of 45 percent carbohydrates, 35 percent proteins, and 20 percent fats. This ratio was determined through extensive research to be the optimal macronutrient profile for creating the most favorable hormonal, metabolic, and thermogenic effects from food.

The ratio and selection of macronutrients (carbohydrates, proteins, and fats) are critical when composing meals. Each macronutrient influences different hormones and chemical processes. Optimal hormonal homeostasis (“balance”) can be achieved only with the right ratios and sources of macronutrients in each meal. Macrobiotic Nutrition’s 45/35/20 lean-mass equation creates the ideal hormonal and metabolic response for building muscle and burning body fat. In the chapters that follow, I will examine how each of the macronutrients influence chemical processes within the body and how they react with one another.

CHAPTER 2

Carbohydrates— Don't Count Them Out

SINCE MANY OF YOU MAY BE STRUGGLING to believe you can get lean with 45 percent of your calories coming from carbohydrates, they will be the first macronutrient discussed. You'll soon find out how it is possible to lose body fat and consume 45 percent of your calories from carbohydrates. But let's first discuss how important carbohydrates are for strength, muscle growth, and performance.

Carbohydrates are the most efficient nutritional source for the body's energy requirements, because carbohydrates are more easily converted to glucose than proteins and fats. Glucose is used by every cell in the body as fuel for energy. Some of the glucose is carried around in your bloodstream to supply your brain and other organs, but most of it is stored as glycogen in the liver and skeletal muscle. When blood sugar levels are low, this glycogen serves as a reserve to replenish blood sugar to meet your body's energy needs. Another very important role of carbohydrates is their "protein-sparing" effect. Carbohydrates protect your protein from being converted to glucose when blood sugar and glycogen are low.

Your body burns more energy when you exercise, so its need for blood glucose increases. Your body taps into the muscles' glycogen reserves to meet these energy demands, so it is critical that bodybuilders consume adequate amounts of carbohydrates to support their energy demands. A 2002 review on diet and anaerobic exercise (weightlifting) in *Strength and Conditioning Journal* concluded that diets containing less than 42 percent carbohydrates do not meet the energy demands or provide adequate glycogen stores for bodybuilders, given their intense workouts. The physical demands bodybuilders place on their bodies creates a greater need for the kind of heavy-duty nutrition that repairs muscle fiber and sustains growth.



Photo by Clark Jackson.

Testimonial

“Most bodybuilders make the mistake of restricting carbs from their diet. Carbs are essential for glycogen replenishment and sparing aminos in muscle tissue. Another common mistake bodybuilders make is avoiding soy as a protein supplement. Soy is one of my favorite protein sources because of its high bioavailability. And unlike meat, milk, and whey proteins, which are acidic, soy is less acidic and easier on your stomach. When you are taking in large amounts of protein, this is very important. Macrobiotic Nutrition stresses the importance of low-glycemic carbs and utilizing a variety of protein sources, which are the backbone of my nutrition program.”

—**Gary Strydom**, *IFBB Professional Bodybuilder*

THE LOW-CARB CONUNDRUM: BURNING FAT AT THE EXPENSE OF MUSCLE

The huge amount of media attention on low-carb diets may have led you to believe that the low-carb diet was the route to getting a lean, muscular physique. Unfortunately, you probably found out the hard way that the low-carb diet you tried didn't give you the results you wanted. Rather than becoming big and hard with full dense muscles, you feel flat, small, and weak!

The fundamental theory behind low-carb diets is that when carbohydrates, the body's preferred source of energy are restricted, the body resorts to using stored body fat as an energy source. In the process known as ketosis, the body breaks down triglycerides for use as a source of energy. Ketosis is not nearly as efficient as using carbohydrates for energy or glycogen replenishment. This lack of efficiency in the absence of carbohydrates can result in the body's also using protein (which is made up of amino acids linked together) from food and muscle tissue for energy. Certain amino acids in the protein you eat and in the protein that makes up your muscle tissues can be converted to glucose. As mentioned earlier, this process is called gluconeogenesis. Amino acids are the building blocks of protein and muscle tissue. Two such acids used for gluconeogenesis are glutamine and alanine. Later in the book, you'll learn more about these amino acids and protein's importance. We already know that body-

builders need adequate protein to support training, muscle growth, and recovery, so having your body cannibalize protein for energy from the foods you eat, or even worse, from your “hard-earned muscle,” is a catastrophe for any bodybuilder.

Carbohydrates are too important a nutrient for bodybuilders and athletes to completely restrict from their diets. You will never make the gains in size and strength you are looking for on a low-carb diet. True, low-carb diets can be effective in reducing body fat, but they can also burn off precious muscle. That's not what you want to achieve from your diet. Remember the goals of Macrobiotic Nutrition are to build muscle, increase strength, and control body fat.

ALL CALORIES ARE NOT CREATED EQUAL!

Macrobiotic Nutrition's fundamental principle “All calories are not created equal” applies to all the macronutrients: carbohydrates, proteins, and fats. Mac-




Photo supplied by Dave Hawk.

Before

up to a show. During my layoff I really let my physique go and put on a lot of body fat. My first instinct was to cut carbs; however, Gerard convinced me that if I wanted to be bigger and better than ever, I had to change my diet. Amazingly, using the Macrobiotic Nutrition regimen along with my Return to Dominance supplementation program, I competed bigger, harder, and fuller at the age of 40 than I did in my last pro show at the age of 28.”

—**Dave Hawk**, IFBB Professional Bodybuilder

Testimonial

“Coming back to professional bodybuilding after a twelve year layoff and competing in the 2002 Masters Mr. Olympia was a dream. Thankfully, I had the opportunity to work with Gerard Dente to help structure my nutrition and some of my supplementation program. In my earlier competition days I followed a high-protein/low-carb diet while preparing for contests. I was usually cut but always lost

a lot of size leading




Photo by Sumio Yamaguchi.

After

robolic Nutrition places great emphasis on choosing your carbohydrate sources wisely. While all carbohydrates yield four calories per gram, their impact on the body can greatly differ.

Carbohydrates can be broken down into three general categories: monosaccharides, disaccharides, and polysaccharides. Monosaccharides and disaccharides are commonly referred to as sugars or simple carbohydrates, while polysaccharides are called complex carbohydrates.

Monosaccharides are carbohydrates that have one sugar molecule. Common sources include glucose, fructose, sorbitol, galactose, mannitol, and mannose.

Disaccharides are carbohydrates with two sugar molecules. Common sources include sucrose and lactose.

Polysaccharides are carbohydrates with three or more sugar molecules. Sources include dextrin, cellulose, and starches.

Another kind of carbohydrate is fiber, which is composed mainly of undigestible polysaccharides. Fiber plays an important role in the next topic of discussion, the glycemic index (GI). Though the classifications of monosaccharides, disaccharides, polysaccharides, and fiber help differentiate carbohydrates, Macrobiotic Nutrition puts great emphasis on the glycemic index when choosing carbohydrate sources.

Table 2.1 on the next page lists the carbohydrate sources I recommend and their caloric, carbohydrate, protein, and fat content.

What Is the Glycemic Index?

The glycemic index (GI) was developed in 1981 as a way to classify carbohydrates. As defined by Jennie Brand-Miller, Associate Professor of Biochemistry at the Human Nutrition Unit of Sydney University, Australia, the glycemic index of a food is a measure of the power of the carbohydrate content in a specific food to raise blood glucose levels after being eaten. The glycemic index is a ranking of carbohydrates based on their immediate effect on blood glucose (blood sugar) levels. It compares carbohydrate foods gram for gram. Carbohydrates that break down quickly during digestion have the highest glycemic indexes and a blood glucose response that is fast and high. Carbohydrates that break down slowly, releasing glucose gradually into the bloodstream, have low glycemic indexes.

The GI of a carbohydrate is determined by measuring blood sugar levels after ingesting 50 grams of the carbohydrate in a fasting individual. Foods that measure a GI of 70 to 160 are considered high GI foods, foods that measure 56 to 69 are considered medium GI, and foods that measure 55 or less are low GI. The GI of a carbohydrate can be influenced by a number of factors: the fiber content, the ripeness, methods of cooking and processing, the types of pre-

TABLE 2.1 CARBOHYDRATE SOURCES RECOMMENDED FOR MACROBOLIC NUTRITION

Source	Portion	Calories	Protein	Fat	Carbohydrates
Grains					
Couscous	1 cup	175	6.0	—	36.0
Oats	½ cup	155	6.0	2.5	27.0
Rice, brown cooked	1 cup	216	5.0	2.0	45.0
Rice, white cooked	1 cup	200	4.0	0.5	43.0
Pasta (semolina)	2 oz	200	7.0	1.0	41.0
Wheat bread	1 slice	70	3.0	1.0	13.0
Vegetables					
Potato, baked with skin	1 med	160	4.0	—	37.0
Sweet potato	1 cup	206	3.0	1.0	49.0
Fruit					
Apple with skin	1 med	80	—	—	21.0
Banana, just ripe	1 med	110	1.0	0.5	28.0
Blueberries, fresh or frozen	1 cup	80	1.0	0.5	20.5
Cantaloupe	1 cup	55	1.0	—	13.0
Grapefruit	½ med	40	0.7	—	10.0
Orange, Florida	1 med	65	1.0	—	16.0
Pear with skin	1 med	100	0.6	0.6	25.0
Raspberries, fresh or frozen	1 cup	60	—	—	15.0
Strawberries, fresh or frozen	1 cup	50	—	—	12.0

servatives used, and the types and amounts of macronutrients (other carbohydrates, proteins, fats) and micronutrients (vitamins and minerals) ingested with the carbohydrate.

What Is the Significance of the Glycemic Index in Macrobiotic Nutrition?

The reason it is so important to measure a carbohydrate's impact on blood sugar is because of its influence on insulin production. Insulin is a hormone released by the pancreas in response to changes in blood sugar levels. The faster carbohydrates are digested and converted to glucose, the more rapidly insulin is produced to stabilize blood sugar levels, using the glycemic index as a measure.

The GI and Insulin

- Low GI means a smaller rise in blood glucose levels after meals.
- Low GI diets can help people lose body fat and increase lean muscle mass.
- Low GI diets can improve the body's sensitivity to insulin.
- Low GI diets can improve the ratio of insulin to glucagon.
- Low GI foods can help replenish carbohydrates stores after exercise.
- Low GI can prolong muscle endurance and energy levels.
- Low GI can improve diabetes control.
- Low GI foods keep you feeling fuller for longer.

Using the glycemic index as a measure, carbohydrates that are digested faster have a higher GI because they cause a greater increase in blood glucose and insulin levels. A bodybuilder wants to avoid high insulin levels, because insulin suppresses fat utilization and promotes fat storage. The rapid increase in blood sugar caused by high GI carbohydrates can also have a negative impact on performance.

High GI carbohydrates also elevate free fatty acids in the blood, further promoting increased body fat. They have also been shown to predispose the development of type 2 diabetes in insulin-resistant individuals. In response to fast-rising blood sugar levels, your pancreas releases large amounts of insulin in an effort to compensate and stabilize the blood sugar. Often, the overproduction of insulin can result in low blood sugar levels or temporary hypoglycemia, a condition in which blood sugar levels drop below normal range. Hypoglycemia can cause fatigue, anxiety, perspiration, light-headedness ("delirium"), and, in severe cases, coma.

These fluctuations in blood sugar levels from very high to very low can hinder your ability to train and perform at maximum capacity. Nothing will zap your energy, strength, and performance like a bout of hypoglycemia during your workout. Temporary hypoglycemia can usually be corrected by consuming more carbohydrates, but it is a condition you want to avoid. If you ever get to this point of temporary hypoglycemia, you will know it. You'll feel super weak and disoriented, and you will usually break out in a cold sweat. If you experience these symptoms, grab yourself some carbohydrates and scoff them down.

Macrobiotic Nutrition places great emphasis on consuming carbohydrates with low to moderate GI for these reasons. Low-to-moderate GI carbohydrates are digested more slowly, providing a gradual, steady supply of blood sugar. This slow, steady supply of blood sugar is critical for peak energy and performance. It also maintains muscle glycogen stores and helps to regulate two very important hormones, insulin and glucagon, for optimum muscle growth and fat loss.

Insulin and glucagon are both influenced by blood sugar levels. High-glycemic carbohydrates such as sugar cause insulin levels to be high and glucagon levels to be low. Since insulin promotes the increase in body fat and glucagon mobilizes and burns body fat, this obviously isn't the metabolism we want. Lower glycemic carbohydrates, especially when they are consumed with protein, shift the levels of these two hormones, lowering insulin and slightly raising glucagon. This is the correct "hormonal profile" for effective fat burning.

Table 2.2 on the next page lists the GI of common carbohydrate sources, broken down into high, moderate, and low GI classes. Try to limit your selections to moderate and low whenever possible. The items included in these tables are primarily foods considered to be "carbohydrate sources," which means most of their caloric content comes from the carbohydrates and they provide proportionately very little protein or fat. These listed carbohydrate sources would need to be consumed with other food sources higher in protein and/or fat to provide the Macrobiotic Nutrition 45/35/20 ratio. When these carbohydrates are consumed with these other foods as part of a Macrobiotic meal, they will yield lower glycemic values.

Keep this in mind when looking up the glycemic index of specific "junk foods." Their glycemic index may seem suitable for Macrobiotic Nutrition from a GI standpoint, but they may fall short in many other areas of Macrobiotic Nutrition, with inadequate protein content and high levels of saturated fat.

TABLE 2.2. GLYCEMIC INDEXES OF COMMON CARBOHYDRATE FOODS

HIGH GLYCEMIC			
Food Item	GI	Food Item	GI
White bread	70	English muffin™	77
Pop Tarts™	70	Cornflakes™	77
Golden Grahams™	71	Corn Pops™	80
Bagel, white	72	Special K™	84
Wonder Bread™	73	Rice cakes, plain	94
Cheerios™	74	Glucose	100
Total Cereal™	76	Maltodextrin	107
MODERATE GLYCEMIC			
Food Item	GI	Food Item	GI
Cranberry juice	56	Banana (just ripe)	62
Baked potato, russet	56	Long grain rice, white	64
White rice, boiled	56	Spaghetti, durum wheat	64
Sourdough rye bread	57	Cantaloupe, raw	65
Pita bread, whole wheat	57	Wholemeal rye bread	66
Blueberry muffin	59	Cream of Wheat™	66
Sweet corn	59	Croissant	67
Bran muffin	60	Grapenuts™	67
Couscous	61	Shredded Wheat™	67
Just Right Just Grains™	62	Cornmeal	68
LOW GLYCEMIC			
Food Item	GI	Food Item	GI
Cashew nuts	22	Banana, slightly unripe	42
Cherries, raw	22	Spaghetti, whole meal	42
Grapefruit, raw	25	Spaghetti, white	44
Barley	27	Pumpernickel, whole grain	46
Lentils, boiled	28	Spaghetti, semolina	46
Peach, raw	28	Sweet potato	48
Milk (skim)	32	Orange	48
Pear, raw	33	Brown rice, steamed	50

LOW GLYCEMIC <i>(continued)</i>			
Milk (whole)	36	Durum wheat	50
Yogurt	36	Oat bran, raw	50
Pinto beans, boiled	39	100% whole-grain bread	51
Apple, raw	40	Kidney beans, canned	52
Strawberries, fresh	40	Spelt wheat-flour multigrain bread	54
Chickpeas	41	Oatmeal	54

GI Table Explanation

When looking up GI values, you may run into what seems to be conflict between this GI table and that of other established GI tables. For example, you may type in “GI value for white rice” on the Internet and you may find multiple values for the same food. This may seem very confusing, but there is an accepted and reasonable explanation for this.

Different methods of processing used by particular manufacturers can lead to significant differences in the rate of carbohydrate digestion by the human body. (The degree of starch gelatinization used in the process may differ, for example.) Also, there may be botanical differences in the type of food being tested. Rice can be from anywhere in the world, and one “white rice” may contain more amylose than another. Amylose is digested more slowly than amylopectin, which is another starch found in other “white rice.” Another difference may even be related to the methods that the research team used to test the GI of a food. GI values can vary greatly due to the amount of time the researchers used to conduct the test or different portion sizes of the test foods, or even due to the source of the drawn blood (venous versus capillary).

So use your better judgment, but don’t be fanatical. Macrobolic Nutrition has devoted valuable time and resources to creating what I believe is the most accurate and relevant GI table available. This GI table lists the foods that you will most commonly eat on your Macrobolic Nutrition program. So, you can use this guide with a great degree of certainty.

INSULIN’S POWERFUL EFFECTS ON BODY COMPOSITION

Insulin has both anabolic (muscle-building) and hyperlipidemic (fat-storage) properties. The key to building lean mass is to control insulin to promote muscle growth and burn body fat. Is insulin needed for muscle growth or does it just store fat? How does insulin affect performance? What are some of the other negative effects of insulin? What is insulin resistance? These questions are addressed in the following sections.

Fiber—The Forgotten Carbohydrate

Fiber plays a critical role in Macrobiotic Nutrition. Fiber is the most recognized of all carbohydrates in terms of being important for disease prevention and general health. Fiber is classified into two different types: soluble and insoluble. Insoluble fiber aids in intestinal cleansing and slows down digestion through the gastrointestinal tract, an important concern for a bodybuilder, since it allows for more thorough nutrient absorption.

Soluble fiber, on the other hand, is an invaluable factor in Macrobiotic Nutrition's theory that a calorie is not just a calorie. According to a study published by M. Chandalia, et al., in *The New England Journal of Medicine* (2000), a diet containing 50 grams of fiber (25 soluble, 25 insoluble) can:

- lower cholesterol
- improve glycemic control
- decrease hyperinsulinemia (the overproduction of insulin by the pancreas)

These findings are very interesting to a bodybuilder wishing to control the glycemic index of meals. By ingesting soluble fiber at mealtime, we can prevent spikes in blood sugar levels and reduce the overall secretion of insulin.

Natural oat fiber reduced blood glucose and insulin levels in a study by J.T. Braatan, et al., discussed in the *American Journal of Clinical Nutrition* (1991). The soluble fiber used in this study was beta-glucan. Oats and barley are good natural sources of beta-glucan. In a 1994 study published in *Diabetic Medicine* by Braatan, et al., patients who were fed wheat farina with oat bran showed a marked decrease in post-prandial (after a meal) blood glucose levels and lower overall insulin secretion.

These studies support the great importance of fiber in Macrobiotic Nutrition. Its impacts on maintaining lower blood glucose and lower insulin levels are conducive to increased fat burning as well as to decreased cortisol and increased growth hormone (GH) levels. Cortisol and GH are two important hormones for bodybuilders, and are described in detail later in this chapter. These are all the things your body needs to pack on the lean mass you want.

Insulin: Is it Anabolic or Hyperlipidemic?

One of the primary concerns that Macrobiotic Nutrition's ratio of calories and low-glycemic carbohydrates addresses is the effect blood sugar levels have on insulin. Insulin is a very powerful hormone, and it needs to be regulated through proper nutrition. Not everything about insulin is bad. In fact, insulin is very anabolic! Insulin increases the transport of glucose, amino acids, and other nutrients into muscle, so it has a very anabolic effect. The increased transport of glucose to muscle serves to increase glycogen stores, which are important for muscle performance. The increased shuttling of amino acids results in increased protein synthesis and nitrogen retention, both essential for muscle growth. So far, it sounds like we should want to load up on insulin, doesn't it? But that's not the case. Elevated insulin can be very hyperlipidemic, which means it can increase body fat. One way insulin increases body fat is by assisting in the formation of glycerol and fatty acids into triglycerides, which are stored as body fat. Elevated insulin not only promotes the storage of triglycerides, it also inhibits the body from breaking them down into burnable fatty acids. So, elevated insulin promotes the storage of body fat, and also inhibits the body from breaking down body fat into usable energy. It's easy to see that elevated insulin levels will make you fat if they are not managed properly by controlling blood sugar.

Controlling Insulin for Optimum Performance

Insulin release must be managed carefully to take advantage of insulin's powerful anabolic effects and to avoid its fat-storing effects. Macrobiotic Nutrition will elicit the ideal controlled blood sugar level to keep insulin in the desirable "anabolic zone." The slow, steady release of blood sugar from a Macrobiotic meal causes the pancreas to gradually release insulin into the bloodstream to manage blood sugar. This is what you want, to get the anabolic effects of insulin while avoiding the storage of body fat.

Figure 2.1 illustrates the effects on blood sugar of a low-glycemic meal versus a high-glycemic meal. Blood sugar levels were tested at 15, 30, 45, 60, 90, and 120 minutes after the meals. As you can see, the Macrobiotic meal provides a much more gradual and steady blood sugar level, while the high-glycemic meal causes a rapid increase in blood sugar and then a quick drop off (even below baseline).

The steady blood sugar levels resulting from the Macrobiotic meal causes a gradual release of insulin from the pancreas. This insulin shuttles the glucose (sugar) provided by the carbohydrates to your muscles to be stored as glycogen. The insulin also shuttles the amino acids into the muscle, a process that stimulates muscle growth by keeping you in the positive nitrogen balance needed for




Figure 2.1. Blood Sugar Values of Macrobiotic versus High-Glycemic Meals

muscle development. These are the effects we want from a meal, and the reason why I call it the “anabolic zone.”

On the other hand, the rapid increase in blood sugar caused by the high-glycemic meal causes your pancreas to produce a lot of insulin to try to quickly lower the blood sugar. These highly elevated insulin levels put you in the “hyperlipidemia zone” and promote the storage of body fat. But that isn’t the only problem such insulin levels create.

The Negative Effects of Insulin

Hypoglycemia

The rapid release of insulin to compensate for high blood sugar actually results in low blood sugar, or hypoglycemia, as mentioned earlier. Some of the symptoms of hypoglycemia include low energy levels, dizziness, moodiness, light-headedness (“delirium”), shakiness, and sweating followed by extreme hunger. Bodybuilders refer to this state as crashing, and this is obviously a condition we want to avoid. I don’t know if you have ever experienced this, but it is definitely not a good feeling. Depending on how low your blood sugar drops, you can feel like you are going to pass out. In the most severe cases you can pass out and wind up in a coma. The remedy for this condition is to slam a bunch of carbohydrates in the hope of bringing your blood sugar back up. This is only a Band-

Aid on the problem, however, because your body will now release more insulin to compensate for all the carbohydrates you just ate, and in a little while, you will be right back where you started.

Your body will never get rid of stored body fat if your meals continually do this to you. Regular bouts with hypoglycemia resulting from large consumption of simple, high-glycemic carbohydrates can eventually lead to insulin-resistance syndrome, or to type 2 diabetes, which comes with a long list of permanent health issues. Dr. Joseph Mercola's paper on this topic, "Reduce Grains and Sugar to Lose Weight and Improve Health," recommends consuming 40 to 45 percent of your calories from carbohydrates to prevent this "crash." This is exactly the level of carbohydrate intake used as the basis for Macrobiotic Nutrition. Using low-glycemic carbohydrates in conjunction with protein and fat in meals prevents the spikes in insulin and rapid drops in blood sugar.

Insulin Resistance

If there ever was a valid medical argument to support Macrobiotic Nutrition, insulin-resistance syndrome (IRS) is it—the reason you should take advantage of the caloric ratios we advocate. (Syndrome X is a common name for insulin resistance.) I choose to speak about it here as it is a primary concern of a high-carbohydrate diet.

With IRS, muscle cells and tissues in the body develop a reduced sensitivity to insulin's actions of delivering blood glucose to them to use as an energy source. As a result, the body senses a high blood sugar level, which signals the



Photo supplied by Gene Rychlak.

Testimonial

"Macrobiotic Nutrition is incredible! It allowed me to increase my bench from 750 pounds to a world record 900 pounds in less than one year. And I actually lost body fat in the process. Okay, so maybe I don't have six-pack abs nor do I want or need them. But the point is that Macrobiotic Nutrition helped me add 150 pounds to my bench press and I did lose body fat while doing it. Nobody ever thought I could bench 900 pounds—I proved them wrong!"

—**Gene Rychlak**, *World Record 900-Pound Bench Press*

pancreas to secrete more insulin to lower it. Eventually a person may develop type 2, or noninsulin-dependent diabetes, because the pancreas just can't make enough insulin to handle the high blood sugar levels. Along with the diabetes come high blood pressure, high triglycerides, and high cholesterol.

Sounds like some scary stuff, doesn't it? I think so. If you think about it, this is all due to your blood sugar's being too high all the time, and your body's secreting too much insulin in an attempt to control it. This is exactly what happens when you eat too much sugar and high-glycemic carbohydrates. Macrobiotic Nutrition takes advantage of a diet that is moderate in carbohydrates (only 45 percent, not like the typical 80 percent most Americans eat) and generous in the muscle-building protein and fat your highly trained body will crave. As far as restoring insulin sensitivity and getting your body back on track, Macrobiotic Nutrition has the prescription for success: frequent meals (every two and a half to three hours) with each meal containing 45 percent low-glycemic carbohydrates, 35 percent protein, and 20 percent fat; a diet high in fiber; and regular exercise. Not only does Macrobiotic Nutrition address the demands your body experiences through athletics, it also gives you the plan you need to be healthy and fit for your lifetime.

MACROBOLIC NUTRITION'S EFFECT ON KEY HORMONES: INSULIN, GLUCAGON, GROWTH HORMONE (GH), AND CORTISOL

I've already established the importance of managing insulin due to its very anabolic (muscle-building) and hyperlipidemic (fat-storing) properties, as well as its influence on health. However, insulin has an even bigger impact on your ability to build muscle and burn fat through its influence on other key hormones: glucagon, growth hormone (GH), and cortisol. All of these hormones can be manipulated through Macrobiotic Nutrition to achieve the ideal hormonal profile to build a "rock-hard" physique.

Here is how it works. Macrobiotic Nutrition keeps blood sugar steady while supplying amino acids and glycogen to muscle tissue. Keeping blood sugar under control also keeps insulin levels low. When insulin is low and amino acid levels are high, the pancreas releases a hormone called glucagon. Glucagon is responsible for mobilizing stored body fat and utilizing this fat as an energy source. Figure 2.2 on the next page shows the intimate relationship between insulin, glucagon, and blood sugar levels.

Low insulin levels also increase GH levels. Every bodybuilder is familiar with GH. If you aren't so sure about what GH is or does, let's just say its name says it all—or I should say almost all. You see, not only will GH make you big like its name implies, it will also burn fat and get you shredded. GH stimulates




Figure 2.2. Normalization of Blood Glucose in the Body

muscle growth by increasing protein synthesis and nitrogen retention, so you'll get even better utilization of your quality protein sources and create an even more anabolic environment. What most people don't know is that GH also has lipolytic (fat-burning) properties. Elevated GH levels have been shown to increase fat oxidation and mobilization, resulting in a reduction in body fat stores. You will optimize glucagon and GH levels and further enhance muscle building and fat burning by keeping your insulin under control.

In addition to its importance to glucagon and GH, insulin also has a direct effect on another hormone, cortisol. Cortisol is extremely catabolic and will actually break down and eat away your muscle tissue. High levels can send your training and growth into a declining tailspin. As a bodybuilder, the last thing you want is elevated cortisol. Keeping cortisol levels under control is no easy task, however, because the two things that shoot cortisol levels through the roof are intense training and insulin. If you want to get big you have no choice but to train hard, so pussyfooting around the gym and training like a wimp isn't the solution to keeping cortisol down. The good news is that Macrobiotic Nutrition will take care of it for you. The first way Macrobiotic Nutrition helps keep cortisol under control is by keeping insulin levels down. Cortisol and insulin

have a direct effect on each other, and elevations in insulin induce elevations in cortisol. You can see that avoiding cortisol's catabolic effect on your muscle tissue is another good reason to keep insulin levels low and under control.

The other way Macrobiotic Nutrition helps keep cortisol in check is by supplying large amounts of glutamine and amino acids from its sources of protein. Your body produces cortisol in response to physical and emotional stress, so you are vulnerable to its catabolic effects after an intense workout. That is why it is supercritical to supply your body with a Macrobiotic meal that supplies high concentrations of amino acids, especially glutamine, after your workout. Glutamine has been shown to neutralize the catabolic effects of cortisol. Supplementation with additional L-glutamine and phosphatidylserine post-workout is also a good idea to help suppress cortisol.

Macrobiotic Nutrition primes your hormonal environment for serious muscle building and fat burning. The 45/35/20 ratio of nutrients provides the optimal hormone profile:

- Low and controlled insulin release
- Slightly elevated glucagon levels
- Increased growth hormone
- Low cortisol levels

TABLE 2.3 MACROBIOTIC NUTRITION'S HORMONAL EFFECTS ON BODY COMPOSITION

⇒ Insulin	Body Fat ↓	Muscle Mass ↑
↑ Glucagon	Body Fat ↓	Muscle Mass ⇔
↑ GHG (IGF-1)	Body Fat ↓	Muscle Mass ↑
↓ Cortisol	Body Fat ↓	Muscle Mass ↑
↑ INCREASE	⇔ STABLE	↓ DECREASE

Table 2.3, although it might be new to you, is no secret to professional bodybuilders. They've been optimizing and manipulating the levels of key hormones for years to get in critical contest shape.

Insulin's impact on glucagon, growth hormone, and cortisol makes a very strong case for Macrobiotic Nutrition's emphasis on controlling blood sugar. Optimizing the levels of these hormones is the only way you're ever going to be able to pack on the mass and keep your body fat in check. If you're wondering about testosterone, don't worry. I didn't forget about it. It will be covered in the discussion on fats in Chapter 4.

CHAPTER 3

Protein— Choosing the Right Building Blocks

PROTEIN IS ONE OF THE KEY NUTRIENTS your body needs to support muscle growth and repair, to form neurotransmitters in the brain, and to create hormones. Protein serves as the source for the amino acids your body craves after you train. Since growth occurs after you lift, it only makes sense that you give your body a constant supply of protein to make sure there is a constant supply of amino acids. The trick for a bodybuilder is making sure to consume different types of protein. Different proteins are made up of different concentrations of amino acids, but some protein sources lack certain amino acids. With regard to muscle growth, the value of a particular protein depends on the proportion or number of essential amino acids (the ones your body cannot synthesize) it contains. Most animal-based proteins are high in these particular amino acids, while vegetable source proteins tend to lack at least one of the essential amino acids.

As a rule, athletes and individuals who work out have higher demands for protein than do sedentary individuals. After a workout, muscle growth is stimulated by insulin's increased transport of amino acids into the muscle tissue. Increased amino acid transport increases the amount of nitrogen present, a critical factor for muscle growth. Additionally, certain amino acids actually help increase the release of anabolic hormones, like growth hormone. This is the metabolic environment your body needs for you to pack on the mass. Current research on strength athletes shows that you need to ingest about 1 to 1.5 grams of protein per pound of body weight daily.

Using this research as a guide for protein requirements, a 200-pound strength athlete or bodybuilder would need between 200 and 300 grams of protein per day. Macrobiotic Nutrition provides even more precise requirements, because it takes into account other important criteria such as your goal, your lifestyle (activity level), and the length of your workout sessions. As you

will see from the Macrobiotic Nutrition caloric calculator in Chapter 7, the same 200-pound athlete would have the protein requirements set forth in Table 3.1.

TABLE 3.1. 200-POUND ATHLETE CALORIC/PROTEIN REQUIREMENTS

Activity Level	Workout Time	Total Calories	Protein Requirements (35% of Total Calories)
GOAL: GAIN MUSCLE/LOSE BODY FAT			
Sedentary	60 minutes	2,600	228
Moderate	60 minutes	3,000	263
Active	60 minutes	4,000	350
Sedentary	90 minutes	2,900	254
Moderate	90 minutes	3,300	289
Active	90 minutes	4,300	376
GOAL: GAIN MUSCLE/MAINTAIN BODY FAT			
Sedentary	60 minutes	3,200	280
Moderate	60 minutes	3,600	315
Active	60 minutes	4,600	403
Sedentary	90 minutes	3,500	306
Moderate	90 minutes	3,900	341
Active	90 minutes	4,900	429
GOAL: GAIN MUSCLE/GAIN BODY FAT			
Sedentary	60 minutes	3,800	333
Moderate	60 minutes	4,200	368
Active	60 minutes	5,200	455
Sedentary	90 minutes	4,100	359
Moderate	90 minutes	4,500	394
Active	90 minutes	5,500	481

Training intensity is a key component in determining the amount of new muscle growth you can achieve. Workout intensity needs to be high enough to challenge the body's growth, but it can't be so intense that it causes an overload and shuts down the body's metabolic response for new tissue growth. No amount of protein or nutrition can compensate for overtraining. Most people, myself included, are guilty of overtraining. Your determination to get big

can work against you if you are not careful. Once you understand Macrobiotic Nutrition and realize how these nutritional changes are going to unlock your growth potential, your enthusiasm to get to the gym and train your ass off is going to be at an all-time high. Be careful not to let your determination and enthusiasm lead to overtraining.

Research has shown that high-intensity anaerobic (weight-training) sessions lasting sixty minutes are the most effective for stimulating muscle growth. Of course, not all athletes train for the same purpose. While bodybuilders are mainly concerned with building as much muscle mass as possible and keeping body fat low, other athletes may be just as concerned with muscle strength and endurance. So, you need to train for your sport. The good thing is, Macrobiotic Nutrition will support the training regimen of every athlete.

Let's look at how protein in Macrobiotic Nutrition supports training and muscle growth. Intense training has a huge impact on your body's metabolic processes. The hour immediately after an intense workout is called the "anabolic window." During the anabolic window period, the hormonal landscape is primed for muscle building; testosterone is high, growth hormone is high, and insulin is low. To take advantage of this "hormone heaven," it is important to consume a meal high in quality protein and moderate to low glycemic carbohydrates—yes, a Macrobiotic meal. I will expand further on this later. For now, let's stay focused on protein.

The body's synthesis (utilization) of protein increases for about forty-eight hours after an intense training session. By making sure you take in protein immediately after your workout and continuing to do so every few hours, you will make sure your body stays in an anabolic environment. Protein donates amino acids to the body, and amino acids donate nitrogen. When the body has



Photo supplied by Travis Claridge.

Testimonial

"I have seen a tremendous improvement in all areas of my performance since using the Macrobiotic principles. I am bigger, stronger, and have much better muscle endurance and energy."

—**Travis Claridge**, *Offensive Guard,*
Atlanta Falcons

an excess of nitrogen, it is in an anabolic state and builds muscle tissue. Your body's demand for amino acids can be met by one of two sources: the dietary protein found in the foods or supplements you eat, or by the catabolic process of breaking down your own muscle tissue. The last thing you want is for your body to break down muscle to meet your protein demands. This is exactly the opposite of the anabolic state we want to maintain to maximize muscle growth. To avoid catabolism, adequate protein from quality sources must be eaten every two to three and a half hours throughout the day. What are quality protein sources, you ask? The rest of this chapter will answer this question.

MAKING THE GRADE FOR ATHLETES

The bodybuilding industry has many different ways to grade protein sources. Biological value (BV), protein efficiency ratio (PER), net protein utilization (NPU), chemical score, and protein digestibility corrected amino acid score (PDCAAS) are the most common terms you may come across in various articles and ads while reading through a muscle magazine.

BV is one of the most commonly used and is arguably the best measure of a protein's quality. BV is based on how much of the protein consumed is actually absorbed and utilized in the body. The higher the amount of protein (nitrogen) that is actually retained, the greater the BV. If a protein has a BV of 100, it means that all of the protein absorbed has been utilized, and none has been lost. Whole eggs score the highest of all foods with a BV of 100, while beans have a BV of only 49.

Many advertisements for whey list their products as being between 105 and 107 on the BV scale when it really has a BV of near 100. Since BV is defined as "the amount of N [nitrogen] retained for growth and/or maintenance that is expressed as a percentage of N absorbed."

When a protein has a BV of 100+, the company intentionally manipulates the numbers for marketing purposes. They do this by making references to studies in which chemical scores of the whey protein is compared to the amino acid patterns of an "ideal reference protein" to test protein. In this instance, it is possible for the score to exceed 100.

Surprisingly, researchers and the industry seem to consider PER and PDCAAs as the gold standards in protein grading. PER is determined by evaluating how much weight a rat gains while being fed a particular protein compared with a standard such as egg white. The higher the PER, the greater the quality of the protein. The problem is, we are talking about rats! What good is that for humans? Well, the researchers realized that as well, and as a result came up with the PDCAAs. This method uses human amino acid requirements to calculate the amino acid score of proteins. PDCAA compares the amino acid

profile of a particular protein to the exact amino acid requirements for humans as set forth by the Food and Agriculture Organization. The World Health Organization (WHO) and the U.S. Food and Drug Administration (FDA) use PDCAA. When a protein fulfills the requirements for humans, it receives the highest score of 1.0.

Since proteins that have a PDCAA of 1.0 may have a lower PER, and more important, a lower BV, I felt that I needed to establish an even better criteria for grading protein. As mentioned, BV is a measure of how much of the protein will actually be assimilated by the body, but it pays no attention to the amino acids critical for actual muscle growth. Based on my research, I determined that, along with BV, the most important factors relevant to bodybuilders is the concentration of what I will call the “critical five” amino acids and their absorption rate. The five amino acids most important to bodybuilders are glutamine, arginine, leucine, isoleucine, and valine. These critical five amino acids are those most often and most rapidly depleted by athletes due to their training, and the most crucial to muscle recovery and overall growth. Macrobiotic Nutrition emphasizes the importance of these five amino acids when choosing protein sources. The release rate or absorption rate is the rate at which the protein source is broken down to raise blood levels of these amino acids. A protein that provides fast-, medium-, and slow-release rates is optimal. This is covered in greater detail later in the discussion on absorption rates.

Why Nitrogen Balance Is Important

In addition to carbon and hydrogen, amino acids also contain nitrogen as part of their molecular structure. This is a unique characteristic of protein. Specifically, nitrogen balance refers to the condition in which the amount of nitrogen ingested from protein is equal to the amount of nitrogen eliminated as ammonia. A positive nitrogen balance occurs when the body has a surplus of nitrogen. This surplus allows for increased protein synthesis and an anabolic environment to support muscle growth. A negative nitrogen balance, which results from inadequate protein intake, can result in the body's cannibalizing muscle tissue to get nitrogen from the amino acids in your muscles (a catabolic environment). So, the goal is to maintain a positive nitrogen balance and prevent a negative nitrogen balance. Macrobiotic Nutrition allows you to stay in a positive nitrogen balance and be anabolic—all of the time.

Amino Acids

Amino acids are the building blocks of all proteins. Different proteins are made up of different combinations and concentrations of amino acids. When we ingest protein, whether it is from food or supplements, it is digested in the stomach and broken down into its amino acid constituents. These amino acids enter what is known as the amino acid pool. This is the body's storage mechanism for excess amino acids, which will be used at a later time for either:

1. Muscle growth
2. Conversion into glucose for energy
3. Synthesis of fatty acids and ketone bodies
4. Maintenance of nitrogen balance

Amino acids can be classified as either essential or nonessential. By essential, I mean those particular amino acids that can't be synthesized by the body but must be derived from the foods we eat. The nine essential amino acids are isoleucine, lysine, methionine, phenylalanine, threonine, tryptophan, valine, leucine, and histidine.

As mentioned earlier, the value of a protein for new tissue growth is determined by the presence and concentration of these amino acids. Since some amino acids are found in smaller amounts in some foods, the supply of these amino acids will be used up faster in the growth of new muscle if these foods are your primary protein sources. In foods, these amino acids are considered "limiting." The way to avoid limiting amino acids and therefore inhibiting muscle growth is to vary protein sources taken in throughout the day. Consequently, Macrobiotic Nutrition recommends utilizing various protein sources. This gives the body a better spectrum of amino acids, since different proteins contain different amounts of certain amino acids. Animal source proteins have a better overall concentration of essential amino acids, which becomes a problem for vegetarian bodybuilders. Vegetarians need to complement (combine) certain protein sources in order to make up for the gaps created by limiting amino acids. The problem this creates with Macrobiotic Nutrition is that most vegetable source proteins are much too high in carbohydrate content. This makes it nearly impossible for a vegan (one who eats no animal products) to adhere to the Macrobiotic principles, unless they compensate for these deficiencies through supplementation. Even then, it would be very difficult to get adequate amounts of the crucial amino acids.

For an athlete, the most crucial amino acids are the critical five of Macrobiotic Nutrition: glutamine, arginine, leucine, isoleucine, and valine. As I

stated earlier, these amino acids are utilized by working muscle tissue for a variety of functions and are rapidly depleted in the body. In gluconeogenesis, a catabolic process, the body breaks down protein into amino acids and converts them to glucose when blood sugar levels get too low. This commonly occurs in low-carb diets; that is why Macrobiotic Nutrition places such emphasis on eating carbohydrates to be utilized for glucose, so that the body will not draw on protein or muscle tissue for this purpose.

The Critical Five Amino Acids

Glutamine. This member of the critical five amino acids is quite possibly the most important amino acid for athletes to replenish through food and supplementation. Although glutamine is considered a nonessential amino acid because it can be synthesized (manufactured) by the body from glutamic acid, I consider it essential to athletes because of their increased need and demand for this multifunctional amino acid.

Glutamine is the most abundant amino acid in skeletal muscle. Your body is constantly tapping into your muscles' glutamine reserves to perform various functions. Heavy weight training has been shown to deplete glutamine from muscle tissue, in response to the physical stress involved. Long, strenuous workouts trigger the release of cortisol. Cortisol is a catabolic hormone, which your body releases in response to physical and emotional stress. Supplementing glutamine after workouts has been shown to offset the catabolic effects of cortisol. This is extremely important, because the last thing you want is to be in a catabolic state after a workout. You want the exact opposite, an anabolic state, to occur. Taking glutamine immediately following your workout will help shift you into an anabolic state.

Glutamine can also help buffer lactic acid. You know that feeling you get when you're training hard, and your muscles feel so burnt and exhausted that you can't squeeze out another rep? That feeling is caused by lactic acid buildup in your muscles. Glutamine helps buffer the lactic acid and delay muscle exhaustion. Lactic acid can be even further buffered by taking glutamine in a bicarbonate (effervescent) form, because minerals like sodium bicarbonate and potassium bicarbonate have a buffering effect on lactic acid.

Glutamine also plays an important role in strengthening your immune system by acting as an antioxidant enhancer. Intense training increases the production of free radicals, invaders that attack your cells. Glutamine helps raise levels of the powerful antioxidant glutathione to combat the damage caused by free radicals. In addition, glutamine protects the body from high levels of the toxic compound ammonia, which can increase as a result of exercise.

Glutamine allows more vitamins, minerals, water, carbohydrates (such as glycogen), and protein (such as amino acids) to enter the muscle, thus performing as a cell “volumizer.” The improved nutrient absorption provided by glutamine increases muscle size not only by increasing intercellular capacity, but also by feeding your muscles to stimulate muscle growth and improve recovery.

Another benefit to increased cell volumization is the heightened sensitivity the now “swelled up” cell will have to anabolic hormones (like insulin and testosterone). Researchers measured the response of anabolic hormones in cells that were volumized and cells that were shrunk. First, the cells were increased in volume by changing the osmolarity, or salt content, of the fluid around the cell. When these cells were exposed to the anabolic hormones, they swelled up with water, just like a sponge. Now, we want that water inside the cell. This is what helps to push carbohydrates into the cell and to promote glycogen storage. When researchers ran the same test without the use of the anabolic hormones (that is, increased the volumization of the cells), the cellular response was the same. The cells grew. However, cells that were shrunk down or dehydrated became catabolic and did not respond to the stimulus. So, it is pretty simple: increasing the cell volume is anabolic; decreasing cell volume is catabolic. When you supplement with glutamine, the amino acid transporters on the cells will carry the glutamine into the cell, resulting in swelling. The research tells us that this swelling will put the cell in an anabolic environment, leading to progressive muscle gains.

Wait! There is even more to say about this dynamic amino acid. Glutamine is also one of the most researched and effective natural growth hormone (GH) boosters. Many studies have been performed validating glutamine’s effects in raising GH levels. And we all know that elevated GH means increased muscle and decreased body fat. In most of the early studies, fairly large doses of glutamine were required to show significant raises in GH. This is due to the fact that stomach acids destroy a lot of the glutamine and a considerable amount is used up by the small intestine. However, in a more recent study by Thomas Welbourne reported in the *American Journal of Clinical Nutrition*, only 2.5 grams of glutamine in a bicarbonate (effervescent) drink was found to increase GH levels by 30 percent. What that 30 percent increase means to you is more lean body mass. If I were you, I’d take advantage of this and slam down 2–4 grams of Effervescent Glutamine right after every workout.

So, as you can see, glutamine is invaluable to an athlete. Aside from these many benefits, glutamine is a major contributor to some key principles of Macrobolic Nutrition such as maintaining lower blood sugar levels, higher glycogen levels, and nitrogen balance. Exercise increases your body’s demand for

both carbohydrates and protein, and a sound Macrobiotic Nutrition diet provides the carbohydrates and protein you need to meet these demands. However, glutamine supplementation is even more important if you're not meeting these demands, because in that case you are not calculating your nutrition requirements accurately, you are not following the program correctly, or you are severely overtraining. As you may recall from Chapter 2, your body converts glutamine to glucose through gluconeogenesis in the absence of carbohydrates and glycogen. The glucose is then used as an energy source. If glutamine is not replenished quickly, this can rapidly deplete your body's glutamine stores and result in a negative nitrogen balance and in catabolism.

My recommendation is to choose protein sources high in glutamine (see "Glutamic Acid" in Table 3.10 later in this chapter) to ensure that you get enough of this critical amino acid. Supplementing with additional glutamine postworkout is also a good idea.

Arginine. Arginine is one of our acclaimed critical five amino acids, and it is crucial for muscle growth. For an athlete, arginine is actually a conditionally essential amino acid. Exercise, like intense training in the gym, increases your body's demand for this amino acid. Arginine has been shown to decrease somatostatin in the body. Somatostatin is the hormone that inhibits GH release. GH has tremendous muscle-building and fat-burning properties. If we want to pack on lean mass, we definitely do not want to limit the amount of GH being released from the pituitary. Arginine's somatostatin-inhibiting effects will elevate GH levels.

As far as strength is concerned, arginine is used as a metabolic intermediate in the formation of creatine phosphate. Creatine phosphate has been shown to increase cell ATP (adenosine triphosphate), which is your muscle cells' first source of energy for any action. By increasing your ATP, you will help increase your muscle strength and performance.

The cardiovascular benefits of arginine are well documented. Arginine works as a potent vasodilator via nitric oxide, as shown in a study by A. Calver, et al., reported in *Clinical Science* in 1991. Arginine is a precursor for the production of nitric oxide, which has been shown to increase dilation in the peripheral vasculature (veins), a great asset for bodybuilders. By increasing blood flow, you will maximize and maintain pumps during and after your workout. You will also increase the blood flow into your working muscle cells. This serves two purposes: First, it allows for an increase flow of nutrients, like amino acids, into your muscles. Second, it allows for better removal of free radicals and ammonia from the breakdown of muscle tissue. By ingesting about 4 grams of L-arginine daily in one dose, athletes can increase the production of nitric

oxide. This is a pretty large dose, but it is the amount needed to significantly increase nitric oxide production. Some companies are making supplements utilizing AAKG (Arginine Alpha Ketoglutarate), but no research has been done on AAKG, so stick with pure arginine if you want to be sure of the true benefits and mega pumps.

Another function of arginine for athletes is its role in ammonia detoxification. High levels of ammonia in cells can be toxic. Arginine converts ammonia to urea, which is then excreted by the body. Since exercise can increase the production of ammonia, arginine would be beneficial for highly trained athletes.

Furthermore, arginine had been found to improve wound healing and to play a role in the immune system. For all of the reasons discussed, arginine is one of the critical five amino acids.

BCAAs: Leucine, Isoleucine, and Valine. The branched-chain amino acids (BCAAs) are made up of valine, leucine, and isoleucine. These three amino acids are considered essential because our bodies do not create them, and we need to get them from our diet. We store these amino acids in our liver and in skeletal muscle. For bodybuilders, these three aminos are a very important part of the critical five. Our bodies need them for the maintenance of muscle tissue, preservation of muscle glycogen, and prevention of the breakdown of muscle protein.

Ingestion of BCAAs has been shown to preserve muscle glycogen levels. This occurs because the BCAAs are a primary amino acid energy source for your body. To produce energy, your body burns the carbon skeletons from these amino acids and then converts the nitrogen residues into alanine. The additional alanine formed from this process is shuttled to the liver to produce glucose via gluconeogenesis. The glucose is transported to working muscles to be used as fuel. In the case of bodybuilders and strength athletes, BCAAs preserve muscle glycogen levels and prevent muscle breakdown by giving the body the starting materials it needs for this energy-producing process, so your body does not need to derive the BCAAs from skeletal muscle. Sparing the muscle glycogen also preserves muscle volume and prevents muscle breakdown. Since an athlete's training creates a great demand for BCAAs to replenish glycogen and provide energy, it is important to consume foods or protein supplements high in BCAAs.

Supplementing the Critical Five

A Macrobiotic diet should provide good quantities of the critical five amino acids from whole-food proteins and protein supplements. In addition to what you will get from these sources, I would also recommend taking a 4-gram dose

RESEARCH UPDATE

BCAAs Aid in Increasing Strength and Lean Body Mass

A 1997 study reported in *Medicina Dello Sport* looked at the effect of taking supplemental BCAAs on bodybuilding progress. The study involved thirty-one male bodybuilders between the ages of eighteen and thirty-four, all of whom were natural, drug-free bodybuilders and had at least two years of training experience. The subjects were divided into two groups—one group of sixteen took a placebo and the other group of fifteen took 0.2 gram of a BCAA supplement per kilogram (2.2 pounds) of body weight thirty minutes before training and thirty minutes after training. The results showed that while both groups experienced increases in body weight, the BCAA group had greater weight gains. An analysis of the weight gain in the BCAA group showed increases in the lean body mass in that group whereas the group taking the placebo showed no lean-mass gains. Strength gains and improvements in exercise intensity were also seen in the BCAA group in both the squat and bench-press exercises. The placebo group gained strength only in the squat exercise.

of arginine preworkout and 2–4 grams of glutamine (preferably Effervescent Glutamine) postworkout. I don't feel taking supplemental BCAAs is necessary, because your Macrobiotic meals and protein supplements should provide enough. However, if you eat a lot of vegetable proteins, taking some extra BCAAs may be a good idea.

CHOOSING QUALITY PROTEIN SOURCES

Now that you can clearly see the importance of protein for muscle growth and now that you are aware of the important components to look for in your protein sources, let's look at which protein sources will provide high concentrations of critical amino acids and high biological values, along with low levels of saturated fat.

Whole Foods versus Supplements

Which is a better source of protein: whole foods or quality-protein supplements? Ten years ago this book may have recommended whole-food protein sources over protein supplements. But today, thanks to recent developments

and advancements in protein supplement manufacturing technologies, I feel that both whole-food sources and the right protein supplement sources should be part of your Macrobiotic Nutrition program.

Whole-food protein sources are an extremely important part of Macrobiotic Nutrition. The recommended food sources for protein include lean red meat, chicken, turkey, fish, whole eggs and egg whites, and some low-fat dairy sources. Each of these sources not only provides protein and all of the essential amino acids, but they also provide other important nutrients to support muscle growth, performance, and overall health. Animal proteins are better sources than plant proteins because they have the proper proportions of essential amino acids. They are called complex proteins because they contain all the essential amino acids in amounts that are sufficient for the maintenance of growth. Vegetable sources are considered incomplete proteins and are usually deficient in one or more of the essential amino acids.

Because whole foods require digestion, they provide a slow, steady release of amino acids into the bloodstream. This slow release helps you to maintain a positive nitrogen balance from meal to meal. Good digestion is another important reason to be sure to eat a few whole-food meals and a few protein supplement meals. The human body is designed to digest whole foods. Our digestive systems cannot handle being fed predigested protein powders all day long. You must feed your body whole foods to take full advantage of the di- and tripeptides available after foods are digested. Ingesting whole food is also important to keep your body producing the enzymes needed to break down and digest food and to help elimination processes.

Digestion of Protein and Amino Acids

The mechanical digestion of protein begins in the mouth during chewing. In the stomach, the enzyme pepsin joins in, breaking down the protein into shorter peptides. The partially digested protein then passes into the intestines, where the free-form dipeptide and tripeptide amino acids immediately begin to be absorbed. Enzymes continue to digest any polypeptides as they travel down the intestines.

Once the free-form dipeptide and tripeptide amino acids enter the bloodstream, they are transported to the liver, where a few things may happen to them. They may be converted into other amino acids, they may be used to make other proteins, they may be further broken down and either used for energy or excreted, or they may be placed into circulation and continue on to the rest of the body. Proteins empty from the stomach in two to three hours, depending on how much fat is present.

The other natural benefit to whole foods is the thermic effect of food

(see the inset “The Thermic Effect of Food” in Chapter 6). This refers to the amount of calories your body needs to digest and use the nutrients found in the food. For some protein foods, such as chicken, the thermic effect may be as high as 30 percent of the calories contained in the food. So, to digest and use a 200-calorie chicken breast, your body may need about 60 calories. In actuality, then, the chicken is only providing 140 net calories for energy or fat storage. On the other hand, something like orange juice, which does not require as many calories to break down, will yield more net calories per 200 calories consumed. If you are aware of this factor, you can avoid packing on the fat.

While whole food must be a main source of protein, I feel protein supplements also need to be taken, especially at critical times. Aside from being a quick and convenient way to make sure you’re getting enough protein, protein supplements are now superior to what they once were due to the recent advancement of technologies used to develop them. I remember when I started training in the 1980s how nasty some of the stuff tasted, but an even bigger problem was that the processing techniques compromised the quality of the protein. Today, processing techniques keep the integrity of the protein while removing the components in the food source you don’t want, like the lactose and fat in whey.

Protein supplements can also be designed to combine several protein sources like whey, soy, and milk to get high concentrations of essential amino acids. Protein manufacturers are also adding other ingredients to their protein supplements to increase its effectiveness and provide additional benefits. For example, the addition of fiber slows the absorption of the protein supplement to help maintain nitrogen balance. Some products are adding free-form amino acids to improve the amino acid profile.

Now that technologies allow manufacturers to really custom design supplements, they can be better than whole food at certain times. Look for a supplement that combines whey, soy, and casein. This trio of proteins provides sustained release of high quantities of the critical five amino acids. This is exactly what you need to stay in a positive nitrogen balance and feed your muscles what they need to grow. I feel that the properly designed protein supplement should be taken three times a day: first thing in the morning, postworkout, and before bedtime.

My overall recommendation is to consume three to four whole-food meals and two to three supplement meals each day, as this will provide you the best balance of Macrobiotic Nutrition. Remember, different protein sources have different concentrations of amino acids. By rotating food and supplement meals and eating every two to three and a half hours, you will minimize any limiting of amino acids, maximize the supply of critical amino acids, and keep your muscles primed for growth!

TABLE 3.2. WHOLE FOOD NUTRIENT CONTENT

Source	Portion	Calories (kcal)	Protein (g)	Fat (g)	Carbohydrates (g)
Chicken breast, boneless	4 oz	130	27.0	1.5	—
skinless	6 oz	187	39.0	2.0	—
Pork, tenderloin	4 oz	162	21.0	8.0	1.0
Turkey, ground	4 oz	170	20.0	9.0	—
Beef, tenderloin	3 oz	200	23.0	11.0	—
Venison	3 oz	127	25.0	2.0	—
Beef, ground, 95% lean	3.5 oz	137	22.0	5.0	—
Turkey breast	3.5 oz	110	22.0	2.0	—
Tuna	6 oz	220	41.0	5.0	—
Salmon, filet	7 oz	281	40.0	12.0	—
Whitefish	7 oz	265	38.0	12.0	—
Tofu, raw regular	4 oz	94	10.0	6.0	2.0
Cottage cheese 1%	8 oz	163	28.0	2.0	6.0
Skim milk	8 oz	80	8.0	—	12.0
Yogurt, low-fat	8 oz	140	14.0	0.5	19.0
Chicken leg, meat and skin	1	190	18.0	12.0	—
Eggs, whole, large	1	75	6.0	5.0	0.6
Egg white, large	1	17	3.5	—	0.3
Egg substitute, liquid	3 oz	80	11.0	3.0	1.0

WHOLE-PROTEIN SOURCES

Beef: The Big Mass Protein

Any bodybuilder knows that beef is a staple in the diet when packing on the mass. Beef definitely has its place in the Macrobiotic cookbook. Beef's popularity can be linked to the fact that it is an excellent natural source of both creatine and L-carnitine. Both of these nutrients have been shown to enhance cardiovascular health as well as aid in muscle growth. No wonder athletes love it! Beef is also an excellent source of zinc, vitamin B₁₂, and phosphorus. Beef

has a good amount of the essential amino acids, especially the BCAAs, making it great as far as muscle growth is concerned. And in my opinion, it is the best tasting of all the protein sources. On the flip side, not all is good. Some sources of beef are very high in saturated fat and cholesterol, two things that will definitely sabotage your cardiovascular health. You must choose lean beef cuts such as top round, flank steak. These sources will provide a good protein along with beef's other attributes.

TABLE 3.3. PROTEIN/NUTRITIONAL VALUE OF SELECT CUTS OF BEEF

6 ounces (raw weight)	Tenderloin	Tip round	Top round	Top blade	Flank
Calories	454	303	277	231	306
Protein (g)	30.65	33.58	37.13	32.71	33.51
Fat (g)	31.11	15.24	11.14	8.78	15.85
Saturated Fat (g)	14.53	6.97	5.10	3.39	7.69
Total BCAAs (g)	5.23	6.20	6.41	6.21	5.79
Total Critical Five Amino Acids (g)	11.69	13.86	14.33	13.50	12.94
BV (g)	76	76	76	76	76

Chicken Breast: The Bodybuilder's Easy Choice!

Chicken breast's popularity can be completely linked to the fact that it has an extremely high protein to fat ratio. This food protein is low in fat and saturated fat is almost nonexistent, making it a heart-healthy choice. This makes it safe as a frequent protein source in your diet. It is also a great source of potassium. The natural low-fat content makes it a great protein source in a Macrobiotic meal and leaves room for the addition of good fat sources such as olive oil to your salad or vegetables.

TABLE 3.4. PROTEIN/NUTRITIONAL VALUE OF CHICKEN BREAST

Serving Size	6 ounces
Calories	195
Protein (g)	40.75
Fat (g)	2.25
Saturated Fat (g)	0.60
Total BCAAs (g)	7.24
Total Critical Five Amino Acids (g)	15.84
BV (g)	76

Whole Eggs: Don't Throw Out All the Yolks

As far as food protein is concerned, eggs are the king! A whole egg has a BV of 100, equal to many protein powders, and also is a great source of the omega-3 fatty acids your body craves to decrease cholesterol levels, aid in joint inflammation, and increase hormone production. Whole eggs also serve as a great source of sulfur. Your body utilizes sulfur in the formation of many anabolic hormones in the body. Eggs are high in the BCAAs, which make them ideal for muscle growth, and they score a 1 on the PDCAA scale. This is due to the fact that they have an amino acid pattern very similar to mother's milk, the protein most fully assimilated by our bodies. With such a high level of essential amino acids and BCAAs, egg is the Macrobiotic Nutrition's first choice of food proteins.

Since the egg yolk is the source of fat and some of the essential amino acids, it is necessary to include some yolk in a whole egg/egg white mix to achieve the optimal ratio of nutrition. A whole egg has about 7 grams of protein with 3.5 grams from the yolk and 3.5 grams from the white. The yolk

TABLE 3.5. PROTEIN/NUTRITIONAL VALUE OF WHOLE EGGS AND EGG WHITES

Whole Eggs	
Serving Size	1 whole egg
Calories	75
Protein (g)	6.25
Fat (g)	5.00
Saturated Fat (g)	1.55
Total BCAAs (g)	1.25
Total Critical Five Amino Acids (g)	2.45
BV (g)	100
Egg Whites	
Serving Size	3 egg whites
Calories	50
Protein (g)	10.40
Fat (g)	—
Saturated Fat (g)	—
Total BCAAs (g)	2.13
Total Critical Five Amino Acids (g)	4.08
BV (g)	100

has about 3.5 grams of protein and about 5 grams of fat. I have found that including one yolk with every four whites (one whole egg and three extra whites) delivers 17.5 grams of protein and about 4 to 5 grams of fat. So, if a dozen eggs are used, you will have three whole eggs and nine additional whites. This will deliver approximately 52.5 grams of protein and 15 grams of fat.

Fish: A Great Catch

Tuna is a favorite among bodybuilders. Its most redeeming quality is its almost nonexistent fat content. Tuna has about the best protein to fat ratio of all whole food proteins. It has a pretty good concentration of BCAAs, so it ranks fairly well for a whole-food protein source. Like red meat, tuna also has a high amount of creatine in it, which is a nice bonus from a protein source.

Tuna is not the only fish acceptable in the Macrobiotic plan. Salmon is exceptionally high in omega-3 fatty acids (good fats) and is a solid source of complete protein high in essential amino acids. Macrobiotic Nutrition is also an

TABLE 3.6. PROTEIN/NUTRITIONAL VALUE OF FRESH TUNA AND SALMON

Tuna	
Serving Size	6 ounces
Calories	197
Protein (g)	43.40
Fat (g)	1.40
Saturated Fat (g)	0.40
Total BCAAs (g)	7.76
Total Critical Five Amino Acids (g)	16.83
BV (g)	76
Salmon	
Serving Size	6 ounces
Calories	306
Protein (g)	34.10
Fat (g)	17.75
Saturated Fat (g)	4.26
Total BCAAs (g)	6.09
Total Critical Five Amino Acids (g)	13.23
BV (g)	76

advocate of any white fish such as haddock, flounder, shark, swordfish, and mahi-mahi. All of these are lean choices with a high protein to fat ratio.

Fish, like any other food protein, must undergo digestion in order to release amino acids into the bloodstream. This makes it a slow-release protein source, but it is faster than the harder-to-digest food sources like red meat.

New Research Bulletin

Fatty fish may offer an added fat-loss perk. Incorporating a daily meal of fatty fish may decrease fat cells and enhance fat loss by raising the amount of leptin in the body. Leptin is a hormone that decreases hunger and shrinks the size of fat cells—good stuff!

Pork Tenderloin

Yes, pork is the other white meat. Certain cuts of pork are a very lean source of dietary protein that can add a little variety to your diet. Pork is high in zinc, iron, and B vitamins and has an excellent protein to fat ratio. Most of the fat in pork is on the outside of the actual meat fibers, rather than intertwined within them. This means that when the fat is trimmed off the meat, very little fat remains. Pork has about 3 grams of fat for every 30 grams of protein, which is an exceptional ratio. About two-thirds of the fat in pork is unsaturated. Unsaturated fat cannot be converted into cholesterol, so lean pork will have no impact on your blood cholesterol levels. Pork actually contains less cholesterol than chicken. I recommend using the tenderloin cuts as part of your Macrobiotic diet. Sorry, bacon, pork sausage, and spareribs aren't on the Macrobiotic menu—only lean cuts will do.

TABLE 3.7. PROTEIN/NUTRITIONAL VALUE OF PORK TENDERLOIN

Serving Size	6 ounces
Calories	204.00
Protein (g)	35.70
Fat (g)	5.80
Saturated Fat (g)	2.00
Total BCAAs (g)	6.47
Total Critical Five Amino Acids (g)	14.28
BV (g)	76.00

PROTEIN SUPPLEMENT SOURCES

As I mentioned earlier, various protein supplements are now available, which can be very beneficial in helping athletes meet their protein requirements. In some instances supplements are better than whole foods, thanks to major advancements in manufacturing technologies. Let's take a look at some of the best protein supplement sources out there.

Whey

Whey protein is a very popular protein source used in sports nutrition supplements. Some of this popularity is due to whey's many benefits, while some is also the result of advertising hype by companies looking to sell product. As you thumb through the muscle mags, you see ad after ad from supplement companies touting their whey to be the "superior protein." All hype aside, whey protein ranks high in Macrobiotic Nutrition, but it's not the answer to all your protein supplement needs. Let's take a look at what whey protein really has to offer.

Believe it or not, whey was once looked upon as a waste product of the cheese manufacturing process. Whey makes up 20 percent of milk protein (casein makes up the other 80 percent) and is a byproduct from the separation of the liquid whey from the more solid curd in cheese manufacturing. Cheese manufacturers used to throw away the liquid whey until someone finally realized they were disposing of the most nutritious part of the milk. Depending on the processing techniques used, whey can contain anywhere from 34 to 90 percent protein. The whey with lower concentrations of protein are typically used in food products, while the whey with higher concentrations are used in nutritional supplements.

Whey Grades: Concentrate and Isolate

The whey proteins used in sports nutrition usually contain protein in a range of 70 to 90 percent or more. Those that range from 70 to 89 percent are classified as whey protein concentrate (WPC). A concentration of 80 percent is usually the standard used in most WPC supplement products. Whey with a protein content of 90 percent or higher is classified as whey protein isolate (WPI). The higher the concentration of protein, the lower the fat and lactose content, so WPI obviously has a higher protein content with less fat and lactose than WPC. In fact, WPI is virtually lactose and fat free, containing less than 1 percent of each. (See Table 3.8.) The higher protein content in WPI also yields a higher concentration of amino acids. However, both WPC and WPI contain high levels of BCAAs and glutamine (in the form of glutamic

acid) and a fair amount of arginine. So, it ranks pretty well on our Macrobo-lic critical five score.

Whey ranks very high with a PDCAA of 1.14 (actually over the top 1.0 ranking set by the USDA) and also gets the highest BV score available at 100. Whey's high BV of 100 reflects how efficiently the body will utilize its amino acids to maintain nitrogen balance.

Aside from whey's high protein, high amino acid content, and high BV, it also contains protein fractions. The protein fractions lactalbumin, lacto globulin, immunoglobulin, lactoferrin, and lactoperoxidase offer important benefits. Studies have shown that these fractions have effective antioxidant and immune support benefits, which can improve a bodybuilder's recovery from a stressful workout. These whey protein fractions also work to remove ammonia, lactic acid, and cortisol, three catabolic byproducts of exercise that hinder muscle growth and performance. The removal of these catabolic substances is critical for achieving maximum muscle growth, performance, and recovery.

TABLE 3.8. TYPICAL PRODUCT ANALYSIS OF WHEY PROTEIN

Whey Protein Concentrate (WPC)	Value
Protein (N x 6.38) as is	81.00%
Moisture	4.10%
Ash	2.50%
Carbohydrates as Lactose	2.60%
Fat	7.90%
Whey Protein Isolate (WPI)	Value
Protein (N x 6.38) dry basis	92.00%
Moisture	4.00%
Ash	3.00%
Carbohydrates as Lactose	<1.00%
Fat	<1.00%

The process in which whey is manufactured is critical for obtaining these fractions. Some processing techniques denature whey and destroy these fractions. WPC is typically higher in protein fractions than WPI. Ion-exchanged WPI processing retains lactoglobulins and lactalbumin but is lacking in the other fractions. One type of WPI processing called "membrane processing" passes the whey through a variety of different membrane type filters, which

separates out of the fat and lactose while retaining the protein and immunoglobulin. As you can see, processing also makes a difference in the quality and functionality of whey supplements. The good news is that most reputable supplement companies are using the proper manufacturing processes to produce quality whey products.

The Final Analysis

Whey makes a great source of protein with its BV score of 100, high amino acid content, and protein fractions. The fact that it is a fast (rapidly absorbed) protein needs to be considered. As a fast-release protein, it will quickly supply amino acids to your muscles to stimulate muscle growth and prevent catabolism. This makes whey a great postworkout protein source, as it will quickly replenish depleted amino acid levels. However, whey protein's fast-release property means it will not maintain amino acid levels. Therefore, I recommend using it in conjunction with the other slower-release proteins.

All in all, WPC and WPI definitely make great protein sources for athletes. Whey should definitely be one of your protein sources, but don't believe all the advertising hype and disregard other great protein supplement sources like casein, soy, and milk protein isolate.

Caseinate

Caseinate, or casein, has made a sudden surge into the sports nutrition arena. Once thought of as an inferior protein because of its low BV of 77, casein has recently gained praise from sports nutritionists. For years, the whey industry pounded the comparison of whey's BV of 100 to casein's measly 77. But a study conducted by Dr. Yves Boire reported in the *Proceedings of the National Academy of Sciences* in 1997 gave casein some firepower to fight back. In this study, cows were fed radioactively labeled leucine. The milk from these cows was separated into labeled whey and labeled casein, and both were fed to humans. The study showed that whole-body muscle breakdown was inhibited by 34 percent after ingestion of the casein, while these effects were not seen in the whey group. The whey produced a short-lived elevation in blood amino acid, while casein provided a more level plateau of increased blood amino acids. This resulted in better nitrogen retention and an overall increase in anabolism with a decrease in muscle catabolism. I'd say casein just leveled the playing field on whey's higher BV!

Additional benefits of casein supplements include its high protein content. Most casein yields between 85 and 90 percent protein and is extremely low in fat, carbohydrates, and lactose. Casein is also fairly high in BCAAs and glutamine.

TABLE 3.9. TYPICAL PRODUCT ANALYSIS OF CALCIUM CASEINATE

Calcium Caseinate	Value
Proteins	88.00%
Moisture	6.00%
Total Fat	1.80%
Ash	5.00%
Lactose	0.20%

Casein's slower release rate and its ability to maintain blood amino acid levels for a longer period of time will keep you in a positive nitrogen balance between meals and during the night while sleeping. Combining casein with whey ensures both fast amino acid saturation and sustained nitrogen retention.

Egg Protein Powder

Egg protein is generally sourced from either egg whites or whole egg. Ovalbumin or egg white is generally used as the reference protein for PER (protein efficiency ratio) comparisons when grading protein quality. Egg protein has slowly lost favor, due to its higher price tag in the supplement industry. An article published in *Functional Foods and Nutraceuticals*, by R. Kreider, discusses research by V. Gattas which illustrates that egg protein promotes anabolism by promoting nitrogen retention. The results show that egg is as good as whey, casein, and colostrums for making these anabolic changes happen.

Since egg protein powder may not be high on the list for taste, I must recommend the use of eggs in your diet on a daily basis. Egg whites have classically been a favorite of bodybuilders, but for every four whites, you must have a yolk. The egg yolks strengthen the amino acid profile and add omega-3 fatty acids (the importance of which is covered in the discussion on fats). Twelve egg whites and three yolks are an easy, and relatively cheap, way to get about 50 grams of high-quality food protein for one of your Macrobiotic meals.

Soy

What I'm talking about here is soy protein powder, more specifically, Supro[®] soy protein, which is the only vegetable protein that is a complete protein. The Solae Company has changed the rules as far as soy protein powder is concerned with the introduction of their Supro[®] powder. Unlike other foods (like bread), which are processed to extend shelf life and to improve flavor, proteins are processed to increase BV and actually improve their functionality. Manufacturers can sometimes turn a crappy protein into a muscle-building giant.

This is the definitely the case with (Supro[®]) soy protein isolate. By using advanced manufacturing techniques, Solae, the makers of Supro, have been able to use an inferior source like soy beans to create a superior muscle-building protein. With a BV of 100 and large amounts of glutamine, arginine, and BCAAs, Supro rates high in Macrobiotic Nutrition!

The health benefits of soy protein have been well documented and researched by the Solae Company. For athletes, these benefits are even more important because they serve both general health and physical performance-enhancement functions. A compilation of thirty-eight studies (by J. W. Anderson, et al., published in the *New England Journal of Medicine* in 1995) reported that soy protein consumption resulted in a significant reduction in both total cholesterol and LDL cholesterol as well as in serum triglycerides. The FDA approved a claim linking soy protein consumption to decreased cardiovascular risk in October 1999. The claim states that a diet low in saturated fat and cholesterol in combination with daily consumption of 25 grams of soy protein reduces the risk of coronary heart disease.

Increased physical activity leads to an increase in oxidative stress and damage to the body and an elevation in serum (blood) free radicals. Skeletal muscle is especially susceptible to oxidative stress and damage. In 1995 Anderson, et al., noted that supplementation with Supro[®] soy protein (with high levels of isoflavones) appeared to provide potent antioxidants that delivered protective antioxidant properties. In a whey/soy study conducted by Rossi and colleagues at the Ohio State University in 1998, two groups of subjects were given either 40 grams of Supro[®] soy or 40 grams of whey. After three weeks, the soy group had significant increases in their total antioxidant status, while the whey group had significant reductions.

The view that soy will increase estrogen levels in men is very inaccurate. In fact, it is my belief and the belief of many scientists that soy actually lowers estrogen activity in men. The key word here is “activity.” Let me explain why this is so.

The view that soy functions as an estrogen stems from the isoflavones found in soy. These compounds have chemical structures similar to estrogen, and because of this similar chemical structure, they bind to and block estrogen receptor cells. By blocking the estrogen cell receptors, the estrogen circulating in the blood will not have any estrogenic effects and is inactive. These isoflavones work similarly to the popular antiestrogen drug Nolvadex[™] (Tamoxifen Citrate) used by bodybuilders. Lowering estrogen activity is always a goal for bodybuilders, because estrogen increases body fat, water retention, and gynecomastia (excessive development of the breast in the male), which

are all highly undesirable characteristics for men and particularly for bodybuilders.

Furthermore, the isoflavones in soy are considered the key compounds responsible for many of the health benefits associated with soy consumption. In a recent study (see “Research Update” below), young men consumed soy protein daily over a two-month period. Over the course of the study, soy consumption decreased DHT (the testosterone derivative that is associated with benign prostatic hyperplasia—prostate enlargement) and baldness, but *did not affect* other types of testosterone.

So, let’s set the record straight about soy protein. High-quality soy-based protein supplements have many advantages for bodybuilders and athletes, and should definitely be an important part of their nutrition program.

RESEARCH UPDATE

Effects of Soy Isoflavone Consumption on Reproductive Hormones in Healthy Young Men

Thirty-five healthy men (average age range of 22 to 33 years old) consumed milk protein isolate (MPI), low isoflavone soy protein isolate (low-iso SPI); 1.6 +/- 0.19 milligrams isoflavones/day) and high isoflavone SPI (high-iso SPI; 61.7 +/- 7.4 milligrams isoflavones/day) for fifty-seven days, each separated by four-week washout periods, in a randomized crossover design. Conclusions: Soy protein, regardless of isoflavone content, decreased serum DHT and DHT/testosterone with minimal effects on other androgens, estrogens, gonadotropins, or SHBG [sex-hormone-binding globulin] in healthy young men.

B. L. Dillingham, B. L. McVeigh, J. W. Lampe, and A. M. Duncan,
“Effects of Soy Isoflavone Consumption on Reproductive
Hormones in Healthy Young Men,” Department of Human Biology
and Nutritional Sciences, University of Guelph, Ontario, Canada.
Fred Hutchinson Cancer Research Center, Seattle, Washington, USA

AMINO ACID CONTENT OF PROTEIN SOURCES

Table 3.10 on the following pages provides the amino acid content of various protein sources, along with its total critical five amino acid content based on 100 grams of protein. Note: Probolic™ protein, which will be discussed shortly, has the highest total value of critical amino acids to support muscle growth.

TABLE 3.10. AMINO ACID PROFILES OF COMMON PROTEIN SOURCES

Ingredient	Probiotic™	Soy Protein Isolate	Egg Protein (Dried)	Milk Protein Isolate	Calcium Caseinate
Alanine	4.23	4.30	5.77	3.50	3.00
Arginine	7.00	7.60	5.43	3.50	3.70
Aspartic Acid	11.13	11.60	10.18	8.00	6.90
Cysteine/Cystine	1.25	1.30	2.59	0.60	0.40
Glutamic Acid	18.78	19.10	13.29	20.80	20.90
Glycine	3.83	4.20	3.49	1.90	1.80
Histidine	2.60	2.60	2.26	2.70	2.90
Isoleucine	4.60	4.90	5.66	4.40	4.60
Leucine	9.33	8.20	8.41	10.30	9.10
Lysine	6.50	6.30	6.80	8.10	7.70
Methionine	1.38	1.30	3.44	3.30	2.90
Phenylalanine	5.10	5.20	5.82	5.00	5.10
Proline	5.43	5.10	3.91	9.50	10.40
Serine	5.18	5.20	6.88	6.20	5.80
Threonine	3.73	3.80	4.55	4.50	4.30
Tryptophan	1.28	1.30	1.23	1.40	1.20
Tyrosine	3.78	3.80	3.91	5.20	5.50
Valine	6.10	5.00	6.37	5.70	5.70
Critical Five Amino Acids	45.81	44.80	39.16	44.70	44.00

STAY ANABOLIC: WHAT PROTEINS AND WHEN?

One of the major aspects of Macrobiotic Nutrition is meal timing. The 45/35/20 lean-mass equation is designed to keep your body in an anabolic state throughout the day by optimizing key hormones and maintaining positive nitrogen retention. This is covered in greater detail in the discussion on meal frequency. For now, let's look at protein's role in the equation.

Protein must be ingested frequently throughout the day to deliver a steady supply of amino acids into the blood to be delivered to muscle tissue. This steady supply provides a positive nitrogen balance and an anabolic (muscle-

Ingredient	Casein Hydrolysate	Whey Protein Concentrate	Beef	Chicken	Fish (Tuna)
Alanine	3.00	4.82	6.02	5.44	6.05
Arginine	3.70	3.18	6.83	6.02	5.96
Aspartic Acid	6.90	12.26	10.00	8.89	10.24
Cysteine/Cystine	0.40	2.28	1.06	1.29	1.07
Glutamic Acid	20.90	15.41	16.04	14.98	14.93
Glycine	1.80	2.00	4.74	4.90	4.78
Histidine	2.90	2.41	3.22	3.10	2.96
Isoleucine	4.80	6.41	5.17	5.28	4.65
Leucine	9.10	11.60	8.85	7.50	8.12
Lysine	7.70	9.83	9.40	8.50	9.18
Methionine	2.90	2.35	2.83	2.77	2.96
Phenylalanine	5.10	3.56	4.35	3.96	3.89
Proline	10.40	6.28	4.21	4.12	3.53
Serine	5.80	6.24	4.00	3.45	4.08
Threonine	4.30	8.44	4.64	4.22	4.38
Tryptophan	1.20	1.80	1.06	1.17	1.12
Tyrosine	5.50	3.26	3.68	3.38	3.38
Valine	5.70	6.09	5.42	4.96	5.16
Critical Five Amino Acids	44.20	42.70	42.30	38.74	38.82

building) environment. If blood amino acid levels drop, your body pulls these amino acids from muscle tissue. This puts you in a negative nitrogen balance and catabolic state. Your body actually “eats away” your hard-earned muscle in order to maintain blood amino acid requirements.

This can easily occur if a highly trained athlete is not careful. Intense training depletes amino acid stores. To make matters worse, building more muscle requires more amino acids. Therefore, it is extremely important that you take in adequate amounts of protein from sources with high concentrations of essential amino acids, especially the critical five amino acids.

The whole idea behind Macrobiotic Nutrition is to give your body the tools it needs to grow muscle. Your protein intake must be staggered throughout the day to deliver a steady supply of amino acids to the blood and to provide ample nitrogen (donated by the amino acids) to shift the body into an anabolic, or muscle-building, state. Whether you derive the protein from food or supplements depends on convenience and availability. The only thing we know for sure is that at bedtime a protein drink, lower in carbohydrates and fat, is superior for several reasons. One reason is that food protein increases stomach acidity, which is not good at bedtime and may result in a feeling of heartburn. Conversely, a shake containing Probiotic-engineered protein is lower in acidic proteins and higher in critical-cluster amino acids than any other protein on the market. This provides exactly what we need at bedtime, which includes:

1. Lower insulin levels
2. Increased glucagon levels to stimulate the conversion of stored triglycerides into usable fatty acids, thereby facilitating body-fat reduction
3. Suppressed somatostatin (due to the increased glucagon) for a more efficient GH release
4. Amino acids, supplied by the delayed-release protein blend, as building blocks needed in the newly created anabolic environment

The most critical times of the day are the one-hour periods prior to and after your workout. If you want to grow, two basic aspects of these time periods must be evaluated: energy levels and recovery nutrients. A Macrobiotically-balanced meal eaten about ninety minutes prior to your workout will provide you with the blood sugar level your body depends on when trying to grind through a grueling workout. A blended-protein source with varying release rates will ensure that a steady supply of the critical five amino acids are available for energy and muscle growth.

Postworkout is also best addressed with a Macrobiotic meal or Macrobiotic meal replacement drink. Again, the low-glycemic carbohydrates contained in the meal will supply the muscle glycogen in a steady manner, while a steady source of amino acids aids in muscle recovery and an elevation in protein synthesis to stimulate muscle growth.

ABSORPTION RATES OF PROTEINS

Absorption rates are another important factor to consider to avoid falling into a negative nitrogen balance. The absorption rates of protein have become a hot topic in the supplement industry, spurred by the whey and milk protein indus-

tries in their fight for market share in protein supplements. Different proteins raise blood amino acid levels at different rates. This is termed “release rate.” Proteins can be categorized into three groups: fast release, medium release, and slow release. Which category is the most effective for athletes has been a heated debate in recent years. Each category has its own characteristics:

- **Fast release**—increases blood amino acid levels rapidly, but amino acid levels also fall rapidly. Sources: whey protein isolate, whey protein concentrate.
- **Medium release**—takes a little longer to raise blood amino acid levels and drops slower. Sources: milk protein isolate, soy isolate, soy concentrate.
- **Slow release**—provides a gradual increase and maintains blood amino acid levels for a longer duration of time. Sources: casein, whole foods.




Figure 3.1. This graph illustrates the release rates of whey, soy, and casein. As you can see, each protein source has a different release rate. Whey protein provides a quick release while soy protein provides a medium release and the casein provides a slow release of amino acids. Therefore, combining all three protein sources will provide a constant supply of amino acids to muscle tissue for optimum nitrogen retention.

Combining Fast, Medium, and Slow Proves Best

The truth is, all of these protein sources have their benefits, and ideally are best used together. Remember, if you are following Macrobiotic Nutrition properly, you will be eating 35 percent of your total calories from protein spaced throughout the day into five to seven meals every two and a half to three hours. Out of these meals, three to four should be whole-food meals and two to three should be supplements. Whole-food protein sources such as chicken provide a slow release of amino acids, due to the body's need to break down and digest the proteins. When proteins are eaten with carbohydrates and fats, the amino acids are released even more gradually. Macrobiotic whole-food meals in the 45/35/20 ratio provide a favorable release of amino acids and an anabolic hormonal environment. Protein supplements and meal replacements, on the other hand, have to be chosen carefully, but if formulated properly, supplements can actually be more beneficial than whole foods at certain times of the day.

Morning, postworkout, and before bed are all times in which the properly formulated supplement can offer a tremendous advantage. The first thing in the morning and postworkout are both times when blood amino acid levels are low, either from being in a rested state (during sleep) or from being depleted by exercise. You're in jeopardy of being in a catabolic state at these times, so you want to quickly replenish amino acids to get back to a positive nitrogen balance and shift from being catabolic to anabolic. A supplement containing fast-, medium-, and slow-release proteins such as whey, soy, and casein will provide a quick infusion of amino acids to get you back to positive nitrogen balance and will let you maintain high blood amino acid levels for many hours until your next meal.

Many of us have busy schedules and don't always have the time to sit down and eat a Macrobiotically-balanced meal, let alone cook and prepare one, so supplements can offer a quick and convenient alternative. Who has the time to prepare, cook, and eat seven meals a day? Good thing there have been advancements in supplement technology. When I was competing back in the late 1990s, these types of quality supplements didn't exist. The development of high-quality supplements allows you to get your five to seven Macrobiotic Meals in—no excuses!

Consuming a slow-release protein before going to bed helps prevent catabolism during the many hours of fasting while you sleep. Whole food provides a slow release of amino acids, which is what you want to maintain blood amino-acid levels during the night, but they can wreak havoc on your stomach. Food proteins, especially meat and dairy, are acidic and can cause stomach discomfort and heartburn during the night. A protein shake containing the whey, soy, and casein combination I mentioned earlier will have the preferred sustained-release

RESEARCH UPDATE

Combining Fast- and Slow-Release Protein Improves Nitrogen Retention

In 1997, French researcher Yves Boirie and colleagues introduced the terms “fast” and “slow” dietary proteins to describe the differences in whey and casein digestion. The higher concentrations of plasma amino acids in the three-hour period following a whey meal results in higher rates of protein synthesis. After this period, it is casein and not whey that results in higher rates of protein synthesis, a condition that is maintained for several hours.

Combining whey with casein provides both fast, high concentrations of amino acids and long-term high concentrations of amino acids to maintain muscle nitrogen retention and improve protein synthesis. *This supports Macrobiotic Nutrition’s recommendation to combine protein sources such as whey, casein, and soy for optimal protein synthesis to support muscle growth and prevent catabolism.*

Yves Boirie, et al., “Slow and Fast Dietary Proteins Differentially Modulate Postprandial Protein Accretion,”
Proceedings of the National Academy of Science 94 (1997) 14930–5.

profile, and is also less acidic and easier to digest. This precisely formulated combination is what I refer to as a Probiotic protein. A Probiotic protein is one that yields a sustained-release profile of the critical five amino acids, to optimize the blood amino acid levels conducive to an anabolic environment.


It is also important to note that a supplement taken at night before going to bed should be high in protein, very low in carbohydrates, and high in fiber, which is not the typical Macrobiotic ratio. The Macrobiotic 45/35/20 creates the ideal hormonal environment during the day, but you want a different hormonal landscape while you sleep. A Probiotic protein taken before bed will do the following:

1. Decrease insulin
2. Increase glucagon
3. Suppress somatostatin and increase growth hormone (GH)
4. Supply high levels of amino acids and maintain positive nitrogen balance for many hours

Engineering the Perfect Anabolic Protein: MHP's Probiotic Protein Blend

For an athlete, a variety of protein sources may not only be a good idea, but also a necessity. Different proteins are composed of varying concentrations of different amino acids, so a blend of protein sources, if chosen wisely, provides an even distribution and concentration of the amino acids necessary for optimal growth. For an athlete, protein demands exceed the RDA (recommended daily allowance set up by the federal government) due to the increase in muscle breakdown caused by extensive training and limited recovery. The even distribution of amino acids and their duration in the bloodstream are both of concern. Different proteins have different release rates and different rates of absorption, factors that affect how long the blood amino acid levels remain elevated.

The blend utilized in Probiolic is, in essence, a delayed-release protein. The whey in the formula possesses a fast release rate and rapidly elevates blood amino acid levels. The soy, on the other hand, is classified as a medium-release protein, and will extend the amino acid release into the bloodstream past that of the whey concentrate. The addition of calcium caseinate helps maintain an elevated blood amino acid level beyond that provided by the soy, and provides the slowest release rate of the three proteins. This protein blend provides amino acids that effectively donate nitrogen for up to three hours after the ingestion, preventing the body from slipping into a catabolic environment.



WHAT DOES ALL OF THIS MEAN TO YOUR TRAINING?

The impact your diet (in this case, the protein you consume) has on the results of your training is the key element in determining the outcome of all your efforts. Everyone knows that you need protein to build muscle, but I have given you the “master key” as far as protein consumption is concerned. The information outlined in this chapter tells you why, how much, and when to eat the right proteins, and most of all, what they will do for you. We all know that athletes have a higher demand for protein than the average individual, but how that extra protein is consumed is what will actually make or break you. The rules for success with protein intake are:

1. Consume 35 percent of your total calories from protein sources high in essential and critical five amino acids.
2. Consume different sources of protein. For example, one meal of eggs and egg whites, one meal with tuna, one meal with chicken, some red meat at one meal, and a couple of supplement meals will provide a wide variety of absorption rates and amino acid profiles. This variety will ensure that you receive a steady source of high levels of all the amino acids necessary for muscle growth. Remember, different proteins score differently in different categories, such as in the biological value (BV) category, so a variety of the highest-scoring protein sources is the best.
3. Eat five to seven meals per day, two and a half to three hours apart. Three to four of these meals should be from whole foods and two or three meals per day should be from protein supplements. Research shows that meal frequency is important to maintain nitrogen retention and improve protein synthesis.
4. Choose protein supplements wisely and take them at critical times of the day. A supplement combining whey, soy, and casein provides a quick, medium, and slow release of the essential and critical five amino acids. Optimum times for Probiolic protein supplements are:
 - a) First thing in the morning as part of a 45/35/20 Macrobiotic meal
 - b) Postworkout as part of a 45/35/20 Macrobiotic meal
 - c) Before bedtime: One-quarter gram (0.25 g) of protein per pound of body weight (body weight \times 0.25), high in fiber to further prolong amino acid release

Therefore, a 200-pound athlete would consume a 50-gram serving of Pro-bolic protein supplement high in fiber and low in carbohydrates before bedtime ($200 \text{ lbs} \times 0.25 \text{ g} = 50 \text{ g}$).

If you follow these protein guidelines, you will remain in an anabolic (muscle-building) state twenty-four/seven. When my protein intake is frequent and I am taking my bedtime shake, my muscles feel fuller and harder. I don't know if it is due to the glycogen-sparing effects, or the anabolic effects, or a combination of both, but I definitely see and feel a difference. Remember, your best twenty-three hours of growth potential each day are the twenty-three hours after you train, but you need to supply your body with the amino acid building blocks if you want to grow!



Photo by Per Bernal.

CHAPTER 4

Fat—It Is Essential!

FATS ARE AS ESSENTIAL AS ANY OTHER NUTRIENT in the human body. They are the richest source of energy, donating nine calories per gram as compared with only four calories per gram provided by both carbohydrates and protein. Dietary fat also helps you absorb the fat-soluble vitamins A, E, D, and K. Fat consumption needs to be monitored, though, because while certain sources offer health benefits, others can present serious health risks including cardiovascular disease, cancer, and obesity. Dietary fats vary in their activity and function in the body. You have probably heard the terms or concepts of “good fats” and “bad fats.” The fats you choose should come from the naturally occurring fat found in lean animal protein sources like chicken, beef, tuna, salmon, and whole eggs, along with other select sources high in essential fatty acids such as seeds, nuts, olive oil, and flaxseed oil. Even though certain animal-based protein sources contain saturated fat, selecting the leanest cuts will ensure that you keep saturated fat at a safe level. In *Macrobolic Nutrition*, 20 percent of your calories are recommended to be derived from fat. The body needs some fat, primarily in the form of fatty acids, in order to perform optimally and maximize muscle growth. When a fatty acid cannot be manufactured by the body and has to be derived from the diet, it is called an essential fatty acid, or EFA. EFAs are responsible for the formation of prostaglandins and other hormonelike substances that regulate blood pressure, fight infection, regulate growth in children, and have an anti-inflammatory effect. They are also responsible for skin and hair growth. Fatty acids are long chains of carbon, hydrogen, and oxygen all joined together. How the carbon, hydrogen, and oxygen atoms are joined together determines whether it is a saturated, monounsaturated, or polyunsaturated fat. The structure also affects their stability, shape, and how they work in the body.

Unsaturated fatty acids have carbon chains linked together by double

bonds. Saturated fats like butter and fatty acids with trans double bonds (the carbon chains are on opposite sides of the bond) like margarine tend to be solids at room temperature, while cis fatty acids (both carbon chains are on the same side of the bond) tend to be liquids at room temperature. Vegetable oil is an example of a cis fatty acid.

When the food industry artificially hydrogenates oils, they reduce the double bonds by adding hydrogen atoms. The result of this process is the creation of trans fatty acids. Trans fatty acids are what keep the oil in processed peanut butter from separating out as it does in natural peanut butter. The problem with trans fatty acids is that they reduce the fluidity of your cell membranes and actually make it harder for cells to function. Trans fatty acids are minimized in the Macrobiotic cookbook, and serve no purpose for a bodybuilder trying to maximize hormone production and muscle growth.

Essential fatty acids are very important for health and performance. The two fatty acids that are considered essential are alpha linolenic and linoleic. These two fats make up two distinct families of fatty acids; the omega-3 family, from alpha linolenic and the omega-6 from linoleic. Seeds and grains are the primary source of omega-6 fatty acids in the diet. Sunflower, safflower, and corn oils are excellent choices to increase your omega-6 intake. The American diet usually provides sufficient omega-6, but is often proportionately low in omega-3. Therefore, omega-3 fatty acids are more commonly

RESEARCH UPDATE

Trans Fats Hinder Muscle Growth

A new study describes yet another insidious effect of trans fats: muscle loss. Focusing on the effects of trans-fat ingestion in seventeen women and fifteen men, age thirty-eight to eighty-three, the study found that it interferes with the metabolism of essential fatty acids by inhibiting their conversion into eicosanoids. It turns out that some of these vital eicosanoids control protein metabolism and synthesis in the body. When the trans fats lowered eicosanoid production, protein synthesis declined, leading to amino acid loss and subsequent muscle-tissue loss.

Hubbard, R., et al., "Apparent Skeletal Muscle Loss Related to Dietary Trans Fatty Acids in a Mixed Group of Omnivores and Vegetarians."
Nutrition Research 23(2003): 651-658.

supplemented, and can be derived from dark green leafy vegetables. Even though fish oil is not high in the primary omega-3 fatty acid, alpha-linolenic acid, it is exceptionally high in the omega-3 derivatives EPA (eicosapentaenoic acid) and DHA (docosahexaenoic acid). These two fats are essential for neurological development and brain function. In one study performed on rats, it was proven that a diet high in saturated fats impaired learning, while a diet high in the omega-3 fatty-acid derivatives DHA and EPA heightened mood and improved cognitive function. As we age, levels of DHA decrease in the brain. So it is imperative to pay additional attention to this fat if you want to stay sharp.

Another function of the omega-3 family of essential fatty acids involves insulin secretion and insulin sensitivity. This is extremely important to Macrobiotic nutrition, since our emphasis is on low-glycemic carbohydrate sources and on optimizing insulin levels throughout the day. In Chapter 2, I explained in detail how insulin affects our bodies. Studies show that omega-3 fatty acids actually increase membrane permeability of the beta cells in the pancreas (where insulin is made), thus allowing for insulin to be released easier. This helps prevent the oversecretion of insulin (hyperinsulinemia or Syndrome X). Since the muscle cells also have a phospholipid or fat layer with insulin receptors on it, the omega-3 fatty acids increase the cells' insulin-binding affinity and sensitivity, enhancing glucose transport across the cell membrane. This change results in increased glycogen storage and leads to fuller, harder muscle and increased energy storage.

Joint pain and inflammation are major concerns for many people, especially athletes. Joint pain and inflammation can sideline your ability to train and perform optimally and hinder your progress. In a study by M. J. James, et al., published in the *Journal of Biological Chemistry* in 2000, omega-3 fatty acids were shown to decrease COX-2 (cyclooxygenase), decrease the enzymes associated with joint degradation, and decrease the cytokines (messenger chemicals) associated with inflammation. The COX-2 enzyme is induced by inflammation, and is involved in most diseases associated with chronic inflammation. Joint pain and inflammation as well as joint deterioration will be highest when this enzyme is turned on. The omega-3 derivatives EPA and DHA have exhibited strong anti-inflammatory activity and help shut down the COX-2 enzyme. The studies actually substantiate the anticatabolic effects that the omega-3 family has on joint tissue. Since joint pain and degradation can plague bodybuilders, it's critical to make smart choices (like the omega-3s) for your fat intake.

In all of the studies I have examined, the anti-inflammatory effect of the

NEW SCIENCE UPDATE

Study Proves Eating Unsaturated Fats Instead of Saturated Fats Helps Reduce Body Fat

Piers and coworkers conducted a randomized crossover study of eight overweight or obese men (aged 24 to 49 years with BMI 25.5–31.3 kg/m²). These subjects followed two diets for four weeks each to determine whether substitution of saturated fat with monounsaturated fat affects body weight and composition. The male subjects were provided with all food and beverages during the study period. The percentage total energy from fat for each test diet was as follows: On the saturated fatty acid-rich diet, 24 percent saturated fat, 13 percent monounsaturated fat, and 3 percent polyunsaturated fat; on the monounsaturated-rich diet, 11 percent saturated fat, 22 percent monounsaturated fat, and 7 percent polyunsaturated fat.

During the study period, the researchers kept track of body composition, blood pressure, energy expenditure (resting and postprandial metabolic rates, substrate oxidation rate, physical activity), serum lipids, the fatty acid profile of serum cholesteryl esters, and plasma glucose and insulin concentrations. The researchers used some of these measurements to determine if the subjects followed the diets, for example by keeping track of cholesterol levels and the fatty acid composition of serum cholesteryl esters in their blood.

The study revealed that the men had a lower body weight (–2.1 kilograms) and lower body-fat mass (–2.6 kilograms) at the end of the monounsaturated fat-rich diet as compared with values at the end of the saturated fat-rich diet. No significant differences were detected in energy or fat intake, energy expenditure, substrate oxidation rates, or self-reported physical activity. The researchers concluded that substituting dietary saturated fat with unsaturated fat (predominantly monounsaturated fats) can cause a small but significant reduction of body weight and body-fat mass without a significant change in total energy or fat intake. This study provides more evidence that “all calories are not created equal”!

Piers L.S., Walker K.Z., Stoney R.M., Soares M.J., O’Dea K. “Substitution of Saturated with Monounsaturated Fat in a 4-week Diet Affects Body Weight and Composition of Overweight and Obese Men.” *British Journal of Nutrition* 90, No. 3 (Sept 2003): 717–727.

omega-3 family is best achieved when the omega-3 fatty acids are derived from fish oils. Incorporating salmon into your Macrobiotic plan provides an excellent source of these fatty acids. In the case of the omega-3 fatty acids, supplementing with some MaxEPA (fish oil) would benefit the bodybuilder, ensuring ample intake of this EFA. Fatty acids in the cell membrane also gives the body the ability to produce hormonelike substances called eicosanoids that produce localized effects in any tissue or organ in the body. These short-acting “hormones” have a direct effect on preventing inflammation and on the blood flow into and out of tissue, which is important to maximize nutrient delivery to growing muscle cells.

Cholesterol is another hormone precursor manufactured from dietary fat in the body. I know that cholesterol has gotten a bad rap in the last few years, but the truth is you need some of it to serve as the starting material for all steroid hormones (like testosterone) in your body. Cholesterol also makes up a great deal of the brain matter and is used in vitamin D production and in the formation of bile, which is used to emulsify fats. Dietary intake of cholesterol is not necessary because your liver can manufacture it from the low levels of saturated fats you receive from your lean animal-protein sources. Cholesterol forms the basic ring structure molecule that your adrenal gland will take to manufacture corticosteroids, testosterone precursors, estrogens, and progesterone.

As you know, the formation of hormones is extremely important for muscle growth. Testosterone is an androgenic hormone responsible for male characteristics such as facial hair, body hair, deepened voice, male sexual functions, and muscle growth. All guys, especially bodybuilders, want to have as much testosterone floating around in their bodies as possible. Studies show that you must take in a minimum of 20 percent of your total calories from fat (yes, the same amount recommended by Macrobiotic Nutrition) to maintain normal testosterone synthesis. Saturated fat is the preferred form of fat for the biosynthesis of cholesterol to testosterone. This important need for saturated fat is another reason why I believe in consuming protein sources like red meat and whole eggs. Maybe this is why most hardcore bodybuilders and power lifters say red meat makes them strong.

Since intense training can actually weaken your immune system, it is worthwhile mentioning the effect that fat intake has on immune function. T cells, B cells, and macrophages all have a strong presence of fatty acids in their cell membranes, so the EFAs are essential for the proper development of these immune cells. Omega-3 fatty acids also produce eicosanoids, compounds that have been shown to elicit a strong inflammatory response to injury and infection. Eicosanoids increase fever and swelling along with

immune globulin E (IGE), all-important defenses of the immune system. Higher body temperature kills bacteria, while increased sensitivity to pain inhibits you from moving an area that is injured (which may lead to further damage), and IGE increases the production of allergic antibodies. These all serve as first-line defenses of your body's immune system, and help ensure efficient recovery from intense training.

This makes a great argument for Macrobiotic Nutrition's requirement to derive 20 percent of your total daily calories from fat. Fat actually helps slow digestion of a meal, delaying the release and absorption of nutrients into the blood. This delayed release helps lower overall glycemic response to a meal and keeps blood sugar from spiking and insulin release stable. Slowing down digestion of a meal also prolongs the release of amino acids from protein into your bloodstream, improving nitrogen retention and keeping you in an anabolic state. Fat also increases the release of CCK (cholecystokinin), a hormonelike cytokine that sends a message to the brain that the body is no longer hungry. Couple the anti-inflammatory, hormone-producing, and insulin-modulating effects with the digestive benefits of fat, and it is a no-brainer why fat is essential in your diet. Just be sure to keep track of how much fat you consume, because at nine calories per gram, the calories can add up pretty quickly. Sorry, this isn't the Atkin's diet, so forget about the crisp bacon sizzled in butter!

FACTS ON FAT

Trans fats: Damaged fats formed when oil is hydrogenated or foods are fried. They may raise your risk of cancer and heart disease.

Hydrogenated oil: Oil that has had extra hydrogen added to keep foods like margarine firmer and longer lasting at room temperature. Hydrogenated oils are found in most supermarket margarines, cookies, doughnuts, cakes, and many other processed foods. This type of fat has been linked to an increased risk of cancer and heart disease.

Saturated fat: Type of fat found in abundance in butter, lard, red meat, and whole milk. Saturated fat has been blamed for increasing the risk of obesity, heart disease, and cancer. These fats get their name from the fact that they are saturated with hydrogen atoms.

Unsaturated fat: Type of fat found in plant oils and considered less harmful than saturated fats. However, experts now feel that unsaturated fats rich in

omega-6 fatty acids can increase your risk of chronic disease if you don't consume enough omega-3 fatty acids. They are called unsaturated because chemically, their molecules have room for the addition of more hydrogen. Oils high in unsaturated fats that some experts recommend include olive oil, hemp oil, peanut oil, and macadamia oil.

Omega-3 fatty acids: Type of fat found in fish, hemp seed, pumpkin seed, and flax oil. Experts believe we should consume more of these fats, since they reduce inflammation and produce a wealth of health benefits.

Omega-6 fatty acids: Group of fatty acids abundant in the American diet. While these fats are not necessarily harmful, they may foster heart disease and other chronic illnesses when consumed in large quantities without sufficient omega-3 fats. Corn oil, safflower oil, and soybean oil are rich in omega-6 fats.

Monosaturated fatty acids: The type of fatty acids found in olive oil and nuts like peanuts, macadamias, almonds, and walnuts. These fats are believed to lower the risk of heart disease and cancer.



Photo by Per Bernal.

CHAPTER 5

Water— The Nutritional Paradox

YOU'VE SEEN PICTURES OF PROFESSIONAL BODYBUILDERS working out. Expressions of intense determination convey the pinnacle of athleticism, only to be overshadowed by the sheer size of the muscles being bombed. This level of intensity is magnified by the beads of sweat forming on their bodies from the amount of effort they are putting into every rep to pack on muscle mass.

Have you ever considered what single nutrient makes all of this possible? What nutrient enables energy metabolism, protein synthesis, and virtually all other chemical reactions used by the body for performance? Most average bodybuilders overlook this cardinal nutrient because of its simplicity. But you are not the average bodybuilder. You are on the path to becoming the best and achieving your maximum human performance. With that in mind, let us consider this supreme compound, known simply as water.

Many overlook water as a nutritional staple. It is a nutritional paradox; this simple compound has no caloric value from carbohydrates, proteins, or fats; nor does it contain any vitamins or minerals. It is almost ironic; pure water is a nutrient that clearly has no nutritional value as far as macro- or micronutrients are concerned. Yet, despite this fact, it is so essential to the performance of the human body that if even 2 percent of your body weight in water was lost, you would begin to lose the edge in the gym. In light of this, water cannot be overlooked in Macrobiotic Nutrition.

Certain minerals, such as the electrolytes, need the presence of water to function properly. Potassium (K^+) and sodium (Na^+) are both cations (positively charged ions). Because of their charge, they can interact with the partially negative oxygen part of H_2O . Thus, both potassium and sodium are dissolved in the presence of H_2O . Without proper levels of hydration, the functions of these two electrolytes will falter. The importance of these two electrolytes with reference to performance is covered in Chapter 8.

All body functions that include enzymatic activity need water to function properly, including glucose utilization, nutrient delivery, and waste transport. Due to its chemical composition, water really is the universal solvent; it allows all body functions to proceed smoothly. I could go into great detail about how water is crucial to each of these avenues of performance, but to be practical, I won't do so. Instead, let's briefly view the additional benefits to the body that water provides, aside from the direct biochemical aspects of this nutrient.

Water contained within the cell is called intracellular fluid. This fluid accounts for 30 to 40 percent of a person's body weight, and is typically high in the macrominerals potassium and phosphate. The water between the cells, which is high in the concentrations of sodium and chloride, is called interstitial fluid. Interstitial fluid is a main component of extracellular fluid, which in total accounts for 20 percent of the total body weight of a person. The body will adjust its water intake and excretion to maintain the proper balance of these fluids.

WATER'S IMPACT ON PERFORMANCE

To illustrate the sheer magnitude of water and its "weight" in the body, let's look at muscle tissue. Muscle tissue is made up of roughly 70 percent water. With a decrease of only 2 percent total water weight, nerves that direct the function of the muscle tissue will be compromised, with noticeable decreases in muscle control and strength. Your strength training will decrease by about 21 percent and your aerobic capacity will drop by 48 percent with a water weight loss of about 2 percent. Protein synthesis and anticatabolism are in effect when muscle cells are properly hydrated. So, water is absolutely necessary for your performance in the gym and your growth outside the gym.

To further illustrate the importance of water, consider the following functions. Water flushes lactic acid from muscle cells, serves as the key component of lubricating fluids for the joints, and serves as a shock protector in the joints and vertebral disks. In addition, water stabilizes body temperature through perspiration. Clearly, proper hydration is the fundamental cornerstone for upping the muscle mass in anyone.

MAINTAINING HYDRATION LEVELS IN THE BODY

Dehydration is avoided when one responds to the thirst sensation. Yet in some instances, dehydration has a rapid onset due to various factors. This could include a willful deprivation of water such as when wrestlers stop drinking to make their weight class or bodybuilders stop drinking to lose subcutaneous water to appear more cut. Excessive water loss also occurs due to vomiting,

dehydration, diarrhea, and in the case of the athlete, sweating. The symptoms of dehydration vary with the loss of body water weight (BWW). The thirst sensation, weakness, loss of appetite, and slight discomfort are associated with a 1 to 2 percent loss of water. If the athlete avoids drinking water and loses 3 to 4 percent of BWW, the performance of the athlete will suffer, as described earlier. When the loss of BWW rises to 5 to 6 percent, the body will have trouble regulating its temperature and it will increase its respiratory rate. When it reaches the point of 7 to 10 percent loss of BWW, muscle spasms, exhaustion, decreased ability to maintain balance, and collapse will result. Normally, the athlete who follows a proper diet program can avoid these conditions. Yet endurance athletes should be aware of the conditions that develop when water levels drop in the body.

The best source of water is obviously water, but it is also found in all foods. Foods such as watermelon, strawberries, nonfat milk, spinach, broccoli, and lettuce all contain 90 to 99 percent water. In addition to the water found in foods, the body actually generates water as a byproduct of energy metabolism. When carbohydrates, proteins, and fats are broken down, their carbon and hydrogen components combine with oxygen to form carbon dioxide and water.

Water is lost through many avenues, so it must be replaced constantly. The most influential variable for water loss is sweating, so Macrobiotic Nutrition suggests drinking at least 4 ounces of water per 100 kcals (kilogram calories) burned during exercise. This may seem too little to support proper body function, but this is the bare minimum that must be consumed when compensating for lost fluids caused by excessive perspiration. Excessive perspiration will result from an intense workout and the temperature of the gym. To ensure peak performance, I always recommend sipping on cool (not ice-cold) bottled water throughout the duration of your workout. I recommend cool, as opposed to cold, because cool water is absorbed more quickly than cold water. This is pivotal for keeping energy production at its maximum and lactic acid saturation to a minimum, which means better workouts and better results. You will also want to drink often throughout the day, since the athletes who follow Macrobiotic Nutrition will want to keep their bodies at the optimum level of hydration so they can train at their maximum capacity. Remember, even with Macrobiotic Nutrition, you'll have to hit the weights "heavy and hard"!

Water is essential to life, and it is also essential to maximum human performance, so let's not overlook its importance. You train hard to be your best, so it would be shameful to screw it up by not doing something as simple as drinking enough water!

CHAPTER 6

Macrobiotic Meals to Muscle Mass

NOW THAT YOU KNOW HOW THE RIGHT SELECTIONS of carbohydrates, proteins, and fats react in the body, we need to look at what happens when you eat these food sources in a 45/35/20 Macrobiotic meal. I want you to understand how this method of eating is going to make you bigger and leaner starting with digestion and ending with the nutrients going into your muscle cells. To begin this chapter, I will show how food is mechanically digested in the mouth and chemically broken down in the gastrointestinal tract, and how different hormones shuttle the nutrients released in the digestive process right to your growing muscle cells—all of this while making sure not to overload these processes and promote any storage of body fat.

THE PROCESS OF DIGESTION

When we eat whole foods, the nutrients (carbohydrates, protein, and fat) we need to get from them are not readily available until the food undergoes the process of digestion. Digestion starts in the mouth with chewing, the mechanical breakdown and mixing of foods. Larger particles are broken down into smaller ones. Smaller particles increase the surface area of the food particles so the digestive enzymes can break down the food more thoroughly. After you chew, digestion begins with the release of salivary amylase, an enzyme that is part of the saliva in your mouth. This enzyme is the first step in the breakdown of carbohydrates. After you chew your Macrobiotic meal, swallowing is achieved by a wave of muscular contractions called peristalsis. The muscles of the esophagus propel food down into your stomach like an ocean wave moving through water. As the food approaches the stomach, muscles relax and the food passes through a ringlike structure of muscle called the esophageal sphincter. This muscular ring prevents food from being regurgitated into the esophagus.

The chewed-up food is now in the stomach. The stomach is an organ

that performs three main mechanical functions related to digestion. In the first function, the upper part of the stomach must relax to be able to accept large portions of food and liquids. The second function involves mixing the food and liquid with the digestive enzymes (like hydrochloric acid) produced by the lining of the stomach. This lining produces both the enzymes that break down the food and the hormones that control hunger and digestion. This is the stage of digestion where protein foods begin to be broken down into peptides and amino acids, to be absorbed and used by your muscles. At this point, all nutrients have begun to be broken down, getting closer to the point where your body can actually utilize them. The third function of the stomach is to empty the partially digested contents into the small intestine to undergo further digestion. The rate at which the stomach empties is controlled by several factors. The protein and fat content of the meal, and the overall physical state of the meal (whether it is solid or liquid) determine the speed at which the stomach empties. A fatty meal may take up to three and a half hours to empty out of the stomach, whereas a liquid shake low in fat will empty in about a half hour. The other contributing factor that will slow gastric emptying is the fiber content of the meal. Fiber also slows down transit of the food through the intestine, allowing for more thorough absorption of the nutrients. This is important and will help keep blood sugar and blood amino acid levels where they need to be. Ideally, a Macrobiotic meal's protein, fat, and fiber content allows for optimal absorption and a gradual release of nutrients.

Inside the small intestine, digestive enzymes from the pancreas, the liver, and the intestine break down the carbohydrates further, to result in glucose molecules that the body can readily absorb into the blood. Absorption of these glucose molecules increases the blood sugar. Remember, Macrobiotic Nutrition's low/moderate-glycemic carbohydrates take longer to break down to glucose, and fat and fiber slow down gastric emptying, so the glucose from the carbohydrates are not released in one big load into the intestine. This prevents any spikes in your blood sugar and prevents insulin from shooting up to high. In a sense, by adding protein, fat, and fiber, you are lowering the glycemic index of the carbohydrates in the meal.

Let's take a look at what happens to the other nutrients in the small intestine. Fats are emulsified by some of the pancreatic enzymes and the bile, which is produced by the liver and stored in the gallbladder. The emulsified fats are easily converted into the fatty acids the body needs for efficient absorption. The pancreas also produces some proteolytic enzymes that finish off protein digestion. The protein foods you ate are now amino acids, dipeptides, and tripeptides, all of which can be absorbed in the small intestine and actively transported into the blood, where they circulate to the muscle cells

that crave them. As the amino acids are absorbed into the blood from the intestine, the blood amino acid level goes up, stimulating muscle growth and signaling an increase in glucagon levels. Glucagon mobilizes stored body fat for energy utilization.

Insulin and glucagon are essential for transport of the nutrients to the cells, but there are three other hormones that play a key role in the digestion of your Macrobiotic meal. They are cholecystokinin (CCK), gastrin, and secretin. Gastrin is the hormone that signals the stomach to produce the acid that breaks down the protein you eat into peptides. It is also necessary for the growth of the stomach lining, the intestinal lining, and the lining of the colon. Secretin is the hormone that signals the pancreas to start sending out the digestive enzymes such as pepsin to break the protein peptides into amino acids. Secretin also triggers the release of bicarbonate into the intestinal tract so the amino acids are absorbed efficiently without being denatured (broken down) by the acid from the stomach. In the presence of fat, secretin signals the liver to produce bile. This is its major role in the digestion of fats. CCK causes the pancreas to produce the enzymes that make up the pancreatic juice released into the small intestine. CCK also signals the gallbladder to release bile for the breakdown of the fatty part of the meal. Additionally, CCK is a messenger hormone that tells the brain that we are no longer hungry.

UTILIZATION OF MACROBOLIC NUTRITION

Now that you know how a Macrobiotic meal is broken down into the nutrients we need to support muscle growth and optimize hormonal levels, let's take an in-depth look at how Macrobiotic Nutrition creates the ideal anabolic and metabolic environment to stimulate gains in lean mass.

The 45/35/20 ratio of select nutrition is designed to have all the macronutrients work in synchronization. This is why it is referred to as the 45/35/20 lean-mass equation. This ratio of nutrients will feed your muscles throughout the day and keep muscle-building and fat-burning hormones in check.

Low-glycemic carbohydrates, protein, and fat in a 45/35/20 ratio will provide a gradual and controlled release of insulin. As you recall, you want to avoid high spikes of insulin, but you want a meal to elicit some insulin because it helps shuttle glucose, amino acids, and other nutrients into your muscles. This steady, controlled release of insulin pushes the high concentrations of critical amino acids provided by the protein sources in the meal to the muscles. This constant supply of amino acids will keep you in positive nitrogen balance, and continually supply muscle-building amino acids. Aside from helping slow digestion and the release of glucose and amino acids into your bloodstream, fats improve insulin sensitivity and are used for the production and formation of hormones.

Macrobiotic meals are designed to keep your body running optimally for approximately three hours. Here's a list of what a Macrobiotic meal provides:

1. Sufficient supply of carbohydrates to meet energy demands
2. Controlled insulin release to shuttle amino acids and glucose to muscle tissue
3. Controlled insulin release to prevent the transformation of triglycerides into body fat
4. Steady supply of amino acids from quality protein sources to maintain positive nitrogen balance
5. Raised glucagon to increase fat burning
6. The necessary supply of fat (especially EFAs) to support hormone production, prevent inflammation, and slow digestion to control blood sugar and amino acid release
7. A thermogenic effect

After a Macrobiotic meal, all these great things are going on at once. Your body is running optimally and you are in anabolic state. However, this is where meal frequency becomes important. A Macrobiotic meal only fuels your body for so long. As I said, you'll be running optimally for about three hours. If you want to keep your body running like a finely tuned muscle machine, you're going to have to refuel with another Macrobiotic meal.

Okay, so now your body is running on all cylinders. You're eating every three hours, fueling your body with the right stuff. All systems are in check, you're as anabolic as a bottle of testosterone. Amino acids are being shuttled to your muscles, stimulating new muscle growth, and fat is being emulsified and burned throughout the day.

Nighttime is a different story. You need to keep insulin low and keep nitrogen high during the nighttime fasting hours. At night, you slam down a Probiotic sustained-release protein shake before going to bed, again creating the ideal anabolic environment for muscle growth and fat burning. Let's look at our nighttime checklist:

1. Steady supply of critical five amino acids from sustained-release protein to maintain nitrogen balance during sleep
2. Low insulin levels due to very low-carbohydrate content of shake
3. Elevated human growth hormone (HGH) due to low insulin levels during sleep

The Thermic Effect of Food

By now, you are probably tired of hearing that a calorie is not just a calorie. I cannot emphasize this enough. I've already covered the difference in impact of carbohydrate, protein, and fat sources on hormones and metabolism even though they contain the same amount of calories. And I think that you now understand the importance of choosing the right macronutrient source. But there is yet another factor to consider when estimating calories, and that is food's thermic effect. It may sound strange to hear that food is thermogenic. When we speak of thermogenics, the first thing that usually comes to mind are fat burners like ephedrine, synephrine, and caffeine. You probably wonder why I would be talking about any kind of thermogenic effect in a book devoted to packing on mass. Well, understanding the effect that the food you eat has on energy is essential to figuring out how many calories you need, so you will pack on lean muscle and not become fat. What we now refer to as the thermic effect of food (TEF) was previously called the specific dynamic action. It is the increase in heat, or oxygen consumption, in your body after you eat a meal. Calories are a measure of heat, so an increase in heat production actually means an increase in calories burned.

Each nutrient (fats, proteins, and carbohydrates) has a different effect on the TEF. Additionally, different types of fats, carbohydrates, and protein will have either a greater or lesser effect on TEF. Basically, the harder nutrients are to digest, the higher the TEF. First, let's talk about carbohydrates and the effect different types have on thermogenesis. A meal that contains an elevated percentage of high-glycemic carbohydrates, like white bread, increases blood glucose and insulin levels more rapidly and has a lower TEF than a meal that contains low-glycemic carbohydrates, like oatmeal. Low-glycemic carbohydrates (especially high-fiber foods) increase thermogenesis and the oxidation of the carbohydrates is more efficient. This would explain why people tend to lose body fat while restricting their carbohydrate intake to mainly low-glycemic foods. A study conducted by Schwartz, et al., illustrated the differences in the TEFs of low and high-glycemic carbohydrates.

In a study performed in 1974, subjects consumed protein, glucose (a high-glycemic carbohydrate), or a combination of the two. The results illustrated a 17 percent increase in basal metabolic rate (BMR) in the protein group, a 14 percent increase in BMR in the carbohydrate group, and a 17 percent increase in the group fed protein and carbohy-

drate. This study illustrates that the right combination of carbohydrates and protein will have the same thermogenic effect on your body that plain protein will have, which is why Macrobiotic Nutrition emphasizes the types of food you eat and the proportions and times you eat them. You need to take advantage of this easy way for your body to utilize excess calories. To top it off, the more muscle you put on, the higher this number gets. That's right, you read that correctly—the more muscle you pack on, the more metabolically efficient your body will get.

Table 6.1 illustrates the results of a Swiss study by J. M. Schwartz, et al., reported in the *American Journal of Clinical Nutrition* in 1989, which found that lean men experienced a higher TEF than obese men. This proves my point. The leaner you are, the more calories you need to sustain your body weight and prevent losing muscle.

TABLE 6.1. TEF IN LEAN VERSUS OBESE MEN

Parameter	Lean	Obese
Body weight (kg)	57	84
BMR (cal/hr)	69	74
TEF	+13%	+5.2%

This point was further proven in a study published by Segal in *Endocrinology and Metabolism* where subjects with a higher lean body mass were found to have a higher TEF from meals eaten pre- and postworkout. One thing is for sure, an athlete needs to do two things to maximize the TEF: (1) increase his or her lean body mass, and (2) eat frequently, approximately every two and a half to three hours, so you do not slow down your metabolism.

It's an interesting point, and one that is familiar to dieters: Eat fewer calories, and you will slow down your metabolism, causing fat loss to cease. This is not the case with Macrobiotic Nutrition. Our "muscle-friendly" ratio of calories will prevent slowing your metabolism, while simultaneously packing on lean, hard muscle. Dramatically reducing calories has been shown to decrease muscle mass, which is undesirable. We have already hypothesized that a higher lean body mass (more muscle) actually increases the thermic effect of food and BMR (the amount of energy expended in a resting state).

In a study conducted by Westerterp and published in the *European Journal of Clinical Nutrition*, diets containing different proportions of

carbohydrates and fat were evaluated for their effect on thermogenesis. The researchers found that meals containing carbohydrates, protein, and fat produced the greatest thermogenic effect, while meals high in fat produced a very low TEF. This study further supports the ratios prescribed in Macrobiotic Nutrition.

While all of this is very interesting, protein is still the best source for increasing TEF. In a 1984 study performed by Jequier, fat was found to have a 2 to 3 percent TEF, and carbohydrates a 6 to 8 percent TEF, while protein elicited a 30 to 40 percent TEF. While the protein numbers may appear high, in all of my research, the lowest TEF for protein I found was 25 percent. This means that if you eat 200 grams of protein (which is 800 calories), your body actually requires anywhere from 200 to 320 calories to utilize it. This is unbelievable! The increased TEF of protein is linked to the increase in nitrogen turnover, caused by an increase in protein metabolism. Basically, we want to utilize protein's higher TEF to maximize caloric expenditure at certain times of the day when we are less active. At night, meals higher in protein will generate a higher TEF, allowing your body to expend more calories while at rest. Couple the protein with a little fat and we delay the gastric emptying of the protein, thus stretching this TEF out over a longer period of time. You see, you want your body to have a demand for calories, even while you sleep. Without carbohydrate floating around as glucose in your blood, your body switches to the triglycerides stored in your fat cells. The protein further increases this caloric demand, while increasing glucagon, which signals the release of the stored fat (as discussed in Chapter 2).

So, the Macrobiotic ratio also maximizes the thermogenic effect of food. The prescribed 35 percent of total calories from protein in combination with 45 percent low-glycemic carbohydrates and 20 percent unsaturated fats eaten in frequent meals throughout the day will require a good amount of calories to digest and metabolize. The meal frequency of every two and a half to three hours not only provides a continual thermogenic effect, but it also boosts your metabolism. By increasing thermogenesis and boosting your metabolism, you optimize your BMR significantly.

Additionally, Macrobiotic Nutrition's recommendation to supplement with a Probiotic protein high in fiber and low in carbohydrates allows you to create the ideal hormonal and metabolic environment to build muscle and burn body fat while you are sleeping. How awesome is that?!

4. Increased fat burning due to raised glucagon, energy requirements of basal metabolic rate (BMR), and thermic effect of last meal

Yes, everything is in check and you're primed for growth—even while you're sleeping. When you wake up, it starts all over again. You see what's happening? You're a muscle-building, fat-burning machine. Your body is running optimally twenty-four/seven and it's in the highest possible anabolic state.

RESEARCH UPDATE

Carbohydrates and Essential Amino Acids Are Anabolic

Strength training builds muscle tissue by triggering tension receptors in the muscle and by creating small injuries to the fibers. In response, the body creates special repair units called satellite cells that are incorporated with the contractile fibers of the muscle cells and makes those cells larger. Also, tension triggers the movement of amino acid into the cells, which makes new protein (muscle). Muscle growth or atrophy (cell shrinkage) is a balance between protein synthesis and protein breakdown. After a heavy workout, protein breakdown usually exceeds protein synthesis—at first. Gradually, protein synthesis exceeds breakdown if the right fuels and hormones are available. University of Texas scientists found that feeding a carbohydrate/essential amino acid supplement after weight training slowed protein breakdown and accelerated protein synthesis. This is more evidence of the importance of taking a protein supplement after your workout.

Rasmussen, B.B., K.D. Tipton. "An Oral Essential Amino Acid-Carbohydrate Supplement Enhances Muscle Protein Anabolism After Exercise." *Journal of Applied Physiology* 88, No. 2 (Feb 2000): 386–392.

IT DOESN'T GET ANY BETTER THAN THIS!

On Macrobiotic Nutrition, you can't help but get big and lean. You see, being in an anabolic state throughout the day, day after day, is going to allow you to continually pack on muscle mass. And if you monitor your calories correctly, you *will* continually burn body fat. In fact, as your muscle mass increases, your caloric requirements will increase. It takes additional calories to maintain muscle mass. So, as your muscle mass increases, your BMR and caloric intake also increase.

What Is BMR?

Basal metabolic rate (BMR), also referred to as resting metabolic rate (RMR), is the amount of calories your body needs to support normal body functions to maintain “life” at total rest. Lifestyle would cover the calories needed to support the energy required for your occupation and other “nonexercise” voluntary activities. Exercise energy (calorie) requirements are those needed or expended specifically during your workout. Your BMR, lifestyle, and exercise routine are the three biggest determinants used in calculating daily caloric needs.

SAY NO TO CARDIO—FOR MAXIMUM GROWTH AND STILL GET RIPPED!

Early in this book, I said that Macrobiotic Nutrition could get you big and ripped without cardio. I’m not a big fan of cardio. Besides the fact that I think running on a treadmill or stepper like a gerbil on an exercise wheel is boring, I feel it is counterproductive to building mass. If you train hard and follow Macrobiotic Nutrition, there is no need for cardio! I know that is a bold statement, but let me explain how this is possible.

First, let’s look at the reasons why people do cardio in the first place. They want to:

1. Burn calories
2. Increase metabolism
3. Improve cardiovascular health
4. Increase cardiovascular endurance

True, these all look like great reasons to do cardio, but isn’t building rock-hard freaky muscle what really matters to you? Unless you are an endurance athlete, I see absolutely no need for you to be doing cardio. You can get the same or even better results from intense weight-training sessions. An hour of intense weight training will burn more calories and boost your metabolism better than an hour of jogging on the treadmill, with equal benefits to your cardiovascular health.

Cardio does absolutely nothing to stimulate muscle growth. In fact, it will hinder muscle growth. Aerobic activities like jogging, cycling, and the stepper

deplete your body of glycogen, amino acids, and other precious muscle-building nutrients. Depleting glycogen and amino acids from your muscles can cause your body to spiral into a catabolic state, eating away hard-earned muscle with every pointless step. Long aerobic sessions have also been shown to raise levels of the catabolic hormone cortisol.

I probably still haven't convinced you not to do cardio. And, if so, you forgot to consider one important new factor—Macrobiotic Nutrition. Remember, if you're following Macrobiotic Nutrition, your body is running optimally. Through your diet, you have optimized your metabolism and hormones to efficiently build muscle and burn body fat. You have calculated your caloric needs and each calorie is being utilized to perform its proper function—*every calorie counts!* If you are following Macrobiotic Nutrition properly and training intensely, you will not have to rely on cardio to burn body fat.

Exceptions to the No-Cardio Rule

(Okay, I'll make two exceptions.)

1. If you participate in a sport that also requires endurance conditioning. As I've said before, you have to train for your sport. Cardiovascular training is essential for some sports, like boxing, wrestling, and basketball, because they require high levels of endurance. These athletes usually need to train at a 70 to 80 percent target heart rate. The good news for you is that Macrobiotic Nutrition is the best program to support your endurance needs and still optimize muscle mass, even with high levels of aerobic exercise. Just remember to increase your calories to compensate for the extra calories burned during your exercise sessions.
2. If you're starting the Macrobiotic Nutrition program with more than 15 percent body fat, you may include cardio in an effort to get body fat down more quickly. But don't overdo it. Remember, you should be using the appropriate caloric requirements chart with the reduced calorie adjustments for people with more than 15 percent body fat.

My recommendation for you is forty-five minutes at a moderate pace (60 to 70 percent of target beats per minute) first thing in the morning on an empty stomach, three to five days per week. To figure your target heart rate, see Figure 6.1 and Table 6.1.

Following the diet, accurately counting the calories you consume, and weight training intensely and consistently are ultimately what's going to help you lose body fat and achieve a lean, muscular physique.

Figure 6.1. Target Heart Rate




TABLE 6.1. TARGET HEART RATE

AGE	Low Impact for Fat Burning 60%–70%		Intermediate 70%–80%		Advanced 80%–90%	
	Beats/min	Beats/10 sec*	Beats/min	Beats/10 sec*	Beats/min	Beats/10 sec*
to 19	121–141	20–24	141–161	24–27	161–181	27–30
20–24	119–139	20–23	139–158	23–26	158–178	26–30
25–29	116–135	19–23	135–154	23–26	154–174	26–29
30–34	113–132	19–22	132–150	22–25	150–169	25–28
35–39	110–128	18–21	128–146	21–24	146–165	24–28
40–44	107–125	18–21	125–142	21–24	142–160	24–27
45–49	104–121	17–20	121–138	20–23	138–156	23–26
50–54	101–118	17–20	118–134	20–22	134–151	22–25
55–59	98–114	16–19	114–130	19–22	130–147	22–25
60–64	95–111	16–19	111–126	19–21	126–142	21–24
65–69	92–107	15–18	107–122	18–20	122–138	20–23
70–74	89–104	15–17	104–118	17–20	118–133	20–22
75–79	86–100	14–17	100–114	17–19	114–129	19–22
80–84	83–97	14–16	97–110	16–18	110–124	18–21
85+	81–95	14–16	95–108	16–18	108–122	18–20

*To use six-second counts, multiply by 10 to get beats per minute.

MACROBOLIC MOMENTUM: GETTING BETTER WITH TIME

The longer you are on Macrobiotic Nutrition, the more efficient your body will become. It takes more calories to maintain muscle than fat. As lean body mass increases, your caloric requirements throughout the day will increase. This means that as Macrobiotic Nutrition shifts your body composition to a higher percentage of muscle and a lower percentage of body fat, you will need to feed it more carbohydrates, protein, and fat. I like to call this phenomenon “Macrobiotic Momentum.” Basically what’s happening is that your ability to pack on lean muscle accelerates over time if you stay on Macrobiotic Nutrition, because your body becomes so metabolically efficient. So, in essence, the bigger and leaner you get . . . the bigger and leaner you get.

You’ve probably seen this Macrobiotic Momentum phenomenon happen firsthand and just never realized it. If you look at the progression of some professional bodybuilders, you will see that there was a stage early in their careers where they made some good, steady progress. But then, all of a sudden, they went through a period of explosive growth. This stage of explosive growth occurs because of the accumulation of muscle mass supported by Macrobiotic Nutrition. There is no reason why you can’t experience this same explosive growth yourself.



Photo by Per Bernal.

CHAPTER 7

Guide to Estimating Caloric Requirements

MACROBOLIC NUTRITION HAS TAUGHT YOU about the powerful effects of food. You've learned that all calories are not created equal and the importance of choosing the right ratio and sources of macronutrients. You've also learned that the 45/35/20 lean-mass equation will optimize hormones and metabolic efficiency.

Now that you know how to optimize metabolic efficiency, you must learn the next critical step in achieving lean mass. Like everything in bodybuilding, it comes down amounts: how many sets, how many reps; how much creatine, how much glutamine? And most important, how much food or how many calories should you be consuming? As I explained earlier, the simple premise that caloric intake minus caloric expenditure will determine weight loss or weight gain does hold merit. Macrobiotic Nutrition differs because its influence on metabolic efficiency increases your BMR, allowing you to consume more food "calories" to support muscle growth while burning fat. However, it is still crucial to adjust your caloric intake correctly to achieve your goal of changing body composition. If you eat more calories than you expend (burn), your body fat will increase even if your metabolic efficiency improves. Usually, but not always, the goal for bodybuilders is to decrease body fat while increasing muscle. Bodybuilders who are relatively lean often want to maintain body-fat levels while increasing muscle mass, and strength athletes often want to increase both body fat and muscle mass.

While Macrobiotic Nutrition's caloric distribution of 45/35/20 is the ideal ratio for muscle growth, power, and performance, the total caloric intake must be adjusted for each individual and their goals. Also, keep in mind that as Macrobiotic Nutrition increases muscle mass and your metabolic efficiency, your caloric needs will continually increase, as more calories will be needed to support this new mass.

So, the question remains, how many calories should you consume to reach your goal? Pinpointing any individual's exact daily caloric needs is very difficult, and many factors must be considered. Not only do these factors vary from person to person, but some factors can vary in the same person from day to day.

As explained earlier, your BMR, lifestyle, and exercise routine are the three biggest determinants used in calculating daily caloric needs. However, Macrobiotic Nutrition also takes into account your personal goal, which is, after all, what Macrobiotic Nutrition is all about. This unique feature is referred to as the Metabolic goal variable (MGV). So, in figuring out your caloric needs, we use your MGV in place of your BMR.

MACROBOLIC CALORIC EQUATION

Macrobiotic Nutrition has devised a simple, yet extremely accurate formula for you to use to determine your caloric needs to achieve your goal. We refer to this formula as the Macrobiotic caloric equation.

$$\begin{aligned} &(\text{Bodyweight} \times \text{MGV}) + (\text{Bodyweight} \times \text{Lifestyle}) \\ &+ (\text{Exercise Expenditure}) = \text{Total Caloric Requirements} \end{aligned}$$

Let's walk through this equation, so you can see how easy it will be for you to calculate the Macrobiotic Nutrition calories you need to achieve your specific goal. The Macrobiotic calorie equation uses MGV, lifestyle, and exercise to figure out your caloric needs, so let's define each of them.

Macrobiotic Goal Variables (MGV)

The metabolic goal variable (MGV) is broken down into three possible goals: gain, maintain, or lose. Of course, each of these goals has gaining muscle mass as a constant, but adjusts for your desired body fat.

MGV: Gain, Maintain, Lose

- Gain Body Fat/Increase Muscle
- Maintain Body Fat/Increase Muscle
- Lose Body Fat/Increase Muscle

Gain: The goal is to increase muscle mass and increase body fat. This is usually the goal of strength athletes or "hard gainers." Strength athletes sometimes like to have a slightly higher percentage of body fat covering their muscle mass. "Hard gainers" are those of you who have a genetic predisposition to being lean with low body fat and have a hard time putting on weight. For the gain variable, we use a multiplier of 13. So, in the calculation: $\text{MGV} = 13$.

Maintain: The goal is to increase muscle mass and maintain the current percentage of body fat. Over the long haul, this should be your status. As a bodybuilder, you should strive to get your body fat fairly low. I recommend staying between 6 and 10 percent body fat year-round, and then dropping a few percentage points in your last ten to twelve weeks of contest preparation. This will allow you to make gains and look big, hard, and full year-round. It will also allow you to coast into your contest looking massive and diced, rather than flat and emaciated. For the maintain variable, we use a multiplier of 10. So, in the calculation, $MGV = 10$.

Lose: The goal is to increase muscle and decrease body fat. This is probably the most common goal since most people want to be bigger and leaner. Now, you can! So, in the calculation, $MGV = 7$. If you are currently carrying a large amount of body fat (over 15 percent), use 6 as your multiplier— $MGV = 6$.

MGV Multipliers	
Gain:	13
Maintain:	10
Lose:	7
High percentage of body fat to lose:	6

Lifestyle

Lifestyle includes all forms of daily voluntary activities except exercise. Your occupation, hobbies, recreational activities, and daily chores all come into play, as they all require calories to accomplish. Lifestyle has been broken down into three categories: Sedentary, Moderate/Active, and Very Active.

Sedentary: Believe it or not, many professional and top amateur bodybuilders fall into this category. Remember, lifestyle does not include exercise. You fall into this category if you don't have a full-time occupation or you do very little voluntary or physical activities outside of the gym. I once fell into this category, where all day was spent eating, sleeping, training, or thinking about training. Man, I miss those days.

Moderate/Active: You fall into this class if you are a student or have an occupation that requires little physical activity. You also don't regularly participate in hobbies or sports that require a lot of activity. This is probably the category for most people. This category requires more calories than the sedentary group for your lifestyle outside of the gym.

Very Active: You fall into this group if your job is very labor-intensive, like landscaping, construction, or any manually physical occupation, or if you participate in sports or hobbies that are physical and active.

Lifestyle Multipliers	
Sedentary:	3
Moderate:	5
Very Active:	10

Exercise Expenditure

I have always believed that weight training at a fairly high intensity level is the best way to stimulate muscle growth, elevate testosterone and growth hormone, and boost metabolism. I feel that doing cardio to burn fat is counterproductive to mass building. Intense weight training with Macrobiotic Nutrition takes care of the body fat. Your training time and energy should be applied to stimulating growth with an intense weight-training program.

Under this recommended training regimen, I calculate your caloric expenditure during exercise by taking the length of workout (number of minutes) and multiplying it by 10. So, if you train for one hour (60 minutes), your calculated exercise expenditure would be 600 calories ($60 \times 10 = 600$).

$$\text{Exercise expenditure} = \text{Length of workout (minutes)} \times 10$$

- If you do low-impact cardio (heart rate of more than 65 percent to less than 80 percent of maximum) such as slow jogging, walking, or stepper, add those minutes to the length of your workout time in the calculation.

For example:

60 minutes of weight training

30 minutes of low-impact cardio

$$(60 + 30) \times 10$$

$$90 \times 10 = 900$$

Exercise expenditure = 900

- If you do high-impact cardio (heart rate of more than 80 percent of maximum) like running, take the number of minutes of high impact cardio and multiply it by 12, and add that amount to the exercise expenditure total.

For example:

45 minutes of weight training

45 minutes of intense cardio

$$(45 \times 10) + (45 \times 12) =$$

$$450 + 540 = 990 \text{ total exercise expenditure}$$

Macrobiotic Calorie Equation in Action

Let's say an athlete weighs *180 pounds*, has a *moderate lifestyle*, and trains four days a week. His training sessions are *sixty minutes* of intense weightlifting. His *goal* is to gain muscle while losing his existing body fat. To properly determine his calorie requirements, he will use Macrobiotic Nutrition's caloric equation.

Training Days

$$(\text{BW [180]} \times \text{MGV [lose]}) + (\text{BW [180]} \times \text{lifestyle [moderate]}) \\ + (60 \text{ min} \times 10) = \text{calories}$$

$$(180 \times 7) + (180 \times 5) + (60 \times 10)$$

$$1,260 + 900 + 600 = \mathbf{2,760}$$

Nontraining Days

$$(\text{BW [180]} \times \text{MGV [lose]}) + (\text{BW [180]} \times \text{lifestyle [moderate]}) = \text{calories}$$

$$(180 \times 7) + (180 \times 5)$$

$$1,260 + 900 = \mathbf{2,160}$$

So, as you can see, this athlete should consume 2,760 calories on his workout days and 2,160 calories on his nonworkout days.

Breaking Down Total Calories for Macrobiotic Nutrition's 45/35/20 Ratio

Figuring out the breakdown of carbohydrate, protein, and fat calories to meet Macrobiotic Nutrition's 45/35/20 ideal ratio for muscle growth, power, and performance is simple, once we know the required total calories. Let's use our example of the 180-pound athlete with a moderate lifestyle, and figure out his daily requirement for each macronutrient. These calculations are all based on the 45/35/20 lean-mass equation. Remember, 45 percent of the total calories are from carbohydrates, 35 percent are from protein, and 20 percent are from fat.

Let's figure out our 180-pound athlete's daily carbohydrate, protein, and fat requirements on his training days. This athlete's total daily calorie requirements are 2,760 calories. So, to figure out the total calories of each macronutrient, use these formulas.

$$\begin{array}{l} \text{Carbohydrates: Total calories} \times 45\% = \text{calories from carbohydrates} \\ 2,760 \quad \times 0.45 = \mathbf{1,242} \\ \mathbf{1,242 \text{ calories from carbohydrates}} \end{array}$$

$$\begin{array}{l} \text{Protein: Total calories} \times 35\% = \text{calories from protein} \\ 2,760 \quad \times 0.35 = \mathbf{966} \\ \mathbf{966 \text{ calories from protein}} \end{array}$$

$$\begin{array}{l} \text{Fats: Total calories} \times 20\% = \text{calories from fat} \\ 2,760 \quad \times 0.20 = \mathbf{552} \\ \mathbf{552 \text{ calories from fat}} \end{array}$$

So, this athlete's daily caloric requirements on training days would be:

$$\begin{array}{r} \text{Carbohydrates} = 1,242 \\ \text{Protein} = 966 \\ \text{Fat} = \mathbf{552} \\ \hline \mathbf{2,760} \text{ total calories} \end{array}$$

As you can see, these totals agree with our previous calculation for total calorie intake.

Meal Frequency and Caloric Distribution

As you learned from the previous chapter, eating every two and a half to three hours is important for staying in an anabolic state and optimizing your metabolism. So, once you've figured out your total daily caloric intake, you need to break that total down into meal values throughout the day. The 180-pound athlete with a moderate lifestyle in our previous example would divide his total calories into five or six meals on both his training and nontraining days. Let's assume this athlete eats six meals per day.

$$\begin{array}{l} \text{Training days} = 2,760 \text{ (total calories)} \div 6 \text{ (meals)} = 460 \text{ calories per meal} \\ \text{Nontraining days} = 2,160 \text{ (total calories)} \div 6 \text{ (meals)} = 360 \text{ calories per meal} \end{array}$$

Converting Calories into Grams

To convert calories into grams, you must remember that 1 gram of carbohydrate contains 4 calories and 1 gram of protein contains 4 calories, while fat, being the most calorie dense, contains 9 calories per gram. To figure out the grams of carbohydrates and protein you must divide the total calories of each by 4, and to calculate the grams of fat you need, you must divide the total fat calories by 9.

For example:

Carbohydrates:

$$\begin{aligned} \text{Total carbohydrate calories} \div 4 &= \text{grams of carbohydrates} \\ 1,242 \text{ calories} \div 4 &= 310.5 \text{ grams of carbohydrates} \end{aligned}$$

Protein:

$$\begin{aligned} \text{Total protein calories} \div 4 &= \text{grams of protein} \\ 966 \text{ calories} \div 4 &= 241.5 \text{ grams of protein} \end{aligned}$$

Fat:

$$\begin{aligned} \text{Total fat calories} \div 9 &= \text{grams of fat} \\ 552 \text{ calories} \div 9 &= 61.3 \text{ grams of fat} \end{aligned}$$

So, our 180-pound athlete would have the following daily nutritional requirements on his workout days:

	Total calories:	2,760
Total calories from carbohydrates (total grams):	1,242	(310.5 g)
Total calories from protein (total grams):	966	(241.5 g)
Total calories from fat (total grams):	552	(61.3 g)

To convert these daily totals into requirements per meal, you would divide each number by the number of meals you eat each day. Since we decided this athlete eats six times a day, we would divide these daily totals by six.

$$\begin{aligned} \text{Total calories} \div 6 &= \text{calories per meal} \\ 2,760 \div 6 &= \mathbf{460} \\ &460 \text{ calories per meal} \end{aligned}$$

$$\begin{aligned} \text{Total daily carbs} \div 6 &= \text{carbohydrates per meal} \\ 1,242 \text{ calories} \div 6 \text{ or } 310.5 \text{ g} \div 6 &= \text{carbohydrates per meal} \\ 207 \text{ calories or } 51.7 \text{ g} &= \text{carbohydrates per meal} \end{aligned}$$

$$\begin{aligned} \text{Total daily protein} \div 6 &= \text{protein per meal} \\ 966 \text{ calories} \div 6 \text{ or } 241.5 \text{ g} \div 6 &= \text{protein per meal} \\ 161 \text{ calories or } 40.25 \text{ g} &= \text{protein per meal} \\ \\ \text{Total daily fat} \div 6 &= \text{fat per meal} \\ 552 \text{ calories} \div 6 \text{ or } 61.3 \text{ g} \div 6 &= \text{fat per meal} \\ 92 \text{ calories or } 10.2 \text{ g} &= \text{fat per meal} \end{aligned}$$

Calculations Made Simple

If all of this confuses you, don't worry about it! I have included Macrobiotic Caloric Requirements Tables that will do the calculations for you, based on your current body weight, goal, lifestyle, and workout schedule, so you will know the exact amount of calories, carbohydrates, protein, and fat you need to maximize muscle mass and target your goals. (You'll find these tables in Appendix A.)

Critique Your Physique

If you want to be as accurate as possible when figuring out your caloric requirements, you need to take a serious assessment of your current condition. Remember, muscle mass requires a lot more calories to maintain than body fat. For example, a 200-pound person with 5 percent body fat (190 pounds lean body mass) requires more calories than a 200-pound person with 20 percent body fat. (160 pounds lean body mass). Critique your physique honestly, because it will help you reach your goal faster.

As explained earlier, there are two different caloric multipliers for the goal of losing body fat: A multiplier of 7 is used for people with less than 15 percent body fat, and a multiplier of 6 is used for people with more than 15 percent body fat. This allows you to calculate your caloric intake more accurately if your goal is to lose body fat.

The goal to maintain body fat is usually for people with 6 to 10 percent body fat, who want to stay at their current body-fat level but increase muscle mass. This group uses a multiplier of 10.

Macrobiotic's caloric calculator is as accurate a system as you can find, but it is not an exact science. Your percentage of body fat and your overall metabolism may require you to slightly adjust the recommended calories. Don't make quick changes in your caloric intake, however. Use the recommendations for at least three weeks, and monitor your progress. At first you may gain or lose a few pounds of body fat. You may want to increase your calories if you feel

you are losing body fat too fast. Don't panic if you feel you are gaining body fat; you may have underestimated the amount of body fat you are carrying and be consuming too many calories. You can do one of two things at this point (after the first three weeks):

1. Keep your calories the same, because the frequent Macrobiotic meals will start to boost your metabolism. Keep in mind that even if you put on a few pounds of body fat, you most certainly are also gaining muscle mass. The increase in muscle mass will demand these calories by increasing your BMR, and the body fat will start to melt away.
2. Adjust the calories slightly in the Macrobiotic caloric equation by cutting back to the calorie requirements of a body weight of 10 pounds less than your current body weight. So, if your body weight is 190 pounds, go by 180 pounds when calculating your figures. This will give you a lower daily caloric intake.

Honestly critiquing your body and assessing your progress are critical aspects of any program. The only way you are going to benefit is to make proper assessments and target your goals. The longer you follow the Macrobiotic Nutrition program, the more in tune you will become with your body. You know what your goals are, you know what foods to eat, now make every calorie count!



Photo by Irvin J. Gelb.

CHAPTER 8

The Important Role of Micronutrients

AS MACROBOLIC NUTRITION'S NAME SUGGESTS, its primary focus is on the macronutrients: carbohydrates, protein, and fat. However, micronutrients also play a vital role in metabolic functions and performance. Therefore, it is important that you gain a general understanding of how vitamins and minerals influence important functions of the body.

This chapter covers the two classes of vitamins (water soluble and fat soluble) and the two classes of minerals (macrominerals and microminerals) and their roles in the body with regard to performance and health. Proper intake levels of the individual vitamins and minerals are discussed, and foods containing high amounts of the particular micronutrients are identified.

VITAMINS

For the elite athlete's performance, vitamins are essential. In fact, an athlete's performance will drop considerably if certain vitamins are deficient in his or her diet. Read on to learn what you need to know about these important micronutrients.

Water-Soluble Vitamins

Water-soluble vitamins are unique in that they are absorbed directly into the blood. Because of this factor, they are able to move freely throughout the body, and once they reach the cell, they can circulate inside its water-filled compartments. Since water-soluble vitamins are extremely mobile, they are also easily excreted from the body. The kidney, whose function is to monitor the blood that flows through it, detects excess amounts of water-soluble vitamins and properly disposes of them. In cases of extreme excess, which is usually due to improper supplementation, there will be certain side effects that may become severe.

The B vitamins are the most crucial water-soluble vitamins that aid in the

overall performance of the athlete. While vitamins in general do not yield energy like the macronutrients, the B vitamins actually assist in the utilization of that energy. The B vitamins thiamine, riboflavin, niacin, pantothenic acid, and biotin are all units of compounds called coenzymes. Coenzymes are small organic molecules that are closely associated with certain enzymes, and that aid in their function. Coenzymes are so important that the enzymes cannot function at all without their presence. The other B vitamins are B₆, folate, and B₁₂. B₆ assists enzymes that metabolize amino acids, while folate and B₁₂ help in the multiplication of cells.

Energy-Releasing Vitamins

Each of these B vitamins assists, in one way or another, with the production of energy. As you will see, their roles are so critical that even a small deficiency will be detrimental to your performance.

Thiamine (Vitamin B₁). This B vitamin is the vitamin component of the enzyme thiamine pyrophosphate (TTP). TTP's function in the body is the conversion of pyruvate to acetyl CoA, which is essential to the metabolism of energy. Acetyl CoA enters the tricarboxylic acid (TCA) cycle, which is a series of reactions for the production of energy, responsible for the eventual formation of ATP (the energy molecule). In addition to its direct assistance in the production of energy, thiamine also resides on specific sites on nerve-cell membranes, directly affecting the activities of the nerves, and subsequently, the adjacent tissues (muscles).

If you eat enough food to adequately meet your energy requirements, you are probably meeting your thiamine requirements, too. The average intake of thiamine in both the United States and Canada meets the recommended amounts.

Natural sources of this vitamin can be found in all of the food categories, including breads, vegetables, fruits, milk, legumes, and meats. Of these foods, pork products, such as lean ham and lean pork chops, contain the most thiamine. One can also get adequate amounts of thiamine from enriched, fortified, or whole-grain products. However, muscle weakness can result if B₁ requirements are not met, because of its role in the conversion of pyruvate to acetyl CoA and the production of ATP.

Riboflavin (Vitamin B₂). The role of riboflavin is similar to that of thiamine. Riboflavin becomes the coenzymes flavin mononucleotide (FMN) and flavin adenine dinucleotide (FAD) in the body, both of which release energy from the macronutrients. FAD picks up two hydrogen ions from the TCA

cycle during the metabolism of the macronutrients and its subsequent release of energy. These two hydrogen ions are then delivered by FAD, which is now FADH₂, to the electron transport chain. These two hydrogen ions are “pumped” through the electron transport chain and eventually help produce ATP. In fact, for every FADH₂ that passes through the electron transport chain, two ATPs are generated.

Most people in the United States and in Canada get enough of this nutrient from their food and either meet or exceed the established riboflavin recommendations. Some of the better food sources of riboflavin include milk, yogurt, eggs, and liver. Liver is the best source of riboflavin, with a 4-ounce portion of fried liver containing (245 kcals) in about 4.68 milligrams.

Niacin (Vitamin B₃). The name niacin actually is shared by two distinct chemical structures; nicotinic acid and nicotinamide (niacinamide). The body has the ability to easily convert nicotinic acid to nicotinamide, which is the most prevalent form of niacin found in the blood.

There are two coenzyme forms of niacin: nicotinamide adenine dinucleotide (NAD⁺) and nicotinamide adenine dinucleotide phosphate (NADP⁺). These two coenzymes are essential in the energy production reactions of the macronutrients, fat and carbohydrates. The other compound that releases its energy with the aid of these two coenzymes is alcohol. NAD⁺ follows the same basic function as FAD. It picks up a hydrogen ion, as NADH, and delivers it to the electron transport chain, thus producing ATP.

Niacin is found in a variety of food sources, including enriched breads, cereals, and nuts. An interesting fact about niacin is that it can be synthesized by the body from the amino acid tryptophan. Since tryptophan is an amino acid, any diet that contains adequate amounts of protein, from both a variety of plant sources and animal sources, will not be deficient in this energy-releasing vitamin/coenzyme. Typically, most people in the United States and Canada have dietary protein intakes that are enough to avoid a niacin deficiency.

Biotin. This B vitamin is the last in the line of B vitamins whose function is solely the release of energy. The coenzyme form of biotin is required to carry a carbon to the 3-carbon pyruvate making it the 4-carbon compound oxaloacetate. Oxaloacetate is then combined with acetyl CoA, enabling the TCA cycle to continue producing products needed to manufacture ATP. The other essential role of the coenzyme form of biotin is gluconeogenesis. As mentioned in Chapter 2, gluconeogenesis occurs when the body does not have enough glucose to carry out metabolic functions. When the body is in this state, it starts metabolizing certain amino acids, and then fat, for the production of ATP.

Gluconeogenesis can have a negative impact on muscle building and performance. Macrobiotic Nutrition's 45/35/20 lean-mass equation can prevent this condition. However, gluconeogenesis is critical not only for performance, but also for brain function and other organ functions in the absence of glucose.

Biotin can be synthesized in the gastrointestinal tract by certain bacteria, but the amount of the synthesized biotin that is actually absorbed is not fully known. An elite athlete should rely instead on a variety of foods as established by Macrobiotic Nutrition and on supplementation to achieve an adequate amount of biotin.

Pantothenic Acid. This is the only B vitamin that serves many purposes, which range from the production of energy to the formation of steroid hormones in the body. In the formation of energy, pantothenic acid actually functions as a part of coenzyme A. This compound forms the acetyl CoA in the "bridge" between the end stage of glycolysis (the breakdown of glucose into two pyruvate molecules) and the beginning stage of the TCA cycle. It is essential in the formation of neurotransmitters, which are chemicals produced by neurons used to carry signals to other neurons and to nonneuron cells like skeletal muscle.

Due to its prevalence in food, pantothenic acid deficiencies rarely occur in a person who has a reasonable diet, but they will result in lethargy among other conditions. A diet following the principles of the Macrobiotic Food Guide Pyramid (see Appendix B) has a good foundation in whole grains, which are rich in pantothenic acid. In addition to whole grains, beef and poultry provide good amounts of pantothenic acid.

The Anabolic B Vitamins

Vitamins B₆, B₁₂, and folic acid directly affect your body's growth potential. They cover mostly protein synthesis, but they can serve in other functions such as steroid hormone activity. As you read, you will continue to learn the importance of Macrobiotic Nutrition and how it relates to maximizing your performance.

Pyridoxine, Pyridoxal, and Pyridoxamine (Vitamin B₆). This B vitamin is like niacin in that it has three chemical forms: pyridoxine, pyridoxal, and pyridoxamine. Any one of these three forms of B₆ can be converted into pyridoxal-5-phosphate (P5P). P5P plays a significant role in muscle growth, because P5P is the coenzyme that transfers amino groups. This is absolutely essential in amino acid metabolism. When the cells need to manufacture proteins, P5P takes freely available amino groups (which are provided by a posi-

tive nitrogen balance due to an abundance of protein intake) and synthesizes all eleven nonessential amino acids. For example, as mentioned earlier, the conversion of tryptophan to niacin requires P5P. Vitamin B₆ is unique among the water-soluble vitamins in that it is stored, to an extent, in muscle tissue.

In support of this role, recent studies have shown that B₆ has a positive influence on steroid-hormone activity. Studies have also shown that a large dosage of B₆ does not improve muscular strength or physical endurance, but I must point out that this is in supplemental form. The role of this vitamin is protein synthesis, and an athlete can easily consume his or her RDA of B₆ from a well-balanced diet. Meats such as chicken breast, turkey, and beef are excellent sources of B₆ since it is stored in muscle tissue. So, if you are eating meat sources such as these, you should be getting enough B₆ to carry out these functions

The Dynamic Duo (Folic Acid and Vitamin B₁₂)

These two anabolic B vitamins are considered the dynamic duo because the role of each in supporting DNA synthesis and tissue growth is rendered inactive without the presence of the other.

Folic Acid. This B vitamin also goes by the names folate, folacin, and pteroylglutamic acid (PGA). The coenzyme form of folic acid, tetrahydrofolate (THF), is required to transfer the one-carbon compounds that are produced during metabolism. This process aids in the conversion of vitamin B₁₂ into one of its coenzyme forms, which plays a role in the synthesis of DNA that occurs in all rapidly growing healthy cells. In food, folic acid is bound with a chain of amino acids called polyglutamate. The body hydrolyzes the polyglutamate to monoglutamate and other glutamates with the aid of enzymes on the surface of the intestinal tract. The monoglutamate is then attached to a methyl group for absorption. At this point, the folate is in the body, but in its inactive state. The coenzyme form of B₁₂ removes the methyl group from the monoglutamate, making the folic acid active. If B₁₂ was not available to remove the methyl group, the folic acid would be inactive in the cell and unable to support DNA synthesis and cellular growth.

Folic acid directly affects the formation of new cells and protein synthesis, so a deficiency would lead to impaired tissue growth. It is, therefore, a crucial vitamin for everyone, especially the elite athlete. One can achieve an adequate amount of this key vitamin by including a good selection of leafy green vegetables, legumes (kidney beans, black beans), and folic acid-rich fruits (such as oranges) in the diet.

Vitamin B₁₂ (cyanocobalamin). As mentioned before, vitamin B₁₂ is cen-

tral for the activation of folic acid, and it also removes the methyl group from the monoglutamate. The methyl group released by folic acid bonds to B₁₂, thus forming the coenzyme form of B₁₂. Methylcobalamin and B₁₂'s other coenzyme, adenosylcobalamin, are key participants in the synthesis of new cells, because both folic acid and B₁₂ are used in the synthesis of DNA and RNA.

Vitamin B₁₂ can also be classified in its own category because it is the only vitamin that is almost exclusively found in products derived from animals. B₁₂ deficiencies rarely occur in people with a well-balanced diet, because a person should be able to meet her or his B₁₂ needs with a reasonable consumption of animal products.

Vitamin C/Ascorbic Acid, the Secondary Antioxidant

Vitamin C has a unique role in the body. It serves two main purposes: as an antioxidant and as a cofactor. Vitamin C is also known as ascorbic acid, and for good reason. Ascorbic acid prevents the condition known as scurvy. It does this through its acidic properties. The two hydrogen (H⁺) ions from the hydroxyl groups on ascorbic acid have the ability to be donated to (taken up by) free radicals in the body. Free radicals are molecules that have unpaired electrons, and because of this characteristic, they can cause oxidative damage to body tissue. Since ascorbic acid has two H⁺ ions, it can help minimize oxidative damage due to free radicals. Thus vitamin C plays a key role in the body, protecting body tissue from oxidative stress by balancing the production of free radicals with the body's ability to manage them.

Unlike the B vitamins, which are coenzymes, vitamin C is a cofactor. A coenzyme is a compound that is made up of several different substances, but a cofactor can help an enzyme in a chemical reaction alone. When vitamin C is present with iron, it helps with the formation of collagen, the fibrous protein structure that is found virtually everywhere in the body. The iron works as a cofactor in the hydroxylation (the addition of OH groups) of two key amino acids, proline and lysine, which become both hydroxyproline and hydroxylysine and facilitate in the bringing together of the collagen fibers. The matrix of the bone upon which minerals are deposited is made up of collagen, and it is present in the walls of our arteries as well as in scars, which are the result of the buildup of collagen. Vitamin C has other crucial functions that influence health, performance, and body composition. It is a cofactor in the formation of carnitine, a compound that transports long-chain fatty acids into the cells' mitochondria for the production of energy during metabolism, and it is important in the production of the thyroid hormone thyroxine, whose chief function is to increase the rate of metabolism, thereby raising energy levels and fat burning.

The most noticeable early sign of a vitamin C deficiency is seen in the maintenance of blood-vessel integrity, resulting in bleeding gums around the teeth and dotted hemorrhaging due to pinpoint subcutaneous bleeding. These symptoms occur because of malformation of collagen due to a lack of hydroxylation of either proline or lysine to their hydroxyl forms. In the absence of hydroxylation, the collagen fibers weaken and lose integrity. As a result, the conditions mentioned before will appear. One can easily avoid the unpleasant effects of this deficiency by consuming a diet rich in citrus fruits, strawberries, red bell pepper, and kiwi.

The Fat-Soluble Vitamins

The fat-soluble vitamins differ in a number of ways from the water-soluble vitamins. Vitamins A, D, E, and K (ADEK) are all insoluble in water, thus they require a substance that makes them available for absorption. Your body produces a substance called bile, which is central in the absorption of fats in the intestines. Since ADEK is absorbed in the presence of fat, ADEK is transported with fat. All fat is transported in the lymphatic system, which is a system like blood vessels that branch out through the entire body.

If there is an excess of any one of the ADEK vitamins, the body will store it in one of two body tissues. The liver and adipose tissue will store ADEK and when the body needs any one of the ADEK vitamins, it can obtain them from the body stores. This process has its pros and its cons. The obvious benefit is that you could consume less food than what is required to meet your RDA for ADEK because you can rely on your body stores. One could potentially do this for several days, weeks, or even months before seeing any sign of deficiency. All someone would have to do is ensure that over time they consume the average daily intakes of the vitamins. The potentially harmful drawback to this fact is that if one is not careful, the fat-soluble vitamins can accumulate to toxic levels since they are not excreted as easily as the water-soluble vitamins.

In the following discussions, the function of each fat-soluble vitamin in relation to peak physical performance and optimal health is established. In addition, the levels at which deficiencies can be avoided and where you can find the best dietary sources are addressed.

The Anabolic Fat-Soluble Vitamins

Vitamin A. Vitamin A comes from a family of chemicals called the retinoids. This family of retinoids consists of three different chemicals: retinal, retinol, and retinoic acid. Animal sources of vitamin A contain compounds called retinyl esters, which are converted into the compound retinol in the intestines. The compounds found in food derived from plants are called carotenoids. Some of

the carotenoids such as alpha, beta, and beta-cryptoxanthin carotene all have vitamin A activity in the body. Different cells have different needs for the specific retinoids and each form of vitamin A performs a specific function that the other two cannot do. The body can actually convert retinol into retinal and vice versa, but once any one of these two forms get converted to retinoic acid, it cannot be converted back.

For example, retinol is used in the support of reproduction and it is the major form of vitamin A that is transported and stored in the body. Retinal plays a vital role in vision by maintaining a clear cornea and takes part in the conversion of light into nerve impulses. It is also an intermediate in the conversion of retinol to retinoic acid. Retinoic acid functions much like a hormone in regulating cellular differentiation in developing embryos and in growth. Vitamin A has a role in the process of osteoclast, or the dismantling of bone. This is a natural process that must occur during bone remodeling, which takes place during bone growth. The enzymes that aid osteoclast cells need vitamin A to assist them in taking away parts of the bone that are not needed.

Some vitamin A deficiencies are apparent in people with night blindness. This is actually the first clear indication that a person may be running out of vitamin A stores in his or her body. The retina does not have enough retinal to regenerate pigments that are bleached by the light. Another severe result of vitamin A deficiency is irreversible blindness.

Toxicity develops usually when a person supplements too much vitamin A in the form of the retinoids. If you rely on a healthy, balanced diet, this is not likely to happen. Although in cases where someone would consistently eat foods that are high in vitamin A, toxicity would be a concern, too.

The healthiest way to obtain vitamin A is through the diet. Some of the best dietary sources of vitamin A are any animal-based food sources. Beef liver is the best animal source of vitamin A since liver stores a good amount of any one of the fat-soluble vitamins. Pumpkin is the best vegetable source of vitamin A since any plant that has a deep orange or a rich yellow color contains abundant amounts of carotenoids.

Vitamin D. Vitamin D is unique among all of the other vitamins, since it not an essential nutrient. Essential nutrients are nutrients that can't be synthesized by the body, yet vitamin D can be synthesized with enough sunlight and an intermediate version of cholesterol. Vitamin D plays a critical role in body development and maintenance over one's life span.

Vitamin D is really a member of a team of nutrients that regulate the bone-making process. Yet vitamin D has a key role in that it raises blood concentrations of both calcium and phosphorus. It does this by first enhancing the

absorption of the two minerals from the gastrointestinal tract. It also enhances the reabsorption of these two minerals from the kidneys, and it regulates their mobilization from the bones into the blood.

Since vitamin D is necessary for the absorption of calcium, if a deficiency in this vitamin occurs, a deficiency in calcium will occur. Osteomalacia and osteoporosis are two common effects of lack of exposure to the sun and, of course, a lack of calcium in the diet. If you rely on getting added vitamin D from the diet, toxicity is unlikely to happen. If you supplement vitamin D, you could run into some potentially serious conditions. Since the role of vitamin D is to raise the blood concentration of calcium, too much vitamin D will deposit in soft tissue. This leads to stones in the kidneys and contributes to calcification of blood vessels, such as the major arteries of the heart and lungs, which will lead to death.

Few foods contain vitamin D, which is understandable since we can manufacture it. To ensure that everyone gets enough vitamin D, milk is fortified with this essential vitamin. Other sources of vitamin D are found in veal, beef, liver, and egg yolks.

Vitamin E, the Primary Antioxidant

Vitamin E is only one of two vitamins that act as antioxidants. Yet what separates vitamin E and vitamin C is that vitamin E functions only as an antioxidant. The antioxidant function of vitamin E is to prevent the multiplication of free radicals from other free radicals, as in the case of an athlete's intense training that results in the overproduction of potentially harmful free radicals. Vitamin E does this by donating one of its own H^{+} ions (electrons) to the free radical, thereby neutralizing it. Unfortunately, vitamin E, by this point, is neutralized itself; however, vitamin C can restore vitamin E back to its active form. A diet that is abundant in these two antioxidants will drastically minimize oxidative stress placed on the body by an athlete's training.

Vitamin E naturally occurs as a family of tocopherols. The four forms of tocopherols are alpha, beta, gamma, and delta. Yet out of all the tocopherols, alpha tocopherol is the only one with direct vitamin E activity in the body. The other tocopherols aren't as easily converted into the alpha form.

A prolonged period of inadequate vitamin E consumption leads to issues with the retina of the eye and to neuromuscular dysfunctions that involve the spinal cord. Because of this, clear signs of deficiencies will include a reduction in one's ability to see, and the diminished control of muscles, resulting in poor coordination and reflexes. Treatments of vitamin E can reverse these effects. Toxicity is rare and if it does occur, it is not as serious as the toxic effects of vitamin D or vitamin A. If toxicity does occur, an interference with the blood-

clotting effects of vitamin K will occur, and one can hemorrhage. To ensure that you get enough vitamin E to avoid deficiencies and toxicity, consume a diet that has a good base in food that is adequate to meet these goals. Seeds, nuts, and various vegetable oils like safflower, peanut, olive oil, and canola oil are all fantastic sources of vitamin E and the unsaturated fats, which also have their benefits for performance and health.

Vitamin K, the Coagulant

Vitamin K is like vitamin D in the sense that it can be synthesized in the body. It cannot be synthesized directly by the body, but bacteria in the gastrointestinal tract synthesize vitamin K, and the body can absorb it in the presence of fat. Vitamin K has, quite possibly, the most important role in the body. With a deficiency of the other fat-soluble vitamins, one can live with a minimum of direct threats to one's life. Vitamin K is used for the activation of several key proteins that work along with calcium in the formation of blood clots. With a deficiency of this crucial vitamin, a simple paper cut could lead to severe hemorrhaging.

Like vitamin D, vitamin K can become deficient due to several factors that are not related to inadequate dietary intakes. Secondary deficiencies occur due to other factors. For example, in the case of vitamin K, a decreased secretion of bile will lead to a decreased absorption of fat and the fat-soluble vitamins. A deficiency may also occur because of the interaction of certain drugs with vitamin K, and with its synthesis and its activity. Antibiotics will kill the bacteria in the intestines that are responsible for synthesizing vitamin K, and certain anticoagulant drugs will interfere with the functions of vitamin K.

Toxicity is not common if vitamin K is acquired from the diet. Supplementing vitamin K may lead to conditions such as jaundice—a condition in which bilirubin, a bile pigment, spills over and gives a yellowish hue to the skin and the whites of the eyes.

As previously mentioned, vitamin K is synthesized by the bacteria that naturally inhabit the intestinal tract. Yet they can only produce about half of the vitamin K we need; a good diet can provide the rest. Significant sources of vitamin K can be found in the liver, which is the best source of the fat-soluble vitamins. Other good dietary sources of vitamin K are any green leafy vegetables, cabbages, eggs, and fruits.

MINERALS

Minerals are the other group of micronutrients the body requires. They are in a completely different class from the other micronutrients for several key reasons, including each mineral's chemical makeup, their availability through

absorption from food, the interactions that take place between two or more minerals, and their roles in the body. Since minerals are a dynamic factor in a person's performance, it is essential that we review their functions in the body. In addition to discussing their functions when they are taken in the proper amounts, I describe the symptoms experienced when toxicity is reached. I also indicate which foods provide the highest amounts of the particular mineral you need to avoid toxicity.

Vitamins are carbon based and are therefore are organic compounds. Anything that is carbon based can be easily destroyed by heat, radiation, and most significantly, time. Minerals, on the other hand, never change in chemical makeup. They never undergo a change to another chemical compound such as we see in the conversion of retinol to retinoic acid. However, they can combine with another compound for a specific function, such as the iron in hemoglobin and myoglobin serves to transport and store oxygen. The only way the mineral makeup of a food can be changed is for it to be depleted from the food. For example, the minerals found in vegetables can be diffused into boiling water, and then be completely lost when the water is dumped down the drain. When you cut vegetables, you increase the rate at which the minerals are lost.

The absorption and transportation processes of minerals by the body are a lot like vitamins in the sense that some minerals, like potassium, are easily absorbed, transported, and excreted by the kidneys, and others are not. Calcium is much like a fat-soluble vitamin in the sense that it needs assistance in the form of carriers and transporters for proper utilization. Since minerals can behave like vitamins in terms of absorption, storage, and excretion, there is the possibility that an excess of certain minerals can lead to toxicity.

The bioavailability of the particular minerals present in food can vary. Bioavailability refers to the amount of the particular nutrient that is available for absorption by the body. Some foods contain compounds called binders that actually hold on to the mineral and prevent it from being absorbed. For example, the compound known as phytic acid (phytates) in legumes, grains, split peas, and parsnips can act in this manner. In foods like spinach, rhubarb, plums, blueberries, nuts, and seeds, oxalates reduce the amount of minerals available for absorption. Although all of these foods provide great health benefits, the bioavailability of the minerals they contain will be low. To lessen this disadvantage, an athlete desiring adequate amounts of minerals would follow the diet-planning principles in Macrobiotic Nutrition, along with the Macrobiotic Food Guide Pyramid (see Appendix B).

Secondary deficiencies can occur due to factors unrelated to deficiency in dietary intake. In the case of vitamins, such factors include other substances

such as drugs and the amount of exposure to the environment. Factors producing secondary deficiencies in minerals include too high an intake of one mineral, which can inhibit the absorption of another mineral. Too much of one mineral can also have a negative impact on the metabolism and excretion of another. For example, high levels of phosphorus will bind with magnesium in the gastrointestinal tract, leading to magnesium excretion.

The Macrominerals

The class of minerals termed macrominerals is present in the body's depositories (bone and teeth), soft tissue, and circulating fluids in amounts larger than 5 grams. Of them, magnesium is present in the smallest amount, which is around 30 grams. The macrominerals are divided into two separate classes, those critical for maintaining the body's fluid balance, and those serving important roles in bone growth and bone maintenance. The minerals that play a role in the body's fluid balance are sodium, chloride, and potassium. The other macrominerals important to the athlete are magnesium, phosphorus, and calcium, all of which play key roles in the processes over the life span of bone growth and maintenance. In the following sections, I establish the amounts of each of these minerals you need to achieve optimum performance and health, and describe the symptoms associated with deficiencies and toxicity. I also identify the foods that provide the highest amounts of the particular mineral to help you avoid any real threat of deficiencies and toxicity.

Fluid Stasis Minerals

Each of these minerals plays a key role in the overall performance of the athlete by maintaining water balance and body pH. Water balance and body pH are chief concerns because, as previously discussed, water provides the proper "environment" for all metabolic reactions to take place.

Sodium. Sodium is the major positive ion (cation) in the fluid outside your cells, and it functions primarily as a volume regulator. In addition to controlling extracellular volume, sodium acts as an electrolyte in the body and assists in the maintenance of the acid/base balance and is essential to nerve transmission and muscle contraction. Foods in general contain adequate amounts of sodium, especially processed foods. Typically 75 percent of a person's sodium intake comes from processed foods. Another 15 percent comes from salt added during cooking and eating. Only the remaining 10 percent of sodium actually is found naturally in the food itself. The intestines directly absorb sodium into the blood, and it is able to travel freely in the blood without the aid of carriers or transporters. Sodium then comes in contact with the kidney, which fil-

ters and releases the exact amount that the body needs. If you eat a food with a high sodium content, your body signals thirst to make you drink more water, increasing fluid levels in the body and allowing the kidneys to process the sodium. Any excess present will be excreted along with the water.

The minimum suggested amount of sodium to be ingested is set at around 500 milligrams and reflects the amount needed in a person who is not actively sweating. The maximum amount of sodium to be ingested is set at 2,400 milligrams. This amount is enough for people who lead active lifestyles and who take part in a wide variety of physical activities in different climates.

Hypertension was considered to be related to the dietary intake of sodium for quite some time. Recent studies have shown that sodium in combination with chloride at high levels raises blood pressure. This compound is sodium chloride, also known as table salt. A point of interest is that sodium alone or chloride alone does not affect blood pressure levels in the way that salt does.

Because sodium is present in adequate amounts in most of the foods we consume, deficiencies only occur for other non-diet-related reasons. These cause conditions in which large amounts of fluid are lost, including diarrhea, vomiting, or heavy sweating during intense exercise. One will experience muscle cramps when sodium deficiencies occur because sodium functions as an electrolyte.

Toxicity resulting from too high a sodium intake can happen but is extremely rare. One of the signs of sodium toxicity is edema, which is excessive fluid buildup in the cells. Hypertension is a condition where blood pressure rises beyond healthy levels. Hypertension can be caused by a number of factors, including exercise, stress, and other existing medical conditions; it can be aggravated by an excessive intake of sodium. Both of these toxic conditions can be treated through adequate water intake.

The best source of sodium is table salt, but you should avoid consuming salt in excess. Instead, you should select foods like meat, milk, bread, and vegetables, which all provide moderate amounts of sodium. The largest amounts of sodium can be found in processed foods, but you would be wise to limit the amount of processed foods that you consume. Processed foods typically have little or no nutritional value, are often high in calories, and usually don't meet Macrobiotic Nutrition's recommended 45/35/20 lean-mass equation.

Chloride. Chloride is the negatively charged anion (ion with a negative charge) used by the body to control the levels of extracellular fluids. Chloride has free access to the cell membrane, and is able to move freely in and out of the cell. Outside of the cell, it is in close association with sodium, while inside the cell it can interact with potassium, classifying chloride as an electrolyte. In

addition to maintaining normal electrolyte levels in the body, chloride is a part of hydrochloric acid (HCL), which maintains the acidic levels of gastric juices. The gastric juices play an indispensable role in digestion, so any decreased production of HCL leads to weakened levels of gastric juices. A case of constant vomiting would lead to a decrease in gastric juices.

Like sodium, chloride is found in abundant amounts in foods, especially processed foods, where it is combined with sodium and other salts. Chloride deficiencies are caused by excessive fluid losses from sweating, diarrhea, or vomiting. Dehydration can cause unusually high toxic concentrations of chloride in the body and is accompanied by vomiting. Adequate food consumption and water intake will rectify the situation.

The best source of chloride is table salt, where chloride is bonded to sodium in the form of sodium chloride. Other foods providing moderate amounts of chloride are meats, milks, and eggs. Again, like sodium, the largest amounts of chloride are found in processed foods.

Potassium. Potassium is another cation used by the body to control fluid levels. Unlike sodium, the other cation, potassium is found inside the cell and serves a host of key functions including maintenance of electrolyte balance and cell integrity. Another major function of particular concern to the athlete is for nerve transmission and muscle contraction. When a nerve impulse is fired to contract a muscle tissue, potassium and sodium briefly trade places across the membrane of the cell. This is of critical interest to athletes, because proper levels of potassium and other electrolytes directly affect performance and help maintain a steady heartbeat.

Potassium deficiencies are rarely, if ever, caused by a low dietary intake. The most common route for potassium to be low is through excessive fluid loss from dehydration, vomiting, and diarrhea. Food-related toxicity is not a concern. Since potassium is found in all living cells, any food that is fresh and unprocessed will contain significant amounts of potassium, and relying on a balanced diet will provide adequate amounts of it. Sources include the potato, banana, acorn squash, sirloin steak, and chicken.

The Bone Builders

As highlighted throughout this chapter, certain vitamins and minerals can be grouped together because they are in close association with each other and because they are synergistic—in other words, one increases the effectiveness of the other(s) and vice versa. Calcium and phosphorus together play a primary role in bone health. In addition to bone health, they are individually responsible for many functions in the body.

Calcium. Calcium plays the most important role in the class of bone builders. It is the most abundant mineral found in the body; it amounts to 39 percent of the total body minerals and total calcium measures about 1.5 to 2 percent of the body weight. The majority of the calcium present in the body is found in the teeth and the bones. Calcium plays two critical roles; the most obvious is its role as the key mineral in bone structure. The second role is as a depository for calcium stores. If a drop in blood calcium levels should occur for any reason, calcium can be drawn from the bones and teeth to help raise the level of blood calcium back to its optimum levels.

Calcium appears in various roles when it is in combination with other minerals. For instance, when bones are forming, crystals of calcium and phosphorus called hydroxyapatite deposit on the matrix of collagen. When the bone is going through the process of mineralization, hydroxyapatite and other minerals become denser in their crystal structures, leading to a stronger bone. This process is constantly happening because bone is continually being torn down and remodeled.

While 99 percent of the calcium in the body is held in the body's depositories, the remaining 1 percent is circulating in body fluids. In this form, calcium is ionized and is absolutely critical to life functions. The ionized fluid calcium aids in the regulation of muscle contractions, plays a role along with vitamin K in the clotting of blood, and assists in the transmission of nerve impulses, the activation of certain enzymes, and the secretion of hormones. For example, ionized fluid calcium activates the protein calmodulin, which is responsible for relaying messages from the surface of the cell to the interior portion of the cell. Some of these messages regulate blood pressure. This has led to the suggestion that higher levels of calcium will not only help reduce the severity of hypertension, but will also provide the depositories with enough calcium to prevent osteoporosis. Because of the many critical functions it performs, maintaining proper levels of calcium in the blood is one of the top priorities that the body addresses, and several hormones and vitamin D control the blood level of calcium. Since the depositories can provide adequate amounts of blood calcium despite a low dietary intake of calcium, an important issue related to a deficient intake of calcium is reduced bone integrity. This process is long and slow, so it can take years before an incident demonstrates a lack of bone integrity.

There are instances where abnormally elevated levels of calcium in the blood can result in muscular contractions that can't be relaxed, a condition called calcium rigor. Calcium tetany, which manifests the same symptoms as calcium rigor, will occur when blood calcium levels are well below normal. In either case, the altered levels of blood calcium is caused by either a decreased

production in the hormones parathormone and calcitonin, which increase or decrease the secretion of calcium from the depositories, or a decreased level of vitamin D. These two muscular contraction abnormalities do not result from dietary excesses or inadequacies of calcium.

Like the fat-soluble vitamins, the mineral calcium needs a transporter to help out with absorption. The acidity of your stomach keeps calcium in a soluble state, and vitamin D plays a role in the formation of a specialized calcium transport protein. This specialized protein is called calcium-binding protein (CBP). Whenever the body is in need of added calcium from the diet, it produces more CBP to facilitate calcium absorption.

Under some conditions, insufficient production of vitamin D inhibits the synthesis of CBP, which is why much of the available milk on the shelves of the supermarket is fortified with vitamin D. Other substances that can negatively affect the absorption of calcium are the phytates and oxalates commonly associated with the fiber in whole grains and vegetables. For this reason, whole grains and vegetables are not suitable sources for dietary calcium.

This should not be a concern for the athlete who builds his or her diet using the principles based on Macrobiotic Nutrition and the Macrobiotic Food Guide Pyramid (see Appendix B), which emphasizes the use of all the food groups. The best source of calcium is found in yogurt and milk. Many athletes, particularly bodybuilders, restrict dairy in their diet. This is a big mistake. True, most dairy products contain high levels of saturated fat. However, dairy sources such as skim milk, low-fat cottage cheese, and non-fat yogurt have great macronutrient profiles and are excellent low-fat protein sources. And many people don't realize that these dairy sources are relatively low glycemic (skim milk has a glycemic index of only 32) even though they contain some lactose. An athlete who is a lactovegetarian will severely limit the amount of calcium present in his or her diet. This does not take into account the fact that whole grains and vegetables, the staples of the typical vegetarian diet, bind up calcium and increase the likelihood of calcium deficiency. Calcium deficiency can also be a concern to athletes who are either lactose intolerant or allergic to milk and its associated products. This would include allergic reactions to the proteins casein and whey (both of which are cornerstones to the athlete) and an inability to break down the milk sugar lactose. In these cases, protein needs can be met elsewhere, through the use of meats and other adequate sources of protein like beans and rice, but calcium will still be missing from the diet. This can be easily remedied by the use of calcium-fortified orange juice and/or milk-based products containing added lactase to break down the lactose into the two base sugars galactose and glucose.

As I have established, any prolonged period of inadequate calcium intake

will lead to decreased bone mineral mass and density. The deficiency manifested as osteoporosis, a condition that can be easily prevented, is very prevalent in the older population. The RDA for both men and women ages nineteen to thirty is set at 1,000 milligrams of calcium a day. In addition to the measure for the RDA, a tolerable upper intake level (UL) has been established, which represents the highest supplemental amount of a particular nutrient, where toxicity is avoided and most people remain healthy. The UL for calcium is set at 2,500 milligrams for both men and women ages nineteen to seventy years. If calcium intakes exceed this level, the potential for toxic symptoms like constipation, kidney dysfunctions, and urinary stones will develop. In addition to those conditions, high levels of calcium will decrease the bioavailability of other key minerals.

Phosphorus. Phosphorus is the second most abundant mineral based on the amount present in the body. It operates in association with calcium in the form of hydroxyapatite crystals in the bone mineral matrix and in the teeth. Actually about 85 percent of the body's phosphorus mass is stored in the depositories. The remaining 15 percent is present in a host of compounds found in the body.

Adenosine triphosphate (ATP), thiamine pyrophosphate, pyridoxal-5-phosphate, and phosphoric acid (the body's buffer system) and its various salts, phospholipids like the chylomicrons, and the membranes of cells all contain a phosphate group in their chemical structure. These compounds would be rendered useless without phosphorus, so we can see that this mineral plays a significant role in the performance of an athlete because of its major role in the release of energy at all stages of metabolism.

Dietary deficiencies of phosphorus are unknown. The reason for this is because phosphorus is present in adequate amounts in any diet with an adequate amount of animal tissue, and legumes and milk are also good sources of phosphorus. In addition, phosphorus is present in processed meats and other processed foods, and in soft drinks. Deficiencies can occur because of a drug interaction with phosphorus. Some drugs bind to the phosphorus, making it unavailable for any of the vital functions associated with metabolism. Phosphorus deficiency symptoms are manifested as muscle weakness, a lack of energy, and bone pain.

Magnesium, the Mighty Mineral

Although magnesium is in amounts so low that it barely qualifies to be considered a macromineral, it serves a host of indispensable functions to the athlete. Over half of the body's magnesium is stored in the bone depository, and

most of the remaining amounts are held in the soft tissue of the body. About 1 percent remains in the extracellular fluid of the body. The magnesium in the soft tissue performs a vast range of roles in the body. It is a main component in the synthesis of protein, and a major player in many of the body's enzyme reactions. It functions as a catalyst in the final role of ATP production, where the final phosphate group is added to adenosine diphosphate (ADP). Magnesium is vital for countering the effects of calcium in muscle contraction and blood clotting.

In most people's dietary intake, magnesium is one of the minerals that falls far below what is considered adequate. This potential deficiency is of particular concern to the athlete, because of magnesium's role in protein synthesis and energy production. A diet high in legumes will provide sufficient amounts of magnesium to compensate for this lack. Other significant sources of magnesium are halibut, cashews, artichokes, and millet.

Toxicity is rare when magnesium is derived from the diet. However, when taken improperly in supplemental form, excessive amounts of magnesium can lead to diarrhea, which leads to dehydration.

Sulfur, the Protein Architect

Sulfur is last in the line of the macrominerals and is not used by the body in its original form. Your body derives this mineral from certain B vitamins and amino acids. In the formation of proteins, sulfur plays an important role in protein folding. The way a protein folds determines how it functions. Insulin is the hormone that promotes glucose utilization, protein synthesis, and the formation and storage of neutral lipids. This crucial hormone is a protein created by the use of three disulfide bridges that exist between the six L-cysteine amino acids.

Cysteine is a nonessential amino acid, which means the body can manufacture it in the presence of sulfur. Therefore, there are no deficiencies associated with sulfur as long as protein intake is adequate. Adequate protein intake can be obtained by an athlete who follows the principles of Macrobiotic Nutrition. When an athlete follows Macrobiotic Nutrition, all of these essential macrominerals are provided in adequate amounts to ensure maximum human performance.

The Microminerals

The microminerals are as important to the body's health and functioning as the macrominerals. They are considered microminerals because of the relatively small amounts that are present in the body. These microminerals, also commonly known as trace minerals, are supplied in adequate amounts in people's diets, to both maintain health and avoid toxicity. In the discussion of the trace

minerals, I determine why these elements are so important to the athlete's performance. To optimize the total bioavailability, I also determine which foods contain the greatest concentration of the particular trace mineral.

The category of trace minerals includes the minerals iron, zinc, selenium, chromium, copper, manganese, fluoride, molybdenum, and others. Most of the minerals mentioned are available in supplemental form, but you should avoid excessive use of these products, because most of the time these supplements contain amounts of the trace minerals that are in excess of what is needed to maintain health. Any amount over what is suggested will produce conditions that are undesirable, which could include organ damage, muscle pain, exhaustion, and fatigue. So, it is vital to address the toxic signs of the trace minerals and the amounts at which toxicity can be avoided.

Secondary deficiencies are another issue associated with an overabundance of trace minerals. Remember, secondary deficiencies are not caused by inadequacies of nutrients in the diet; they are caused by other nondietary factors. In the case of the trace minerals, a nondietary factor may be caused by an excess of one trace mineral that is inhibiting the absorption of another equally important trace mineral.

Iron, the Oxygen Transporter

Iron is a mineral that is needed in amounts ranging from 8 milligrams per day for men to about 18 milligrams per day for women. The importance of its role to the athlete is unmatched, despite the fact that this mineral is present in the body in relatively small amounts,

Iron exists in the body in two distinct states of charge, called ionic states. Ferrous iron has a +2 charge, while the ferric iron form has a +3 charge. Since iron has the ability to exist in two distinct states, it can participate in a wide range of enzymatic reactions, and it also plays a principal role in the production of ATP. The enzyme fumarase reduces NAD^+ to NADH in the TCA cycle of metabolism. NADH can then provide H^+ to the electron transport chain. In the electron transport chain, cytochromes, which contain iron, facilitate the movement of H^+ from the matrix to the intermembrane space of the mitochondria. Fumarase and cytochromes are thus key components involved in the production of ATP.

The most common use of iron by the body is in the compounds that transport oxygen to tissues. Hemoglobin and myoglobin are two proteins that have a heme group, which contains ferrous iron. Hemoglobin is the primary transport for oxygen, while myoglobin is used mostly for storing oxygen in the muscle.

When a muscle tissue is active, a buildup of H^+ and CO_2 are produced. Hemoglobin has a high affinity for H^+ when little oxygen is present at the site of the active tissue (conversely, when blood passes through alveoli of the lungs, oxygen concentration is higher than H^+ concentrations, so oxygen binds to hemoglobin). Due to this fact, oxygen is traded for H^+ in a muscle tissue that is actively respiring. Without iron present in hemoglobin, there is a decreased ability to transport oxygen. This has a large impact on athletic performance, and is discussed later in this section.

Since iron is used in the part of hemoglobin that actually accepts oxygen, it is understandable why the body would be designed for the conservation of iron. The body doesn't excrete iron too well because of this; in fact, one of the only ways for your body to lose iron is through bleeding. That is why premenopausal women need 10 milligrams more iron than do men. The body maintains the levels of iron primarily through the manipulation of absorption, absorbing more when it is needed and less when it is not.

Iron absorption is a process that requires three specialized proteins: mucosal ferritin, mucosal transferrin, and blood transferrin. When iron is present in the small intestines, a protein formed in the mucous lining of the intestinal cells binds to the iron. This protein is called mucosal ferritin. When the body signals for the absorption of iron, the mucosal ferritin gives up the iron to another mucus-based protein, mucosal transferrin. Blood transferrin then transports the iron to the rest of the body, where either bone marrow or other cells will use it. The cells of the intestines renew themselves every three days. When they are excreted out of the body through feces, any iron that is not needed by the body at the time will be lost. Having the ability to store some iron is vital to the body. A reasonable amount of the excess dietary iron is stored in the liver or the spleen. The liver creates specialized proteins called ferritin and hemosiderin. The body readily breaks down ferritin, so a constant blood supply of iron is always available. Hemosiderin is formed from ferritin when blood iron levels become abnormally high.

To consider which type of iron has the highest bioavailability, let's quickly examine the storage of excess carbohydrates as fats, and excess fats as fats. Which is easier (in terms of caloric expenditure) for the body to do: to take a carbohydrate and chemically convert it into a fatty acid and then chemically combine it to a glycerol backbone, or to take a fat and store it as fat? The answer is obvious. So too is the bioavailability of heme iron, which is used in hemo/myoglobin. For example, heme accounts for about 10 percent of the total iron acquired from the diet, yet 25 to 35 percent of the heme iron is actually absorbed. This is significantly more than the 10 percent that is absorbed from nonheme iron.

There are several factors to consider that contribute to the overall bioavailability of iron in the diet. Since any flesh-based source of iron is best as far as bioavailability is concerned (heme iron), it is not surprising that these sources also contain a substance that enhances the all-round absorption of iron. It is called the MFP factor, because it is associated with the digestion of meats, fish, and poultry. Vitamin C also enhances the absorption of nonheme iron. This is because ascorbic acid is able to hold on to the reduced form of iron, which is ready for absorption. These two absorption enhancers are met with a degree of opposition from phytates that naturally occur in some foods, but this can be easily compensated for by the use of a wide variety of foods, as demonstrated by the Macrobiotic Food Guide Pyramid. (See Appendix B.)

When the body stores of iron fall to extremely low levels, iron deficiency anemia becomes the major concern for the athlete. In this condition, the red blood cells are small and pale, and because of their physical state, their ability to bind to oxygen in the lungs is severely diminished. Energy production is decreased since the body needs oxygen to perform metabolism at its optimal level, so symptoms of this condition manifest in the form of fatigue, weakness, headaches, and poor core-temperature regulation. Iron deficiency anemia is visibly detected by a pale complexion on a person with fair skin. It can be detected in people with a darker skin tone by checking the lining of the eye, which will become pale rather than pink. Because the body conserves iron, the main cause of iron deficiency would be excessive blood loss.

The potential for greater harm exists with the toxicities associated with iron. A condition known as hemosiderosis will develop when iron is taken in excess. Repeated blood transfusions, which completely circumvent the body's intestinal defense, and naive iron supplementation, which overwhelms the intestinal defense, cause the liver to transform ferritin into hemosiderin. Hemosiderosis is a condition in which buildup of hemosiderin in the liver and other tissues can potentially cause tissue damage. Infections are more common, not because there is a decrease in immune function, but because bacteria thrive on iron-rich blood. Athletes who rely on Macrobiotic Nutrition and its associated food guide pyramid will have no associated iron deficiencies or toxic symptoms, thus ensuring optimal performance.

The best dietary sources of iron are the flesh of animals since it has iron in the same form that is used by our bodies. Ground beef, sirloin steak, and shrimp are all superior choices for iron. Kidney, garbanzo, pinto, and navy beans are also excellent legume sources of iron. Other enriched foods also provide adequate amounts of iron, including oatmeal, spaghetti, and flour tortillas (all of which are low glycemic).

Zinc, the Cofactor

Zinc is a major component of more than 100 different enzymes called metalloenzymes that carry this label because a mineral is included in their structural makeup. Zinc has the ability to stabilize the membrane of the cell, which makes it crucial in the defense against free radicals. It is a participant in the synthesis, storage, and release of the blood sugar regulating the hormone insulin. Zinc is another component of the body's blood-clotting system, and it affects the functions of thyroid hormone. It is also essential for vision, since it produces retinal, the form of vitamin A required for the visual pigments.

Zinc absorption is similar to that of iron in certain ways. When the body has enough zinc to maintain optimal performance and health, it reduces the amount of absorption that occurs. As with iron, fiber and its associated phytates bind to the excess zinc, limiting the bioavailability of the mineral.

When zinc is absorbed into the microvilli (fingerlike projections on intestinal cells), the variety of roles it can take on is great. It can be absorbed into the cell right away, where it can function as a component of either DNA or RNA polymerase. This is significant because the intestinal cells renew themselves frequently and zinc is therefore absolutely crucial to the growth of new intestinal cells. Another possible avenue for zinc is in the storage protein called metallothionein. This specialized binding protein helps regulate zinc absorption, much like iron's mucosal ferritin. Metallothionein is stored in the liver, so the body has some reserves if there is any need for more zinc.

The pancreas produces many of the enzymes with zinc in their structure that are critical for digestion. When you consume a food like steak, the dietary zinc is combined with the zinc from the pancreatic juices. The zinc derived from the pancreas can be reabsorbed into the intestinal cells for growth or return to the pancreas for further use as metalloenzymes.

To illustrate the importance of a proper diet and ill-advised supplementation, let's briefly consider the interaction between iron and zinc. Some plasma zinc binds to transferrin, which is also the preferred mode of travel for diet-derived iron in the blood. Adults who overuse iron supplements will overly saturate the transferrin-binding sites and leave them unable to absorb zinc. The converse is true too; large amounts of zinc will inhibit the absorption of iron.

Large dosages of zinc interfere with copper absorption too. This interference may occur because the two minerals compete for absorption with one another. When an abundance of zinc is ingested, the intestinal-absorbing cells produce more metallothionein, a binding protein that stores zinc in much the same way that iron is stored in mucosal ferritin. Metallothionein also binds to other microminerals, such as copper and cadmium. A higher affinity exists between copper and metallothionein than between zinc and metallothionein.

So increased levels of metallothionein, due to an abundance of ingested zinc, will bind to copper as well, and severely limit the copper that is freely available for absorption. Thus copper availability is reduced because of the larger amounts of zinc. This point illustrates the most common mistake people make with supplements. It can be avoided by relying on a sound diet and some key supplementation.

Since zinc absorption is similar in principle to iron absorption, it follows that its loss from the body occurs in the same basic fashion. Since metallothionein is produced by the microvilli, any tied-up zinc is excreted in the feces when the microvilli are replaced.

When zinc levels are all but nonexistent, growth in general will be retarded because zinc plays such a large role in the area of cellular growth and protein synthesis. Deficiencies of zinc also provoke diarrhea, which worsens not only the zinc deficiency, but also the presence of other essential nutrients. Other symptoms of deficiency include an altered thyroid function and metabolic rate, warped taste, and a lowered rate of wound healing. These zinc-deficiency related conditions can manifest even when only mild zinc deficiencies exist in the diet.

The toxic symptoms of excess zinc may include vomiting, exhaustion, muscle pain, dizziness, and drowsiness. Too much zinc interferes with the absorption of copper and its proper metabolism. When researched in test animals, excess zinc has led to degeneration of cardiac muscle. High zinc levels also reduce the production of red blood cells, which further affect performance.

It is wise to rely on a variety of healthy foods to supply the body with the right amount of zinc and avoid the pitfalls of incorrect supplementation. Zinc is found in the highest concentrations in protein-rich foods, such as sirloin steak, ground beef, ham, dark turkey meat, and yogurt.

Iodine, the Metabolic Trace Mineral

Iodine is the food-based form of the ionized form iodide used by our bodies. Although iodide is found in very small amounts in the body, its role is critical. It is a primary component of the thyroid hormones triiodothyronine (T_3) and tetraiodothyronine (T_4 ; thyroxin). These hormones are used to regulate cellular metabolism and other functions like body temperature, reproduction, growth, and muscular function. These critical hormones actually regulate the amount of oxygen a cell uses and therefore manipulate the amount of energy released during basal metabolism.

Iodine deficiency is not common in developed countries, so we don't have to concern ourselves with this issue in great detail. To provide a basic overview, the hypothalamus controls the production of thyroid-stimulating hormone

(TSH), which is produced by the pituitary gland. During iodine deficiencies, thyroid-hormone synthesis declines. TSH is produced in larger amounts in an attempt to increase absorption of whatever iodine is left in the thyroid gland for thyroxin synthesis. If this condition persists for an extended period of time, the cells of the thyroid gland enlarge to the extent of developing into a large lump on the neck. The symptom of a toxic iodine level is identical to the symptom of deficiency, and it is also identified by a rather large lump on the front of the neck.

To avoid any form of deficiency or toxicity, the athlete would be wise to plan a well-balanced diet with a firm basis in Macrobiotic Nutrition. This diet will ensure an adequate amount of iodine from seafood and other foods cultivated in iodine-rich soil. If you are allergic to certain proteins found some seafoods, you can conservatively use iodized table salt, which contains the RDA in half a teaspoon.

Selenium, the Antioxidant

Selenium is a cofactor in the enzyme glutathione peroxidase. It facilitates the reduction of toxic hydrogen peroxide within the cells. This enzyme works in conjunction with vitamin E in the defense against free radicals, so selenium can prevent oxidative damages to the cell and block free-radical formation. When selenium can't block the formation of free radicals, vitamin E will be able to stop it. The enzyme responsible for converting tetraiodothyronine to triiodothyronine, which is the active form of the hormone, contains selenium.

A deficiency in selenium is associated with a type of heart disease that is prevalent in an area of China; this deficiency is also reported in New Zealand and Finland, areas of the world where the soil is not very rich in selenium. The cardiac muscle is enlarged in this disease of the heart, and the normal tissue that composes the middle layer of cardiac walls is replaced by fibrous tissue. Since we live in a developed part of the world, selenium deficiency is hardly a concern for us. The selenium content of the soil in North America has a significant amount of this essential mineral, so a well-balanced diet will provide enough selenium. Symptoms of selenium toxicity may include vomiting, loss of hair and nails, lesions of the skin, and diarrhea, when people ingest a milligram or higher per day.

Copper, the Oxide Eater

Copper plays a diverse role in enzymatic activities. All of the metalloenzymes containing copper in their structure have similar functional characteristics, and consume oxygen and its associated radicals. The most well-known enzyme that protects against free radicals is a compound called superoxide dismutase

(SOD). This compound facilitates the conversion of the potentially dangerous compound superoxide anion into hydrogen peroxide and oxygen ($2\text{O}_2^- + 2\text{H} \longrightarrow \text{H}_2\text{O}_2 + \text{O}_2$). Another free-radical enzyme that relies on copper for its properties is a dehydrogenase protein called ceruloplasmin. This protein is involved in the transport and storage of copper, but it also can reduce superoxide anions without any intermediate phase, making it extremely efficient.

Another enzyme that must contain copper in order to function is an enzyme called lysyl oxidase. This protein is necessary for an essential step in the cross-linking of collagen strands and for the healing of wounds. Copper works in a similar fashion to iron in the process of energy production. Cytochrome C oxidase is associated with the pumping of H^+ protons and the resulting phosphorylation (the addition of a phosphate to an organic compound) of ADP to ATP. Even though copper is present in the body in small amounts, usually around 100 milligrams, it is evident that it performs in a wide range of vital life functions.

The issue of copper deficiency is of no real concern to people living in the United States and Canada, since the typical diet includes adequate amounts of this trace mineral. Extremely high levels of vitamin C can interfere with copper absorption, and if prolonged, may lead to deficiency. Deficiency can result in anemia due to the reduced hemoglobin synthesis. Copper toxicity is also a rare occurrence, and should not be a concern for any athlete who follows Macrobiotic Nutrition and its associated food guide pyramid (see Appendix B) for guidelines on proper dietary practices. If copper supplementation is abused, toxicity is certainly a concern. It will bring on symptoms like vomiting and diarrhea, and if prolonged will result in liver damage.

The athlete who desires adequate amounts of copper in the diet should make sure she or he eats enough legumes, seeds, nuts, and organ meats, such as liver. Another source of copper is tap water; its value as a source depends on the hardness of the water and whether or not you have copper pipes.

Manganese, the Magnesium Substitute

This trace mineral is found in the bones and in metabolically active tissue like the liver, pancreas, and kidneys. If you were to combine all of the manganese in the body, it would measure about 20 milligrams. However, this mineral plays a role in the formation of energy with the enzyme pyruvate carboxylase in the conversion of pyruvate to oxaloacetate (an intermediate in the TCA cycle), and is a cofactor in the copper enzyme SOD. It can also substitute for magnesium in many of its associated enzymes when magnesium is needed to activate the enzyme.

If an athlete relies on proper dietary principles, a primary manganese deficiency is a rare concern, partly because of the low RDA and its relative abundance in plant-based foods. The same can't be said of secondary deficiencies, however. Overuse of iron and calcium supplements can inhibit the absorption of manganese as well as the phytates found in legumes, grains, and seeds, which, ironically, are the best sources of manganese. A deficiency may become apparent in poor growth and nervous-system disorders. Toxicity should not be a concern for athletes who rely on sound dietary practices for their RDA for manganese. If an athlete oversupplements with manganese, the toxic effects will present themselves as nervous-system disorders.

Fluoride, the Bone Hardener

This abundant trace mineral is found in soil all over the world and is also present in water, plants, and animals. Yet fluoride is found in very small amounts in the human body. However, fluoride is crucial for the formation of a harder bone-mineral matrix. The macrominerals calcium and phosphorus combine to form a crystal compound called hydroxyapatite. These crystals harden with the addition of other minerals on the protein collagen. Fluoride then replaces the OH groups of the hydroxyapatite crystals, forming fluorapatite and making it stable. This compound makes bones harder and teeth more resistant to decay. Too much fluoride will have an aesthetically damaging effect. A condition known as fluorosis causes unsightly white specks on the teeth. In the more severe cases, the enamel will become permanently stained. Toxicity can occur only in areas where fluoridated water exceeds 150 ppm (parts per million). In these cases symptoms include vomiting, associated nausea, diarrhea, pain localized in the abdomen, and tingling in the extremities. The best dietary sources for fluoride are seafoods and most regular teas.

Chromium, the Insulin Augmentation Mineral

Chromium is essential in the metabolism of carbohydrates and lipids. It is similar to iron in that it exists in several different ionic states, but the state that seems to have the most benefit to metabolism is the 3+ form of chromium. Chromium is a component of compounds called glucose-tolerance factors. It helps maintain a balance in blood sugar by improving the hormone insulin's activity in the body. Because of the improved actions of insulin, less of the hormone is needed to maintain proper blood glucose levels. If chromium is deficient diabeteslike symptoms will result, manifesting in impaired insulin response, glucose tolerance, and glucagon response. Glucagon is the counter hormone to insulin that releases stored glucose (glycogen) into the blood-

stream when blood glucose levels are low. A decreased level of stamina will be noted in the athlete who is low on chromium. Toxic symptoms are unknown. The best dietary sources of this trace mineral are liver, whole grains, cheeses, and nuts.

Molybdenum, the Oxidase Mineral

Molybdenum is present in such small amounts that dietary deficiencies and toxicities have not been reported in humans. It is a component of several metalloenzymes, such as xanthine oxidase, aldehyde oxidase, and sulfite oxidase. The best dietary sources of molybdenum are legumes, green leafy vegetables, skim milk, and liver. Toxicity has been observed in test animals, where kidney damage and reproductive problems became apparent.

Notable Mentions

Other trace minerals that have been shown through research on animals to have beneficial qualities are nickel, vanadium, cobalt, silicon, and boron. However, the exact amounts of these trace minerals in the body are extremely difficult to determine, and deficiencies and toxicities are not known. Their functions are known, however. Nickel may be able to serve as a cofactor for some enzymes. Vanadium is necessary for proper bone and growth development. Cobalt is the mineral present in vitamin B₁₂ (cobalamin), and silicon is used in the synthesis of collagen and its subsequent bone formation. Boron may be used to facilitate optimum brain function.

Macrobiotic Nutrition is a concept that allows for optimum health. All of the essential nutrients mentioned within the preceding chapters are found within foods that are central to the Macrobiotic Nutrition philosophy. Food is the ultimate transporter for all of the nutrients that are key to overall fitness. Following the principles of Macrobiotic Nutrition will allow the average athlete to grow into a class all his or her own and attain maximum human performance.

CHAPTER 9

Enhancing the Macrobiolic Nutrition Effect

THE PREVIOUS CHAPTERS HAVE SHOWN YOU how the Macrobiolic Nutrition approach provides the core building blocks for you to experience tremendous increases in muscle building, strength, and performance, while promoting leanness. You will be extremely satisfied with the results you will get from following the core Macrobiolic Nutrition program alone, but this program can be intensified by using the special Up Your MASS muscle-building nutrition products. This combination of program and nutritional product will take your body into a hyperanabolic state that you have never experienced before, and that until now has only rarely been encountered by top bodybuilders. Once your body enters into this Macrobiolic-induced lean muscle-building mode, your body will undergo that Herculean transformation you are looking for, culminating in a massive, superior sculpted physique.

However, if you are like me and other champion athletes, you want to know if you are doing everything possible to pump out that extra rep, to get the results you want—*faster*. If you have the same competitive spirit that I have, it is natural to be wondering, *Is there anything else I can do to enhance the Macrobiolic Nutrition effect?* For example, you may want to get bigger faster, stronger faster, and/or leaner faster. I know that is what I wanted: the fastest and best results possible. I spent years of research and development to determine if any additional ingredients and delivery-system technologies existed to enhance the Macrobiolic Nutrition effect. I determined that the answer is yes, there are extra measures you can take to further enhance Macrobiolic Nutrition, and now I will share the results of this massive research effort with you—my secrets of how to enhance the Macrobiolic Nutrition lean muscle-building effect.

This chapter is written to give you more knowledge and some precise direction on other products to consider including as part of your Macrobiolic Nutrition program. You will be interested to learn that sports nutrition prod-

uct development and production technology has made significant advancements in the past several years. This means you can now get even better results with well-known, clinically proven sports nutrition ingredients. For example, we all know creatine works to increase muscle size and strength, but did you know that there are sports supplement breakthroughs that may make creatine work even better? The same is true for other research-tested sports nutrients like glutamine, and fat-metabolizing dietary supplement formulas. You will also be interested to discover that a new branch of sports nutrition, muscle-building and performance science has developed as a result of the medical insights gained from working with joint and connective-tissue health issues. It deals with certain dietary and supplement ingredients, and how we can enhance joint structure and function and help to reduce exercise-induced pain and inflammation through nutrition.

You will get an inside look into how MHP develops its top-selling sports nutrition products. You will also realize that one major indication that MHP products are top quality and the most effective available is that I use them, the MHP staff uses them, and my friends use them—so I am going to make very sure that only the best ingredients are used in precise combinations! Because of my competitive drive for excellence, I am constantly trying to improve on MHP's sports nutrition products to make sure we receive the best possible results. So get ready to experience more ways to get bigger, stronger, and leaner faster, and develop a physique of the champions.

GETTING BIGGER AND STRONGER FASTER

In the world of bodybuilding, you can never be “too big.” Bigger is always better. Or at least that is the way we view it. The same applies for many other sports. Most athletes always feel they can be bigger and stronger. While we all continually strive to get bigger and stronger, we want to get these results as quickly as possible, so let's look at some supplements to help us get bigger and stronger faster.

Creatine, Creatine Cofactors, and Delivery Systems

Creatine has become one of the most popular sports supplements on the market today, due to the amount of substantiating research published beginning about 1993 that started this explosion of use. The first generation of creatine supplements was based on mimicking these scientific studies. Typically, researchers in such studies use megadoses of the substance they are researching to make sure that if the substance produces any beneficial results, they will be measurable.

The number of scientific research studies documenting the effectiveness of

creatine in muscle development continues to grow. This tremendous body of knowledge has enabled us to determine many facets of creatine's effective activity in the body, and I used these insights and sports nutrition technology to determine how to create a creatine-based product that produces maximum results. These technological breakthroughs in sports supplements included getting better results with less creatine, and maximizing these results with creatine cofactors.

Here's a short overview of creatine supplementation and new research findings: To date, the major scientifically proven benefits of taking creatine supplements include those listed in Table 9.1.

TABLE 9.1 CREATINE SUPPLEMENTATION RESULTS

Creatine Supplementation Has This Effect on the Body	Which Results in Muscle Performance
Increases muscle content of creatine	Increased peak muscle power output
Increases muscle content of phosphocreatine	Increased one repetition maximum
Increases lean body mass	Increased vertical jump
Increases muscle size	Increased strength and power Speeds short duration sprinting

The newest creatine research efforts are focused on fine-tuning our current understanding of creatine and exploring the magnitude of its use. For example, E. S. Rawson and J. S. Volek, at the Department of Exercise Science and Athletics, Bloomsburg University, Bloomsburg, Pennsylvania, conducted a review of twenty-two published research studies and reported their results in the November 2003 issue of the *Journal of Strength and Conditioning Research*. Based on this review, the overwhelming scientific evidence determined that creatine supplementation improves muscle strength and weightlifting performance when combined with resistance training. Of the twenty-two studies reviewed, the average increase in muscle strength following creatine supplementation plus resistance training, at a one-, three-, or ten-repetition maximum, was 8 percent greater than the average increase in muscle strength following placebo ingestion during resistance training. Similarly, the average increase in weightlifting performance following creatine supplementation plus resistance training was 14 percent greater than the average increase in weightlifting performance. The increase in bench press, at one-repetition maximum, ranged from 3 to 45 percent, and the improvement in weightlifting performance in the bench press ranged from 16 to 43 percent. Rawson and Volek concluded that there is substantial evidence that creatine supplementation during

resistance training is more effective at increasing muscle strength and weight-lifting performance than is resistance training alone, but added that the response is highly variable.

Another research review conducted by R. B. Kreider of the Exercise and Sport Nutrition Laboratory, Center for Exercise, Nutrition and Preventive Health Research, Department of Health, Human Performance and Recreation, Baylor University, provides further evidence of creatine's effectiveness. Kreider determined that a review of the scientific literature indicates that more than 500 research studies have evaluated the effects of creatine supplementation on muscle physiology and/or exercise capacity in healthy, trained, and various diseased populations. Short-term creatine supplementation has typically been reported to increase total creatine content by 10 to 30 percent and phospho-creatine stores by 10 to 40 percent. About 70 percent of the approximately 300 studies that have evaluated the potential ergogenic (performance-increasing) value of creatine supplementation report statistically significant results, while the remaining studies generally report nonsignificant gains in performance. For example, creatine supplementation has been shown to improve short-term physical performance activities such as peak power and strength muscle output and all-out-effort sprint performance, and to increase the number of sets and reps performed during resistance-type training. Furthermore, creatine supplementation combined with training has been reported to promote significantly greater gains in strength, fat-free mass, and performance, primarily in high-intensity exercise tasks. Kreider goes on to add that the preponderance of scientific evidence indicates that creatine supplementation appears to be a generally effective nutritional ergogenic aid for a variety of exercise tasks in a number of athletic and clinical populations.

The research results support the use of creatine supplementation for improving muscle strength, size, and power—that is, as a nutrition enhancer for strength athletes and bodybuilders. Creatine increases energy during maximum muscle contraction. During the very first millisecond of a grueling all-out muscle contraction, the ATP is quickly depleted in type 2 muscle fibers. Creatine phosphate (CP) is standing ready in the muscle cells to regenerate new ATP. This process happens in just a few seconds. Once CP is depleted it takes glycolytic energy systems to regenerate ATP and CP; and muscle-energy output decreases, lactic acid builds up, and fatigue sets in, so having a plentiful supply of creatine during those very strenuous short-rep workouts can ensure a maximum supply of CP at the cellular level. This is one main reason why muscle fibers increase in size to become stronger. When fibers increase in size, they increase their storage capacity to hold more ATP and CP. This gives the muscles more energy to increase starting strength and capacity to maintain

contractions for longer periods of time. This is, of course, an oversimplification of the hypertrophy process, but it serves to underscore creatine's role in muscle strength and muscle-fiber content.

The form of creatine that research studies have determined works best is creatine monohydrate, and this is MHP's ingredient of choice, with one improvement. The early research studies used just pure creatine monohydrate, but eventually researchers wanted to determine if there were cofactors that would improve creatine utilization by the body. They first mixed creatine with other nutrients, such as carbohydrates, to see if the insulin boost would increase creatine uptake and delivery to the muscle fibers, and it did. Studies by A. L. Green, in 1996, demonstrated that combining creatine with a simple carbohydrate, such as glucose, will increase creatine transport into the muscle. The solution tested consisted of 5 grams of creatine and 90 grams of glucose, consumed four times per day. When tested against creatine alone, which increased total muscle creatine and creatine phosphate levels, the creatine carbohydrate supplement increased total muscle creatine and creatine phosphate levels significantly more. However, the overconsumption of simple carbohydrates can have detrimental effects on your lean muscle-building progress, as you learned from reading about Macrobiotic Nutrition in previous chapters. This is one example where blindly jumping on the scientific bandwagon is not justified. Bodybuilders and strength athletes who use sugar-loaded creatine add more than 150 grams of pure sugar to their diets per day in the loading phases alone! That amounts to 600 daily calories from sugar alone, and equals the amount of sugar found in four cans of soda! Does that sound like the kind of nutrition you want to feed *your* body? There are other ways to make creatine delivery more effective without overloading the body with simple carbohydrates that only serve to increase your insulin levels too much, as you shall see shortly.

I also wanted to determine if there were ways of reducing the conversion of creatine to creatinine, which is basically useless. Ironically, acid conditions like those found in the stomach stimulate the conversion of creatine to creatinine. This is one reason some researchers have believed that large amounts of creatine had to be ingested to see beneficial results. In fact, a few interesting insights have revealed how creatine behaves in the body. A large portion of the creatine you ingest gets converted to creatinine in the stomach. Additionally, your body can only effectively absorb about 2 grams of creatine into the bloodstream from the intestines, and any excess creatine is excreted. Other nutrients can help to maximize creatine's effects and boost creatine's content in your muscle fibers once creatine is in your body, however.

This information led to the development of MHP's unique proprietary TRT™ (Time Release Technology) microencapsulation process, which pro-

duces a unique release profile for its creatine monohydrate and other ingredients. This new delivery system accomplished these benefits for creatine users: the microencapsulation protects the creatine from being converted to creatinine in your stomach. In your intestines, the creatine is time released to allow for maximum utilization and minimum waste. The timed release allows for convenience of use, because you need to take only one serving per day rather than several servings per day. This technological breakthrough led me to name the MHP product TRAC—Time Released Arginine/Creatine. As the name reveals, arginine is one of the key creatine cofactors that I found will produce additional benefits for creatine supplementation. Early in my research efforts I discovered that besides providing additional benefits of its own, arginine can help boost the effectiveness of creatine. I was the first to promote the use of arginine for benefits associated with improved nitric oxide production. While creatine helps promote muscle-cell volume, size, and strength, nitric oxide has a major influence on muscle physiology, because it increases vasodilation and nutrient uptake. The effect of nitric oxide's action is to help accelerate muscle velocity, amplify power output, improve recovery, and most amazingly, stimulate new muscle-fiber production.

New research shows ADNO (arginine-derived nitric oxide) to have powerful insulin-mediating effects. TRAC contains 4 grams of arginine in order to increase the production of nitric oxide (NO) and stimulate insulin output and sensitivity *without fattening sugar*. This new patent-pending approach to mediate insulin is called “nitro-loading,” and it delivers much needed nutrients to your muscles for increased size, strength, and improved recovery. (See Figure 9.1.)




Figure 9.1. Dose Response Curve for Oral L-arginine

The arginine drives insulin and leg blood flow, causing an increase in muscle perfusion and transport of creatine to muscle. A large enough dose of arginine is necessary to push nitric oxide production enough to mediate insulin and increase blood flow.

TRAC employs the patented proprietary microencapsulation process, TRT, to produce a unique release profile for its main components. The arginine and creatine are sustained release, but the L-arginine release precedes the creatine release. As can be seen in Figure 9.2, the L-arginine release is about 64 percent at the two-hour time point, whereas the creatine release is 43 percent at the same point. At the four-hour time point, the L-arginine release is about 80 percent, while the creatine release is around 66 percent. At six hours, the L-arginine release is basically complete, but the creatine is still releasing and is about 82 percent. The arginine release is always ahead of the creatine to provide the *bioactive shuttle*, stimulating IGF-1 production and facilitating the sodium/potassium channel, to increase creatine production, transport, and preservation. (See Figure 9.2.)




Figure 9.2. TRAC Sustained Release Profile

TRAC contains other synergistic ingredients, including one of my favorite new sports ingredients, referred to as NADH. NADH (coenzyme nicotinamide adenine dinucleotide) is present in every living cell, and is a vital biochemical that is required for production of cellular energy. NADH is involved in the production of ATP. Some research has shown that supplemental intake of NADH can also improve mental function. NADH increases the production of the neurotransmitter dopamine, which plays a role in short-term memory, concentration, and muscle function. NADH also enhances the synthesis of norepinephrine, which functions in alertness, concentration, and mental activ-

ity. Research on athletes indicates that NADH supplementation enhances work capacity. Oxygen uptake and reaction time was also observed to improve in the athletes tested. The possible increase of dopamine levels might explain improvements in reaction time, and increases in performance could be due to NADH's role in energy production. Combining NADH with creatine makes TRAC an "ATP Power Plant" for explosive strength.

A final word about how I use TRAC. I take a serving of TRAC thirty to forty-five minutes before workouts on training days, usually after work around 6:00 P.M. I also take one serving in the early evening on nontraining days to maintain muscle-creatine saturation.

Glutamine

While following the Macrobiotic Nutrition program will result in the dietary intake of glutamine and other amino acids your body needs for lean muscle mass growth and performance, numerous research studies indicate that taking extra glutamine supplements can provide you with benefits above and beyond diet alone. This is especially true if you are exercising or training intensely. In addition to glutamine's benefits, which will be reviewed in this section, I find it interesting that even the military has turned its attention to the way that glutamine supplements can help improve military personnel performance. In addition to the role of glutamine in supporting immunologic defenses, a 1999 military research review by the Institute of Medicine mentioned other benefits of taking glutamine, notably: glutamine supports skeletal muscle protein synthesis; it enhances bicarbonate production, which may neutralize the acid load generated by moderate to severe exercise or catabolism; and glutamine supports glutathione synthesis, an antioxidant that attenuates the tissue damage associated with free-radical production. When it comes to improving health and physical performance the military is right on in identifying these key roles of glutamine.

Glutamine's multifunctional role and its involvement in various biochemical processes makes it one of the most important amino acids for athletes and individuals interested in maximum physique development, performance, and general health. It is a neurotransmitter and an energy source in the brain. Glutamine is a mediator of glutamic acid and GABA activity and can be converted back to glutamic acid in the brain, where it is essential for brain function. Glutamine is also vital to immunity function. New studies show that cell replication in the immune system requires glutamine. Most glutamine is stored in the muscles, however, so your muscles have to supply a large amount of glutamine to the immune system. Supplemental use of L-glutamine by athletes is known to have a strong anticatabolic effect that neutralizes the highly cata-

bolic cortisol generated by strenuous exercise. Its anticatabolic action allows anabolism (muscle building) to take place more efficiently. L-glutamine also plays an active role in the recovery and healing process.

First generation glutamine sports nutrition products contained free-form L-glutamine, which research studies report works well. However, after reviewing the scientific research and making observations of my own, it became apparent that many of glutamine's effects can be potentiated by increasing its bioavailability and absorption. L-glutamine is very susceptible to the acid environment of the stomach, and MHP developed Effervescent Glutamine to deal with this issue. MHP's commitment to research and our willingness to go the extra mile has led to the development of some innovative and incredibly effective products. Effervescent Glutamine is at the top of this list. Studies show the utilization of a pharmaceutical-grade bicarbonate delivery system can improve glutamine uptake by almost 400 percent. Recognizing this, MHP's team of biochemists set out to find a way to neutralize the acidic gastric medium and improve bioavailability, and subsequently developed a glutamine product with an advanced bicarbonate-buffering system. Utilizing a pharmaceutically proven source for gastric alkalization (sodium bicarbonate), MHP's Effervescent Glutamine takes advantage of the acid neutralization power of the bicarbonate ion (HCO_3^-). Bicarbonate bonds with excess hydrogen ions produced by the HCL acid to form carbon dioxide and water. This process allows a buffered state to exist in the stomach, permitting more L-glutamine to pass into the intestines where it can be absorbed and utilized by intestinal cells for increased glutamine delivery to muscles. It is interesting to note that bicarbonate has been clinically proven to help increase strength and power of athletic performance on its own, making it a truly synergistic ingredient to team up with L-glutamine.

I take two servings of Effervescent Glutamine a day to ensure optimum glutamine levels, avoid catabolism, and stimulate growth hormone (GH) production. I take one serving in the morning, between my first and second meal, and the second immediately following workouts or in the early evening on nontraining days.

Testosterone Optimizers

Exploring ways of optimizing the body's production of testosterone is one of the oldest approaches to enhancing athletic performance and muscle growth. A primary goal of all bodybuilders and other strength athletes is to increase testosterone levels and optimize the anabolic effects of testosterone. MHP has focused its research efforts in two primary directions in this area:

1. providing the athlete with natural testosteronelike “prohormone” substances that are more anabolic than testosterone
2. developing a product that will optimize the body’s testosterone production and utilization

T-BOMB

MHP research and expertise on prohormones resulted in what I believe is the most powerful anabolic substance legally available in the sports nutrition market today. Its chemical name is 17-beta-hydroxy-5-alpha-androst-1-ene-3-one tetrahydropyranyl ether, also known as 1-Testosterone. The fact is that 1-Testosterone has been proven to be about seven times more anabolic than the testosterone your body makes. Another benefit of 1-Testosterone is that it does not convert to estrogens in your body. There was one major challenge to overcome to make MHP’s 1-Testosterone the most powerful legal anabolic supplement available, and this involved stabilizing the 1-Testosterone substance in a tablet formulation for oral ingestion. MHP’s research and development team found the solution by applying a special pharmaceutical enteric coating on the tablet. This enteric coating protects the 1-Testosterone from degradation while it is in the stomach, thereby ensuring optimal stability and maximum potency. This 1-Testosterone product is so potent MHP named the finished product T-Bomb. (See Figure 9.3.)

Suggested use: 1 tablet twice daily. T-BOMB should be cycled for two months on and one month off. You may repeat the cycle on an ongoing basis.

Note: As with all testosterone formulas, T-BOMB should be used only by




Figure 9.3. The Stability of 1-Testosterone in Enteric-Coated Tablet versus Capsule

males over the age of eighteen, and is not recommended for women or minors. Always consult your physician before taking any supplement.

T-BOMB II

Often the greatest discoveries are inspired by necessity. This is the case with MHP's T-BOMB II. The category of prohormones has been under scrutiny by the media as to whether the hormonelike compounds should be considered dietary supplements or drugs. I personally think that this scrutiny is unjust. It seems that any time supplements provide "very measurable results," they become subject to attack by the press, which forces the FDA to respond.

In any event, I decided to be proactive and looked to develop a "pro-hormone free" testosterone formula that could parallel the effectiveness of T-BOMB. This was not going to be an easy task. T-BOMB definitely set a standard that was going to be difficult to match. Hundreds of thousands of satisfied T-BOMB users reported exceptional gains from T-BOMB, making it the number-one-selling prohormone formula in America.

I knew I would need to dig my heels into some research if I was going to develop a "sequel" equal to T-BOMB. After researching thousands of compounds and examining the way they worked in the body (and consulting with scientists and endocrinologists), it became evident to me that all of the products on the market were way off the mark. Let me explain: Though some of these products do contain effective testosterone-boosting ingredients, none of them addressed the full complexity of hormone regulation by the endocrine system. The endocrine system is a complex network of checks and balances. Testosterone elevation triggers other important hormonal responses, all of which must be considered. I discovered through my research that if I was going to develop a supplement to elevate testosterone, it must also address estrogen, 5 alpha-reductase, DHT, and sex-hormone-binding globulin (SHBG).

Note: These other hormonal responses (estrogen, DHT, and SHBG) did not have to be addressed with T-BOMB, because of 1-Testosterone's unique chemical structure. Though 1-Testosterone is anabolic and attaches to the androgen receptor, it does not convert to estrogen or DHT (the testosterone derivative associated with prostate enlargement). However, testosterone does make this conversion, which is why these factors must be addressed if you elevate testosterone.

Now I'm going to explain why all of these hormones come into play. As testosterone production increases, an active enzyme called *aromatase* converts a portion of testosterone to estrogen. As you already know, a high level of estrogen is definitely not what a guy wants, especially a bodybuilder. Estrogen

causes increased body fat, water retention, and gynecomastia (excessive development of breasts in men). If that isn't bad enough, estrogen also stimulates the production of sex-hormone-binding globulin (SHBG), which binds testosterone. This binding process lowers free testosterone making less available to attach to the androgen receptor where it can exert its effects on sex functions and muscle building. Another enzyme, 5 alpha-reductase, converts a portion of testosterone to DHT (5-dihydrotestosterone). This compound is directly related to hair loss and prostatic hypertrophy, and can also occupy the receptor for testosterone and inhibit free testosterone from attaching. You can clearly see why you want to keep estrogen and DHT levels down.

T-BOMB II is the first formula to contain select key ingredients to do the following:

1. Increase the production of testosterone by stimulating the production of luteinizing hormone
2. Reduce DHT
3. Reduce estrogen
4. Reduce SHBG

A New Frontier—Signal Transduction: Second Messengers “Amplifying the Signal at the Receptor”

Aside from being the first and only testosterone formula to optimize endocrine functions by improving the testosterone-to-estrogen ratio and maximizing the anabolic effects of testosterone, T-BOMB II introduces a new frontier called “second messengers.” This new advance in nutrition technology is going to be the biggest bodybuilding breakthrough since anabolic steroids.

In order to get the full anabolic benefits of testosterone, or any hormone, steroid or prohormone, it must first attach to the receptor cells in the body. But, so-called second messengers are responsible for actually carrying out the hormone's tasks. In other words, the binding of a hormone to the receptor initiates a series of events, which lead to the generation of second messengers within the cell (the hormone is the first messenger). The second messengers then trigger a series of molecular interactions that alter the physiological state of the target cell. This process is called “signal transduction.” It sounds complicated, because it is a very complicated system.

Simply stated, second messengers amplify the signal at the receptor from a specific hormone. Therefore, a small amount of a given hormone will result in a strong hormonal response. This is why I refer to our proprietary complex as

“Second Messenger Hormone Amplifiers.” Increasing signal transduction at the receptor makes the given amount of hormones like testosterone and growth hormone circulating in your body and attaching to the specific cell receptors more powerful. T-BOMB II’s proprietary “Second Messenger Hormone Amplifier Complex” further increases the anabolic and androgenic effects of elevated testosterone.

T-BOMB II is by far the most advanced testosterone-booster/hormone-optimizing formula to date. The proper manipulation of endocrine function is critical if you want to improve the ratio of testosterone to estrogen. This can be achieved only by using the right amounts and combination of synergistic ingredients that are present in T-BOMB II, coupled with the groundbreaking discovery of signal transduction and second messenger hormone amplifiers. If you’re looking for an anabolic edge, try T-BOMB II.

Suggested use: 3 tablets twice daily. Since T-BOMB II improves hormonal homeostasis in a natural fashion, you do not have to cycle this product. However, T-BOMB II can also be cycled for three months on and one month off and then repeat cycle.

Note: As with any testosterone formula, T-BOMB II is recommended only for males over the age of eighteen, and is not recommended for women or minors. Always consult your physician before taking any supplement.

Growth Hormone and IGF-1 Enhancers

Next in line after testosterone/anabolic steroid alternatives are human growth-hormone stimulators, which are also included in the story of how I founded MHP. As a competitive bodybuilder, I looked for every possible way to improve my progress in gaining lean muscle mass. Human growth hormone caught my attention when I looked at the medical research reports on lean body mass-enhancing substances. My path for finding natural growth-hormone stimulators led me to Vincent Giampapa, M.D. Dr. Giampapa was involved in some intriguing research involving a unique natural growth-hormone-stimulating product he had developed to use in his clinic, which specializes in antiaging medicine and hormone replacement therapy.

Dr. Giampapa introduced me to his natural product and explained how he was using it in his clinic to optimize growth hormone levels in some of his older patients for antiaging purposes. What led him in this direction was the impressive research on the antiaging (youth-enhancing) benefits of human growth hormone. Human growth hormone or HGH (also known as somatotropin) is produced by the pituitary gland, and is essential for growth and regeneration of cells and tissues. Growth hormone has both anabolic and fat-

burning effects, so it's easy to see why athletes and bodybuilders try to enhance their HGH levels.

I learned a lot about HGH while working with Dr. Giampapa during this period, both from sorting through the medical research and from experimentation. The body's production of HGH is very high during childhood and adolescence. Unfortunately, after age twenty, the pituitary gland's release of HGH falls at a rate of approximately 14 percent every ten years. A loss of 75 percent or more of HGH is not uncommon by age sixty. The usual physical symptoms of aging, such as wrinkles, increased body fat, loss of muscle mass, loss of energy, and other signs are directly related to this decrease in the production of HGH.

Research on stimulating the production of growth hormone first started with children who had inadequate growth hormone production. Researchers examined ways of increasing growth hormone production with natural and synthetic substances, and amino acids were found to be the most effective and promising substance. Some of the amino acids would actually increase the body's production of growth hormone. Yet other amino acids helped block the breakdown of the growth hormone produced by the body. From your reading so far, it is obvious that most amino acids have a few to several functions in the body. In addition to their other beneficial effects, some amino acids you've already read about, such as L-glutamine and L-arginine, are natural growth-hormone stimulators. Once researchers determined that these growth hormone-stimulating amino acids, also referred to as GH secretagogues, worked in children, they started to see how they worked in adults of all ages and activity levels. This massive research effort revealed that these GH secretagogues benefited both athletic and nonathletic adults. Of particular interest is that they benefit bodybuilding, by helping stimulate lean muscle-mass development.

But in 1990, when *The New England Journal of Medicine* published the results of a study by Daniel Rudman, M.D., the world turned its attention to the many potential benefits of HGH. Dr. Rudman led a research team that examined the effects of administering biosynthetic human growth hormone to healthy men age sixty-one to eighty-one years old with low levels of IGF-1. Note that IGF-1 is related to HGH concentrations in the human body, so when HGH levels in the body increase, IGF-1 levels also increase. The biosynthetic HGH was administered to the experimental group for six months. At the end of the study it was determined that blood levels of IGH-1 rose to that normal for a youthful range in the group of people taking the biosynthetic HGH. This would be expected from taking the biosynthetic HGH, but what shocked the medical world were the results of the increased IGF-1 levels in

these older men. They increased lean body mass by about 9 percent, reduced body fat by about 14 percent, and also increased their bone density. Wow, taking this stuff really did result in a “fountain of youth” effect.

Of course, the results of the Rudman study attracted the interest of athletes, who want to increase their lean muscle mass and reduce excess body fat. It also stimulated interest in the medical community to use synthetic HGH or promote natural levels of HGH for improved general health, well-being, and longevity. Since Dr. Rudman’s landmark research, hundreds of studies have been undertaken to explore the benefits of taking synthetic HGH and of using natural HGH stimulators, including the studies by Robert Goldman, M.D., and Ronald Klatz, M.D., who are authors, researchers, and founders of the American Academy of Anti-Aging Medicine. They note the following benefits that HGH has on the body:

- Reduced body fat
- Increased muscle mass
- Higher energy levels
- Faster wound healing
- Regrowth of hair
- Elevated mood
- Stronger bones
- Restoration of youthful immune function
- Lower cholesterol and blood pressure
- Enhanced sexual performance
- Regrowth of vital organs
- Smoother, firmer skin
- Improved cognition
- Sharper vision

Back to Dr. Giampapa. Once I brought myself up to date on HGH, I better appreciated why Dr. Giampapa and others were so enthralled with using HGH for antiaging, enhancing physique, and improving athletic performance. Dr. Giampapa actually created a natural, amino-acid-based HGH-stimulating formula to use in his clinic. I expressed my interest in putting this natural HGH-stimulating formula to the test by using it myself. Dr. Giampapa agreed, and established a monitoring program to track my results. To my surprise and delight, Dr. Giampapa measured a significant increase in my natural HGH and IGF-1 levels. My IGF-1 levels increased by 42 percent. Benefits to my physique became measurable after just a few weeks, and I began to increase muscle mass and decrease body fat at a faster rate than I could before taking the amino acid product.

I was blown away by these results. It is a rare occurrence for a bodybuilder at my level to be able to achieve such noticeable results in a short period of time by taking natural substances. I asked Dr. Giampapa if I could license his formula and bring it to the market for bodybuilders and other athletes, and he

agreed. I incorporated MHP and launched Secretagogue-One in January 1998. The rest is history. Secretagogue-One remains the number-one-selling HGH-releasing formula in retail stores to date—because it works. You get what you pay for in this world and I believe in making an effective, quality product.

From this experience working with Dr. Giampapa, I established the foundation for MHP product development. MHP products are scientifically developed, medically validated, and bodybuilder tested to produce results fast.

As with other MHP product development, we relied on Dr. Giampapa's assistance during the product development and verification process for Secretagogue-One. Dr. Giampapa went to work in his own clinic to verify that Secretagogue-One was the best he ever used. He confirmed that it is a breakthrough in natural hormonal manipulation, designed to support the body in releasing more of its own HGH through the use of secretagogues and other precursors. Coupled with an effervescent delivery system to improve absorption, these secretagogues are delivered in a proprietary "Glucose Polymer Matrix," *providing nearly 100 percent assimilation*, as opposed to only 10 to 15 percent absorption for competing secretagogue products.

Secretagogue-One activates the pituitary gland naturally, and allows for optimum hormonal levels, while maintaining homeostasis. It is nontoxic and *does not* contain synthetic hormones, which can weaken your endocrine system. Secretagogue-One allows your body to make more of its own HGH in a balanced and natural way.

I use Secretagogue-One every day. I take one serving before bedtime on an empty stomach. This is the best way for bodybuilders and other athletes to take advantage of the natural nighttime high that occurs in HGH and IGF-1 levels, which Secretagogue-One increases even more. Note that women also produce HGH. Though HGH is anabolic, it does not have any androgenic (masculizing) effects. So, HGH supplementation is also excellent for women.

Joint Health and Reduction of Exercise-Induced Pain and Inflammation

No pain, no gain? Think again. Muscle-building progressive-resistance training requires overloading the muscles to a point where muscle fiber damage is imminent. The result is pain and inflammation, and muscle growth. Exercise physiologists use the phrase DOMS (delayed-onset muscle soreness) to describe this exercise-induced muscle pain. As an iron-pumping athlete, I wanted to discover what natural solutions existed beyond anything currently available to help relieve this exercise-induced muscle soreness. The many years of extensive training has done its share of wear and tear on my joints. I tried various types of joint remedies, such as MSM and chondroitin, and though I had some ben-

efits, none of these products seem to address the pain and inflammation I was experiencing. I wanted to formulate a product that helped repair and minimize the pain and inflammatory response. It would then be possible to decrease recovery time physiologically and allow for greater progress. Psychologically, reducing or eliminating exercise-induced pain and inflammation translates into more frequent, better, and more pleasurable workouts. Breaking the pain barrier allows you to maximize your potential.

In search of the answer, I turned my attention to areas of medicine that have a track record of dealing with similar situations. In particular, my journey led me to the area of medicine that deals with combating the pain and inflammation of arthritis. The body tissues, especially the connective tissues of people with arthritis, are being broken down in a process that is accompanied by much pain and inflammation. Upon examination, exercise has a similar effect on the body and will eventually wear down even the healthiest joints if it is unchecked.

I began to experiment with different substances, including supplements and over-the-counter drugs like ibuprofen, in this research effort. In fact, a recent research study published in the *Journal of Strength and Conditioning Research* by S. P. Tokmakidis of the Department of Physical Education & Sports Sciences, Demoritus University of Thrace in Greece, reported on benefits of this drug. It was determined that taking ibuprofen helped reduce muscle soreness after exercise (weight resistance against muscle in the lowering motion). This motion is commonly referred to as negative rep or movement. But turning back to the lessons from my research in arthritis, I focused my attention on the inner workings of a special class of nonsteroidal anti-inflammatory drugs (NSAIDs), called COX-2 inhibitors (COX stands for cyclooxygenases).

Cyclooxygenases are enzymes needed for the synthesis of hormonelike substances called prostaglandins. There are two types of cyclooxygenases: the COX-2 enzyme that mediates inflammation and pain, and the COX-1 enzyme that helps maintain other physiological functions in the body. Traditional NSAIDs inhibit both enzymes. The new NSAIDs, however, mostly block the COX-2 enzyme, offering a new treatment option for people who have difficulty tolerating the old NSAIDs. The FDA approved the first COX-2 inhibitor, Celebrex (celecoxib), in 1998 to treat rheumatoid arthritis and osteoarthritis. Vioxx (refecoxib) became the second COX-2 inhibitor to receive approval, in 1999, for the treatment of osteoarthritis, dysmenorrhea (pain with menstrual periods), and the relief of acute pain in adults, such as that caused by dental surgery. I was amazed to find that these two are among the most frequently prescribed drugs, with billions and billion of dollars per year being spent on them. COX-2 inhibition is something people look for, sooner or later.

In the last ten years, this COX-2 paradigm has emerged as one of the most important biochemical systems of the twenty-first century. The COX-2 enzyme has become a primary model for understanding and treating numerous different conditions, including pain and inflammation. Researchers found that pain and inflammation can be dramatically reduced by inhibiting COX-2. Reducing the pain and inflammation also allows the body more resources to help speed up the muscle and connective tissue-rebuilding process. COX-2 inhibitors should be part of the daily routine of athletes and bodybuilders, who put a tremendous amount of stress and strain on their joints and cartilage. During and after every workout, damaging COX-2 is being created or induced. Daily use of COX-2 inhibitors can arrest this process and provide for long-lasting joint and overall body health.

In my search for the ideal natural COX-2 inhibitor, I reviewed the research and tried a variety of botanicals traditionally used for relieving pain and inflammation, but which also had modern scientific research proving their efficacy. I found two botanicals, boswellia and turmeric, that worked okay, but then I came across a new natural extract that boasted almost 100 percent selective COX-2 activity with almost no COX-1-inhibiting activity. This substance is called IsoOxygene™, and from the first day I used it, I experienced the best possible pain-relieving and anti-inflammatory effects ever, and I have been using it ever since.

In studies with cells, IsoOxygene has been shown to significantly reduce the production of PGE-2 produced by the COX-2 enzyme when stimulated by a mediator or cytokine. In comparison tests, IsoOxygene at a fairly low concentration was shown to reduce this degenerative chemical (PGE-2) to virtually zero. Other botanicals typically used for this purpose were inferior in both selectivity for the COX-2 enzyme, and in potency or efficacy for reducing PGE-2. Generally speaking, experts in the field agree that for an ingredient to be considered a true COX-2 inhibitor, it should selectively inhibit the COX-2 form of the enzyme five times more than the COX-1 form of the enzyme. None of the other botanical substances tested came close to the thirty times more selective ingredient in MHP's joint formula. The nearest commonly used botanical, curcumin, had a selectivity ratio of COX-1:COX-2 of about 1 to 2.8, while IsoOxygene has a selectivity ratio of 1 to 30. IsoOxygene is patent pending, which will eliminate would-be imitators and ensure authenticity of products claiming to contain this precious COX-2 inhibiting ingredient. (See Figures 9.4, 9.5, and 9.6.)

With the COX-2-inhibiting ingredient taken care of, my attention focused on including a key ingredient clinically proven to be the champion of connective-tissue rebuilding. After sorting through the research I started to




Figure 9.4. Botanical COX-2 Inhibitor Comparison

COX-2 inhibition as measured by inhibition of PGE2, a pro-inflammatory prostaglandin, in RAW 264.7 cells. The IC50 is the amount of the ingredient needed to reduce PGE2 by 50 percent.

Nexrutine™ is from Next Pharmaceuticals. Curcumin is a commonly used botanical anti-inflammatory ingredient with COX-2 to COX-1 ratios of 2:1, resveratrol is a phytoalexin found in grapes and other botanicals. IsoOxygene™ is the proprietary new COX-2 specific inhibitor. As can be seen from this graph, a much lower concentration of IsoOxygene is needed to reduce PGE2 by 50 percent than the other botanicals. Therefore, it is a much more potent COX-2 inhibitor.

Botanical COX-2 inhibitors such as curcumin have a COX-1:COX-2 ratio of about 1 to 2.8, which is called the COX-2 specificity. IsoOxygene has a COX-2 specificity of about 30.




Figure 9.5. COX-2 Inhibitor Comparison: Whole Blood Assays

The potencies of the various drugs were also tested according to the William Harvey Human Modified Whole Blood Assay (WHMA) at the William Harvey Research Institute, Royal London School of Medicine in the United Kingdom.

The IC50 is in micromoles or 1×10^{-6} moles. The lower the amount (concentration) necessary to inhibit PGE2 by 50 percent (IC50), the more potent the drug. The more potent the drug, the lower the effective dose will be for reducing COX-2 activity.




Figure 9.6. Selectivity for Cox-2 (WHWBA Data)

When compared to the OTC pain relievers aspirin, ibuprofen, and naproxen, IsoOxygene™ appears to be more selective for COX-2 than COX-1 and has a lower IC50 value for COX-2 than any of the OTC pain relievers. The selectivity for COX-2 exhibited by IsoOxygene appears to be comparable to the prescription COX-2 pain relievers Rofecoxib and Celecoxib. Conversely, the COX-1/COX-2 IC50 ratio for IsoOxygene using the WHMA protocol is about 92.85, which qualifies it as a selective COX-2 inhibitor in the topmost category of those select compounds that have greater than fifty-fold selectivity for COX-2 over COX-1. Since IsoOxygene inhibits COX-2 at the level of gene transcription, the results in the A549 cells were even more significant than in blood alone.

examine how I personally responded to the top candidate ingredients supported by good research studies: chondroitin sulfate, MSM, and glucosamine sulfate. Glucosamine sulfate was my first choice, following several months of comparison and successful use, and because it was the most clinically proven of the three.

What really convinced me to use glucosamine sulfate in my new MHP products was, in addition to its being the most clinically proven for rebuilding connective tissues in people with arthritis, my discovery of a study in which glucosamine sulfate was effective in young athletes with athletic-induced wear and tear of the cartilage of the knee. A group of researchers in Germany headed by D. Bohmer examined the effects of glucosamine sulfate in young athletes with an exercise-induced knee problem. In the young athletes taking the glucosamine supplements, the knee pain was almost gone after just four weeks, and completely gone after twelve weeks of supplementation. Glucosamine's repair and rebuilding of the knees of young athletes' connective tissues allowed them to recover to the point of resuming an intense-training program.

A short review of glucosamine and related topics will explain how it works in your body to produce numerous benefits. There are several types of connective tissues. Cartilage, tendons, ligaments, intervertebral discs, the pads between joints, and cellular membranes are all composed of connective tissue. All connective tissues have two common components, chief of which is collagen. One-third of your body's total protein volume is composed of collagen, making it the most common protein in the body. The other component is proteoglycans (PGs). PGs form the "framework" for collagenous tissue. These huge structural "macromolecules" are composed mainly of glycosaminoglycans (GAGs), which are long chains of modified sugars. The principal sugar in PGs is hyaluronic acid, 50 percent of which is composed of glucosamine. The principal amino acids that form collagen are proline, glycine, and lysine. Collagen and PGs must somehow "get together" during the production of new connective tissue. Of the multitude of biochemical reactions that take place during the synthesis of connective tissue, there is one critical "rate-limiting" step, which, once reached, guarantees that new connective tissue will be successfully synthesized. That rate-limiting step is the conversion of glucose to glucosamine, so glucosamine is the single most important substance in the synthesis of connective tissue. Over thirty years of research has gone into understanding how glucosamine acts as the precursor of GAG synthesis. Scientists have long known that simply ingesting purified glucosamine from connective tissue allows the body to bypass the critical rate-limiting step of converting glucose to glucosamine. In human clinical trials, glucosamine taken orally was observed to initiate a reversal of degenerative osteoarthritis of the knee after two months. Normalization of cartilage was documented by taking biopsies of the tissues and scrutinizing them with an electron microscope. Of greater interest to athletes, glucosamine aids in feeding GAG to your injured connective tissues. GAG is the most critical precursor for rebuilding the collagenous matrix, which forms connective tissue; glucosamine is the preferred substance in synthesizing PGs, your connective tissue's framework. Glucosamine as a supplement clearly aids in connective-tissue synthesis. All athletes need such a substance, because the repair and growth of connective tissue is never-ending.

After more use of this prototype formula, I found that adding a few more ingredients improved the total effects. The formula was further enhanced with baikal skullcap, N-acetylcysteine, and alpha lipoic acid, a synergistic combination of ingredients that work together to provide fast-acting relief and anti-inflammatory action.

The resultant formula worked so well to reduce exercise-induced pain and inflammation, I decided to call it RELEVE. In order to stimulate muscle

growth, you have to train heavy and hard. This training is sure to cause joint pain, inflammation, and/or injury over time. If you're looking to make serious progress in your training without suffering debilitating soreness, consider trying RELEVE.

GETTING LEANER FASTER

Meal-Replacement Powders (MRPs)

As was previously discussed, a primary goal of Macrobiotic Nutrition is to give your body the tools it needs to grow lean muscle. You have learned the enormous impact food has on your body, and how Macrobiotic Nutrition is scientifically developed and bodybuilder-tested to provide the nutrients you need for optimum lean muscle growth.

However, putting any nutrition plan into practice is not always as easy as one, two, three. While this book contains thorough nutrition plans and convenient recipes, you don't always have the time to plan and prepare meals and snacks that meet the Macrobiotic Nutrition criteria. The same is true for me, too. That's why I developed the Up Your MASS products. The first generation of Up Your MASS products consists of the Up Your MASS Muscle Building Macrobiotic Nutrition Formula in the form of nutrient dense, delicious meal-replacement powders. The Up Your MASS formulations are no ordinary MRPs, as you will soon discover from the following overview of how they were developed and of their special muscle-building characteristics.

Meal-replacement powders have a long history of use in the United States to help people with weight control and to serve as a quick meal or meal/snack on the go. The MRPs you are most familiar with are the low-calorie mass-market products used by millions to get slim fast. These and the so-called more advanced MRPs designed for athletes fall short on quality nutrition. On the surface, the mass-market MRPs offer some weight-loss benefits for the general population. However, when you examine these products with a more sophisticated eye, they are nothing more than sugar drinks, with a minimal amount of protein and essential vitamins and minerals. The weight-loss MRPs are definitely not the first choice for anyone, especially the health-conscious athlete. Although using these mass-market MRPs may result in weight loss from a reduction in daily calorie intake, there is no focus on the quality of the weight lost and the future negative impact that these unbalanced, high sugar content, poor-quality nutrition products may have on your health. The sports nutrition MRPs are not much better. When you examine sports nutrition MRP products, especially those in the weight-gain category, you will commonly discover that many of these products are loaded with sugar and are deficient in the

protein-content department. These formulas are also devoid of another major group of macronutrients that you have learned are very essential: carbohydrates and fiber.

So, to keep us on track with our Macrobiotic Nutrition plans, I developed Up Your MASS meal replacement powders, which are based on the 45/35/20 lean-mass equation. Following is a summary of some of the exciting lean muscle-building, nutritional technological breakthroughs found in the Up Your MASS MRPs.

The first goal was to develop a natural-based, low-glycemic-index blend of carbohydrates, because carbohydrates are the primary source of energy and are vital to exercise performance and health. To accomplish this daunting task, I directed my research team to conduct a rigorous examination of the scientific research. We discovered that the secret was to develop a virtually sugar-free natural-based blend of low-glycemic-index carbohydrates from barley, oats, and oat fiber. I call this low-glycemic-index carbohydrate blend Glycemix LGI™. When you ingest MHP products containing Glycemix LGI, you get a sustained release of energy into your body, and keep your blood-sugar levels in the anabolic zone.

In addition to increasing sustained energy, Glycemix LGI increases glycogen storage, stabilizes blood sugar levels, and improves your “hormonal profile” of glucagon and insulin, for accelerated muscle building and fat burning. Glycemix LGI is also high in heart-healthy fiber. The special soluble fibers contained in Glycemix LGI help to lower your cholesterol levels, and may even help reduce your risk of developing cardiovascular diseases. Fiber also helps prolong the supply of nutrients to muscle.

The second goal was to design a protein blend that works best for muscle building. The fact is that all proteins are not created equal. Furthermore, your body experiences different protein needs, based not only on the extent of physical activity, but also on the type of physical activity or exercise. This means that weight-lifting, lean muscle-building athletes have different protein needs, or more exactly, different amino needs. After years of researching and experimenting to learn how to best meet the protein/amino acid needs for optimal lean muscle building, MHP’s Probolic protein was developed. Probolic, as discussed in Chapter 3, is a proprietary amino acid-enhanced, sustained-release blend of high-quality proteins sources including proteins derived from choice, high-quality whey concentrate, casein, and soy protein isolates, fortified with crucial amino acids. As Figure 3.1 and Table 3.10 in Chapter 3 illustrate, this makes Probolic win in amino acid content, including arginine, BCAAs, and glutamine. The other key development was to make Probolic contain the ideal amino acid profile while displaying the precise release rate. This dynamic

protein-delivery matrix thus provides a fast, medium, and slow release to feed your muscles over hours and help to improve nitrogen balance, growth, and recovery.

The third major goal was to develop a precise blend of lipids that best meet the Macrobiotic Nutrition criteria, which demanded the inclusion of healthy, performance-enhancing fatty acids. This was no easy task, as there are many variables to conquer when attempting to contain healthy lipids in a dry, meal-replacement powder environment. The resulting Lipid Complex™ includes some of the healthiest and most sought-after lipids on the market today, such as conjugated linoleic acid (CLA), borage seed-oil powder, evening primrose oil, flaxseed powder, omega-3s, and medium-chain triglycerides. The Lipid Complex is, therefore, a synergistic blend of lipids that are vital for muscle growth, promote a desirable hormonal balance, improve nitrogen balance, provide the body with a high-energy source, and help stabilize insulin release.

All of this nutrition research and technology has resulted in making the Up Your MASS meal-replacement products a major advancement in lean muscle-building science. I usually take two servings of Up Your MASS a day; one between breakfast and lunch, and another about one hour before workouts. I find that this helps fuel my muscles during my grueling workouts, as well as being an easy-to-digest premeal workout. For added convenience, MHP has also developed Up Your MASS nutrition bars. They are also nutrient dense and based on the 45/35/20 lean-mass equation. These bars taste awesome and are great on-the-go meals or snacks.

Thermogenic Fat-Loss Aids

In addition to wanting faster improvements in muscle size and strength, most of the people I encounter also want to get leaner faster. While following the Macrobiotic Nutrition program will intrinsically get you leaner, you may want to move this process along at a more rapid rate. If so, this section will bring you up-to-date on what has been termed the thermogenic fat-loss aids.

When you read this and other sections of this book regarding losing body fat, you will note that losing too much weight too fast is not the healthy way, nor is it the Macrobiotic Nutrition way. A safe rate is a few pounds of *fat loss* a week. Don't confuse weight loss with fat loss. Fat loss is what you want, and not at the expense of losing any lean body mass, which includes muscle mass. When you go on a fat-loss program that results in losing too much weight too fast, most of this fast weight loss comes from water and muscle mass. My intention here is not to turn your fat-loss efforts into a race, but for you to realize that sometimes your body needs a boost in its fat-burning metabolism to better release fatty acids from body fat stores and use them for energy. This is typ-



Reproduced by permission of WWE, Inc.

Testimonial

“When I was an Olympic wrestler, putting on a ton of muscle mass wasn’t a concern because I needed to stay in my weight class. But once I entered the WWE, I found myself going against 300-pound monsters. So, I needed to pack on serious size, but I didn’t want to put on body fat or screw up my endurance and quickness. Macrobiotic Nutrition has allowed me to pack on over 20 lbs of lean body mass.”

—**Kurt Angle**, *Olympic and Professional Wrestler*

ically the case if you are overfat and inactive. Being overfat and inactive makes your body develop a fat-building metabolism. What you want to do is reverse this condition, or better yet, prevent it from occurring in the first place. Macrobiotic Nutrition helps you make your metabolism a fat-burning and muscle-building metabolism.

There are many supplements that help improve your body’s fat-metabolizing rate. Sometimes just ingesting certain essential vitamins and minerals you are deficient in helps improve your body’s use of stored body fat for energy. For example, chromium is a trace mineral shown to help many people turn ingested nutrients into energy instead of storing them as body fat. Chromium is a cofactor for the proper function of insulin, the hormone that signals cells to let in nutrients that are circulating in bloodstream after a meal. Adequate intake of chromium allows insulin to function at full capacity. Other studies have shown that for some individuals, just ingesting calcium and magnesium supplements along with following a weight-loss diet and exercise program helps improve their rate of fat loss. There is also derived banaba leaf, a botanical ingredient with insulinlike function that I like to use to help get as much of the ingested nutrients into your cells as possible. Banaba contains a substance called corosolic acid. Scientists who have conducted clinical research on products containing corosolic acid found that it promotes improvements in lean body mass. It is believed that corosolic acid has a pseudo-insulin effect, working independently of insulin to help stimulate the cells in your body, including muscle cells, to actively take in nutrients like amino acids, essential fatty acids, and glucose. I was so impressed with the research on corosolic acid and the benefits I directly observed that I added it to the Up Your MASS products.

Fat-loss science has also turned its attention to exploring other ways to optimize your fat-loss metabolism. What has become known as the thermogenic fat-loss aids work to stimulate your body to use and metabolize body fat and ingested fats beyond what a healthy body, fortified with essential nutrients, would otherwise experience. The positive impact essential nutrient intake has on losing body fat underscores the importance of maintaining an adequate, nutrient-dense eating program like Macrobiotic Nutrition, because thermogenic fat-loss aids work even better when you are well nourished. So first and foremost, get your Macrobiotic Nutrition program perfected and stick to it. Only then should you try using thermogenic fat-loss aids. When discussing thermogenics, I contend that ephedra is hands-down number one. Regardless of all the negative press on ephedra, I personally think that when it is used properly by healthy individuals it is the most effective thermogenic fat-loss aid available. The most recent research results agree with my position. For example, the results of one of the highest quality research studies on ephedra-containing products was recently published. In 2002, C. N. Boozer and co-workers of the New York Obesity Research Center, at St. Luke's-Roosevelt Hospital and Columbia University, found that during a six-month period, healthy, obese people taking the herbal ephedrine/caffeine product lost more weight than those taking a placebo pill.

Research Findings Concerning the Safety and Effectiveness of Ephedrine/Ephedra Products

The results of research from a Harvard Medical Research team headed by Diane R. Krieger on how thermogenic aids can stimulate significant losses in body-fat mass got the attention of the medical community. Krieger reported the results of this study in an article titled "Ephedrine, Caffeine and Aspirin Promote Weight Loss in Obese Subjects," in *Transactions of the Association of American Physicians*.

It was in this research article that Krieger revealed the model of how the ECA (ephedrine, caffeine, and aspirin) combination might work in conjunction with a reduced-calorie diet to increase fat loss by stimulating the thermogenic effect. Ephedrine stimulates the production of norepinephrine release in the nervous system. This increased release of norepinephrine causes a chain reaction, which results in increasing the activity of a substance in the mitochondria called uncoupling protein that stimulates the release of energy from fatty acids. However, due to the fact that the body has to regulate overproduction of substances, the ephedrine-stimulated increase of norepinephrine production causes the release of two substances that work to reduce it: adenosine

and prostaglandins. This is where aspirin and caffeine come in. Aspirin works to inhibit the prostaglandin synthesis, and caffeine to reduce adenosine activity. The result is prolonged stimulation of increased norepinephrine release and the thermogenic fat-burning process.

Supplement companies, inspired by the results of this research, sought to create herbal EC and ECA products, with ephedra as the cornerstone ingredient to supply herbal ephedrine and guarana, kola nut, green tea, and maté being the primary source of herbal caffeine. Willow bark is an herbal source of naturally occurring aspirinlike substances called salicin.

Unfortunately, all of the controversy spearheaded by the irresponsible media regarding ephedra-based products led to public uncertainty regarding its safety. As the debate continued, I decided that we should start researching options other than ephedra (even though it continues to be my favorite thermogenic fat-loss aid). MHP product development was steered toward formulating an ephedra-free thermogenic fat-loss aid that would duplicate the thermogenic effects of ephedra. MHP was at the forefront in developing a safe and effective alternative. When the FDA banned ephedra's use in dietary supplements, MHP already had an excellent substitute available.

The Next Generation Thermogenic Fat-Loss Aids

Once again, I directed MHP's research efforts toward examining the scientific research and what traditional natural medicine had to offer as ephedra substitutes. I was impressed with the botanical candidates we identified, and narrowed it down to two of the substances that I found worked best for me, along with the time- and research-tested guarana. These new substances included synephrine from bitter orange (*Citrus aurantium*), and certain polyphenols from green tea (*Camellia sinensis*). As it turns out, synephrine works like ephedrine in that it stimulates the body's use of fats for energy and increases caloric-burning rate. But, unlike ephedrine, it does not have the potent central nervous system and heart-stimulating effects. Simply put, synephrine works in the brain and in the body to increase the use of fatty acids for energy.

Medical studies on overweight people have demonstrated that they have a slow metabolic rate and do not use much fat for energy. This citrus solution takes care of these two major weight-loss obstacles, stimulating the use of fats for energy, and increasing the number of calories the body uses each day. Synephrine selectively turns up the switches in the body that control the rate at which fat is released from body stores (lipolysis) and the metabolic rate. This results in more fat and more calories being used for energy. In a 2002 issue of *Journal of Medicine*, researchers H. G. Preuss and coworkers from the Depart-

ment of Physiology, Medicine and Pathology of Georgetown University reviewed the existing published research on *Citrus aurantium* products in their article titled “*Citrus Aurantium* as a Thermogenic, Weight Reduction Replacement for Ephedra: An Overview.” They reached a similar conclusion to mine; there is scientific evidence that *Citrus aurantium* can stimulate thermogenesis and promote a significant increase in the reduction of body-fat mass.

Research is also available regarding the green tea polyphenols. A. G. Dullo, one of the ephedra researchers previously mentioned, has taken the lead in proving that substances in green tea extract can increase energy expenditure and fat oxidation in humans. The neurochemical (brain chemical) norepinephrine (noradrenaline) is a key player in stimulating thermogenesis, energy, and appetite control. The green tea polyphenols block an enzyme (catechol O-methyltransferase) that normally degrades norepinephrine and helps maintain and prolong its thermogenic effect. MHP’s TakeOFF formula also contains caffeine from guarana to further help stimulate and maintain the thermogenic effect, since caffeine has synergistic properties. Caffeine on its own is thermogenic and promotes lipolysis; it also helps maintain the release of norepinephrine by inhibiting adenosine, another substance that breaks down norepinephrine in the neuroeffector junction.

With the ECA activity taken care of, additional synergistic ingredients were considered for addition to the TakeOFF formula. We developed a special triple ginseng concentrate—Chinese ginseng (*Panax ginseng*), American ginseng (*Panax quinquefolius*), and Siberian ginseng (*Eleutherococcus senticosus*; eleutherosides)—to include in the TakeOFF thermogenic fat-loss aid formula for a few reasons, but the main one centered around MHP’s holistic product development philosophy. We recognized that although the *Citrus aurantium*, guarana, and green-tea extracts maximize the body’s thermogenic effects, such prolonged, stimulated production of norepinephrine could lead to stress of the adrenal glands. One of the ways these ginsengs help promote health and energy in the body is by supporting adrenal gland function. It is through this activity that ginsengs are thought to exert their adaptogenic effects. The term “adaptogen” was coined by researchers to describe a substance that helps to increase the body’s resistance to adverse physical and environmental influences, that is, a cure-all. To be a true adaptogen, the substance must prove safe for daily use, increase the body’s resistance to a wide variety of harmful factors, and have a normalizing action in the body. Adaptogens are useful to healthy individuals as a method of coping with daily stresses and workload, and also as a tonic support to help the body normalize when it is ill (in addition to primary medical treatment). Ginseng is an example of one of the more popular adaptogenic herbs.

The “active” substances of ginseng are a group of sponin compounds called ginsenosides/panaxosides, and eleutherosides in Siberian ginseng. Ginseng also contains the trace mineral germanium, which has been shown to exhibit overall beneficial health effects and to increase the body’s supply of oxygen.

Recent research has also determined that taking ginseng on a regular basis helps to keep blood sugar levels within a healthy range. Keeping your blood sugar levels under control is an important factor when trying to lose body fat. On average, when daily blood sugar levels are elevated, the body can’t liberate and use its stored fat as well as it can when blood sugar levels are lower. In fact, high average blood sugar levels put you in a fat-building mode. The Macrobiotic Nutrition program helps you keep your blood sugars in the right range for maximum energy and fat loss, but some ginseng each day can help this process.

I was pleased with the initial results in the TakeOFF prototype formula at this point in product development. But I wanted to further enhance the effects and duplicate the boost in physical and mental energy I got from my ephedra-containing formulas. After some further research and development, we concluded that the addition of ginkgo and L-tyrosine does the trick, with both possessing multiple functions. Ginkgo improves mental focus, concentration, and memory. It also improves circulation, which further enhances the fat-metabolizing effects of the other ingredients. L-tyrosine is involved in nervous system function and supports the thyroid to stimulate weight loss, thermogenesis, and energy production, and to improve mental focus. TakeOFF’s synergistic ingredients work together to provide an immediate burst of energy to fuel you through an intense workout and keep you on the go through your busy day. Its scientifically balanced performance formula was designed to increase energy levels and boost metabolism—*without ephedra!* This revolutionary formula maximizes thermogenic fat burning, improves concentration and physical performance, and dramatically increases energy. TakeOFF really gets me energized and focused for my workouts, and helps increase the fat-burning effects of exercise.

Nighttime Lean Muscle-Building Requirements

An important part of the Macrobiotic Nutrition approach is nutritional consideration of your nighttime metabolic needs. In Chapter 3, I reviewed how protein intake must be staggered throughout the day to deliver a steady supply of amino acids in the blood to keep your body in an anabolic muscle-building state. Satisfying the metabolic needs during sleep can be tricky, however. As previously mentioned, this is one time we depart from the 45/35/20 rule, because it is more advantageous to consume a supplement that is high in

protein and lower in carbohydrates and fat, at bedtime. The goal is to provide a steady supply of amino acids, keep insulin levels down, and help raise growth hormone and glucagon levels. Since you are not working out in your sleep, you do not require carbohydrates for glycogen replenishment at night.

Developing a nighttime protein supplement to maximize anabolic and fat-metabolizing processes during sleep became the most recent research challenge of the MHP product-development team. In fact, it was during the writing and completion of this book that I signed off on the final formulation. This new formula is called Probolic SR. It contains the same highly effective Probolic-engineered protein found in Up Your MASS, but in a special base nutrient matrix compatible with your nighttime metabolic needs. Therefore, including Probolic SR in your Macrobiotic Nutrition program will help maximize your lean muscle-building efforts and help you experience faster results.

A Final Word about Enhancing the Macrobiotic Nutrition Effect

Macrobiotic Nutrition will provide the core of your nutritional needs. However, additional supplementation with key ingredients can help enhance your muscle-building and fat-burning efforts. The supplements described in this chapter are the ones I think are the most important to consider adding to your program. In addition to these performance-enhancing supplements, I also recommend a well-formulated multivitamin/mineral supplement. I reviewed the importance of these supplements earlier in the book. Be sure to choose a high-quality brand. I also recommend taking 2,000 to 3,000 extra milligrams of vitamin C and a good antioxidant formula to combat the free radicals produced by your intense training program.

If this chapter comes off as hype for some of my products, believe me, it's not. I am conveying my passion, commitment, and confidence in these formulas. Keep in mind that the reason I started MHP was to find ways to improve the quality of supplements available. As a bodybuilder, I relied heavily on supplements to enhance my physique. The supplements I used prior to MHP did work, but I knew they could be better.

Unlike most supplement companies, which are owned by people who never compete (or even touch a weight, for that matter), MHP supplements are backed by science from leading experts and universities, and proven by a bodybuilder—me—and many other world-class athletes. Every MHP product described in this book was inspired by my own personal quest to find a better means of supplementation, to improve my physique as well as yours. I'm not bashing all other supplements or supplement companies, because there are

a few good companies out there who make good products. Unfortunately, however, there are some poor-quality products out there that are “all hype.” It can be difficult, as a sports nutrition consumer, to determine and choose which products are best for you. Hopefully, this overview on supplements has shown you that you can’t go wrong with MHP when it comes to sports nutrition.

It is very rewarding to hear the positive feedback I get from satisfied customers. MHP receives thousands of calls and e-mails from customers, telling us about the gains and progress they have made by using MHP products. There is nothing more rewarding than receiving these testimonials. Helping others achieve maximum human performance has been and will remain my commitment to all athletes and fitness enthusiasts.



Photo by Irvin J. Gelb.

CHAPTER 10

Macrobiotic Meals

AN ADDED BENEFIT OF THE MACROBOLIC 45/35/20 ratio is that it allows you to choose from a wide variety and great selection of foods. The more flexibility and variety in a diet, the easier it is for people to follow. Knowing this, I teamed up with Roger Warn, a nutritionist from Montclair State University, to create some delicious Macrobiotic meals. These meals taste great—from my personal favorite, Mexican Omelet with Granola for breakfast, to King Crab Roll Sushi for dinner.

The sample meals I include offer a great variety. Some are quick and easy to prepare, while others take a little more time. You can try to create some of your own favorites; it's not that hard. Just read the nutrition information label on the foods you prepare and try to stay within the 45/35/20 ratio. You don't have to be exact; you will see from my sample meals that I sometimes am a little off in the ratio numbers. It is nearly impossible to prepare meals to the exact ratio, but try to stay within the range of 42 to 48 percent carbohydrates, 32 to 38 percent protein, and 17 to 24 percent fat.

The meals I have provided and the ones that you create will probably need to be adjusted to meet your calculated calorie requirements for each meal. Choose your favorite meals, prepare and enjoy!

BREAKFAST

Mexican Omelet with Macrobiotic Granola

YIELD: 1 SERVING

5 large eggs, whites only

¼ cup chopped sweet green pepper

¼ cup chopped sweet red pepper

2 tablespoons chopped onions

1 ounce shredded low-fat cheddar cheese

1 serving Macrobiotic Granola*

1 cup Skim Plus Milk or skim milk

1. Preheat an 8-by-12-inch skillet over medium-low heat.
2. Place egg whites in a bowl and beat with a fork until blended.
3. Spray preheated pan with nonstick cooking spray and pour in the egg whites. Flip eggs when the underside starts to get firm or it reaches the desired consistency (about 2–3 minutes).
4. Place the vegetables and cheese on top of one side of the omelet. When the underside reaches desired consistency, fold one half neatly over the other and allow the cheese to melt.
5. Serve with a side of Macrobiotic Granola mixed with milk.

*See recipe for Macrobiotic Granola in this chapter.

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Egg whites	5 large	83	1.70	17.36	—	—
Sweet green pepper	¼ cup	10	2.40	0.33	0.07	0.70
Sweet red pepper	¼ cup	10	2.40	0.33	0.07	0.70
Onions	2 tbs	8	1.73	0.23	0.03	—
Cheddar cheese, low fat	1 oz	49	0.54	6.90	1.98	—
Macrobiotic Granola	1 serving	218	29.34	9.42	8.10	6.22
Skim Plus Milk	1 cup	101	13.68	9.74	0.62	—
	TOTALS	479	51.79	44.31	10.89	7.62
			43%	37%	20%	

==== Egg White Omelet with Macrobiotic Granola ====

YIELD: 1 SERVING

- 5 large eggs, whites only
 2 tablespoons chopped onions
 1 serving Macrobiotic Granola*
 1½ cups Skim Plus Milk or skim milk

1. Preheat an 8-by-12-inch skillet over medium heat.
2. Place egg whites in a bowl and beat with a fork until blended.
3. Mix half of the onions into the egg whites.
4. Spray preheated pan with nonstick cooking spray and pour in the egg whites. Flip eggs when the underside starts to get firm or it reaches the desired consistency (about 2–3 minutes).
5. Place the remaining onions on top of one side of the omelet and fold one half neatly over the other.
6. Serve with a side of Macrobiotic Granola mixed with milk.

*See recipe for Macrobiotic Granola in this chapter.

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Egg whites	5 large	83	1.70	17.36	—	—
Onions	2 tbs	8	1.73	0.23	0.03	
Macrobiotic Granola	1 serving	218	29.34	9.42	8.10	6.22
Skim Plus Milk	1½ cups	151	20.52	14.61	0.92	—
TOTALS		460	52.63	41.62	9.07	6.22
			46%	36%	18%	

==== Macrobiotic Granola Mix with Side of Eggs ====

YIELD: 1 SERVING

- 4 large eggs
 ½ cup Macrobiotic Granola*
 ½ cup nonfat, plain yogurt
 1 scoop Up Your MASS CinnaBun (measuring scoop included)

1. Boil the eggs until they are hard-boiled (approximately 15 minutes after the water comes to a boil and the heat is reduced). Peel the eggs, cut them in half, and discard the egg yolks. Place the egg whites in a bowl.
2. Mix the Macrobiotic Granola with the yogurt and one scoop of Up Your MASS CinnaBun in a separate bowl. Serve.

*See recipe for Macrobiotic Granola in this chapter.

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Macrobiotic Granola	½ cup	218.0	29.34	9.42	8.10	6.22
Plain nonfat yogurt	½ cup	64.0	8.27	6.50	0.20	—
Up Your MASS CinnaBun	1 scoop	127.5	14.50	11.00	2.75	0.30
Hard-boiled egg whites	4 large	66.0	1.36	13.89	—	—
TOTALS		475.5	53.47	40.81	11.07	6.52
			45%	34%	21%	

Macrobiotic Morning Mix

YIELD: 1 SERVING

½ cup low-sodium instant oatmeal,
cooked

1¼ cup low-fat (1 percent fat)
cottage cheese

1 small apple, chopped

1 tablespoon flaxseed

1 cup Skim Plus Milk or skim milk

1. Place the oatmeal in a cereal bowl and mix in the cottage cheese, chopped apple, and flaxseed.
2. Serve the milk on the side.

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Low-sodium oatmeal	100 g	93	16.67	3.95	1.79	2.3
Low-fat cottage cheese	1¼ cup	203	7.68	35.00	2.88	—
Apple	small	63	16.16	0.20	0.38	2.9
Flaxseed	1 tbs	59	4.11	2.34	4.08	3.3
Skim Plus Milk	1 cup	101	13.68	9.74	0.62	—
TOTALS		519	58.30	51.23	9.74	8.5
			45%	38%	17%	

Macrobiotic Pancakes

YIELD: 8 SERVINGS

- 1 cup whole-wheat flour
- 4 scoops Up Your MASS CinnaBun
- ⅓ cup wheat germ
- ½ teaspoon baking soda
- 2 large whole eggs
- 7 large eggs, whites only
- 2½ cups Skim Plus Milk or skim milk
- 1 tablespoon safflower oil
- Water, as needed*

1. Mix the whole-wheat flour, Up Your MASS CinnaBun, wheat germ, and baking soda together in a bowl.
2. Mix the whole eggs and egg whites together in a separate bowl.
3. Add the Skim Plus Milk to the dry mixture and place in a blender. Set the blender on low and blend for 3 minutes.
4. Add the egg mixture to the blender mixture and blend for 2 minutes.
5. Add the safflower oil and increase the blender speed. Add water until the batter reaches the desired thickness, approximately 2 minutes.
6. Lightly coat a pan with nonstick cooking spray and heat.
7. Pour some of the mixture into the heated pan and allow to cook until a corner of the pancake lifts easily. Flip the pancake and cook other side. Repeat with the remaining mixture.

*Note: For thicker pancakes, use less water. For thinner pancakes, use more water.

Macrobiotic Pancake—per serving

(Suggested Serving Size: 2 pancakes and 1 cup of Skim Plus Milk)

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Macrobiotic Pancake	2 pancakes	389	45.80	30.40	9.60	3.64
Skim Plus Milk	1 cup	101	13.68	9.74	0.62	—
TOTALS		490	59.48	40.14	10.22	3.64
			48%	33%	19%	

Macrobiotic Pancakes—full recipe ■ 8 servings

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Whole-wheat flour	1 cup	407	87.08	16.44	2.24	14.6
Up Your MASS CinnaBun	4 scoops	510	58.00	44.00	11.00	10.5
Wheat germ	1/3 cup	1	0.17	0.08	0.03	5.0
Baking soda	1/2 tsp	—	—	—	—	—
Skim Plus Milk	2 1/2 cups	252	34.20	24.35	1.54	—
Eggs, whole	2 large	149	1.22	12.49	10.02	—
Eggs, white	7 large	116	2.38	20.30	—	—
Oil, safflower	1 tbs	120	—	—	13.60	—
TOTALS		1,555	183.05	121.66	38.43	30.1
<i>per pancake</i>		<i>194.39</i>	<i>22.90</i>	<i>15.20</i>	<i>4.80</i>	<i>3.76</i>
			47%	31%	22%	

Macrobiotic Granola

YIELD: 24 HALF-CUP SERVINGS

7 cups quick oats

1 1/2 cups wheat germ, crude

1 1/2 cups wheat bran, crude

6 scoops Up Your MASS CinnaBun

1/4 cup safflower seed oil

1/2 cup water

1 tablespoon vanilla extract
 ¼ cup honey
 1 teaspoon ground cinnamon
 1 teaspoon ground nutmeg
 ½ cup chopped pecans
 ½ cup sliced or chopped almonds

1. Preheat oven to 275°F.
2. In a large bowl, mix the oats, wheat germ, wheat bran, and Up Your MASS CinnaBun.
3. In a separate bowl, mix together the remaining ingredients except the pecans and almonds. Combine with the oat mixture and place into a shallow baking dish.
4. Bake for 30 minutes, stirring every 15 minutes. After 30 minutes, stir in the pecans and almonds, and continue baking for 15 minutes more, or until the mixture is lightly brown. Remove and cool and store in an airtight container.

**Note: Macrobiotic Granola is nutritionally enhanced, but does not meet the 45/35/20 ratio. Macrobiotic Granola should be used with other high-protein food sources as part of a meal to balance the ratios to 45/35/20. (See breakfast omelet with Macrobiotic Granola recipes.)*

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Quick oats	7 cups	2,177	379.89	9.77	35.72	60.1
Wheat germ, crude	1½ cups	621	89.36	39.93	16.77	22.8
Wheat bran, crude	1½ cups	188	56.12	13.53	3.70	37.2
Up Your MASS CinnaBun	6 scoops	765	87.00	66.00	16.50	18.0
Safflower seed oil	¼ cup	482	—	—	54.50	—
Water	½ cup	—	—	—	—	—
Vanilla extract	1 tbs	37	1.64	0.01	0.01	—
Honey	¼ cup	258	69.83	0.25	—	—
Cinnamon	1 tsp	6	—	0.09	0.07	—

Nutmeg	1 tsp	12	1.08	0.13	0.80	—
Pecans	½ cup	411	8.25	5.46	42.82	5.7
Almonds	½ cup	275	9.38	10.10	24.05	5.6
TOTALS		5,232	704.39	226.27	194.90	149.4
<i>per ½ cup serving</i>		<i>218</i>	<i>29.34</i>	<i>9.42</i>	<i>8.10</i>	<i>6.2</i>
			51%	17%	32%	

Scrambled Eggs with Oatmeal and Strawberries

YIELD: 1 SERVING

½ cup quick oats

4–8 ounces water

6 large eggs, whites only

1 large whole egg

1 cup fresh strawberry slices

1. Place the oats in a microwavable container. Add water and place in the microwave. Cook for about 1 minute or until the oatmeal reaches desired consistency.
2. Preheat a skillet over medium heat.
3. In a bowl, beat the whole eggs and egg whites with a fork until well blended.
4. Coat the skillet with nonstick cooking spray and pour in the eggs. Gently mix the eggs with a wooden spoon until cooked.
5. Serve the eggs alongside the bowl of oatmeal and the strawberry slices.

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Quick oats	½ cup	156	27.00	6.00	2.50	4.3
Egg whites	6 large	99	2.04	20.38	—	—
Whole egg	1 large	75	—	6.25	5.00	—
Strawberries	1 cup	50	11.65	1.01	0.61	3.8
TOTALS		380	40.69	33.64	8.11	8.1
			44%	36%	20%	

LUNCH

Chicken Parm Pasta Salad

YIELD: 1 SERVING

- 2 ounces melted shredded mozzarella
- 1 cup cooked whole-wheat spaghetti
- 2 large ripe tomatoes, sliced
- 2 large raw mushrooms, sliced
- 1 teaspoon dried parsley
- $\frac{1}{2}$ teaspoon minced garlic
- 2 ounces sliced roasted chicken breast, meat only, skin and visible fat removed

1. Mix the mozzarella cheese into the cooked spaghetti.
2. Mix the sliced vegetables into the spaghetti and add the parsley and garlic.
3. Place the roasted chicken breast slices over the spaghetti.
4. Serve either hot or chilled.

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Mozzarella	2 oz	144	1.58	13.76	9.02	—
Whole-wheat spaghetti	1 cup	174	37.16	7.46	0.76	6.3
Tomatoes	2 large	76	16.89	3.09	1.20	4.0
Mushrooms	2 large	12	1.88	1.33	0.15	—
Parsley	1 tsp	1	0.15	0.07	—	—
Garlic	$\frac{1}{2}$ tsp	1	0.23	0.04	—	—
Roast chicken breast, meat only	2 oz	94	—	17.59	2.02	—
	TOTALS	502	57.89	43.34	13.15	10.3
			44%	33%	23%	

Grilled Chicken–Bean Salad

YIELD: 1 SERVING

$\frac{1}{2}$ cup chopped grilled chicken breast,
meat only, skin and visible fat removed

$\frac{1}{2}$ cup canned black beans

$\frac{1}{2}$ cup canned kidney beans

$\frac{1}{2}$ cup chopped sweet green peppers

$\frac{1}{2}$ cup chopped sweet red peppers

1 teaspoon olive oil

$\frac{1}{8}$ teaspoon dried ground basil

$\frac{1}{8}$ teaspoon dried ground oregano

1 tablespoon grated Parmesan cheese

1. Place chicken, beans, and peppers in a bowl, and toss. Mix in the olive oil, basil, and oregano.
2. Sprinkle with the cheese.
3. Serve either hot or chilled.

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Grilled chicken breast, meat only	$\frac{1}{2}$ cup	116	—	21.71	2.50	—
Black beans	$\frac{1}{2}$ cup	114	20.39	7.62	0.49	7.5
Kidney beans	$\frac{1}{2}$ cup	112	20.19	7.67	0.44	5.7
Sweet green pepper	$\frac{1}{2}$ cup	20	4.79	0.66	0.14	1.3
Sweet red pepper	$\frac{1}{2}$ cup	20	4.79	0.66	0.14	1.5
Olive oil	1 tsp	40	—	—	4.50	—
Basil	$\frac{1}{8}$ tsp	—	—	—	—	—
Oregano	$\frac{1}{8}$ tsp	1	0.12	0.02	0.02	—
Parmesan cheese	1 tbs	23	0.19	2.08	1.50	—
	TOTALS	446	50.47	40.42	9.73	16.0
			45%	36%	19%	

Tuna Salad Sandwich

YIELD: 1 SERVING

- 1 6-ounce can white tuna in water, drained
- 1/2 cup chopped carrots
- 1/2 cup sliced cucumber
- 1 teaspoon flaxseed
- 1/2 cup sliced celery
- 1/2 cup chopped onion
- 2 tablespoons lemon juice
- 1 cup shredded loose-leaf lettuce
- 3 tablespoons fat-free mayonnaise
- 2 slices pumpernickel bread

1. In a large bowl, mix together all of ingredients except the bread.
2. Place the salad between the slices of bread and serve.

Note: If there is any extra, place on a side dish and serve with the sandwich.

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Tuna, white, canned in water	6 oz	220	—	40.63	5.11	—
Carrots	1/2 cup	28	6.49	0.66	0.12	1.9
Cucumber	1/2 cup	7	1.44	0.36	0.07	0.4
Flaxseed	1 tsp	19	1.36	0.77	1.35	1.1
Celery	1/2 cup	10	2.19	0.45	0.08	1.1
Onions	1/2 cup	30	6.90	0.93	0.13	1.4
Lemon juice	2 tbs	6	1.94	0.12	0.09	—
Loose-leaf lettuce	1 cup	10	1.96	0.73	0.17	—
Fat-free mayo	3 tbs	34	5.95	0.10	1.30	2.0
Pumpernickel bread	2 slices	160	30.28	5.50	1.98	4.2
TOTALS		524	58.51	50.25	10.40	12.1
			44%	38%	18%	

Beef Mass Burger

YIELD: 2 SERVINGS

8 ounces lean ground beef or turkey
 ¼ cup shredded low-fat cheddar cheese
 ⅛ teaspoon parsley
 ⅛ teaspoon basil
 2 rolls whole-wheat hamburger rolls
 2 leaves loose-leaf lettuce
 4 medium slices ripe tomato

1. Preheat a skillet or grill to desired heat range (145°F for rare; 160°F for medium; 170°F for well-done).
2. Mix the beef or turkey with the cheese and spices in a large bowl. Form into two equal patties.
3. Coat skillet or griddle with nonstick cooking spray and cook patties until desired doneness.
4. Place on whole-wheat roll with lettuce and tomato. Serve.

Per 2 servings

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Lean ground beef	8 oz	311	—	48.60	11.35	—
Low-fat cheddar cheese	¼ cup	49	0.54	6.88	1.98	—
Whole-wheat hamburger rolls	2 rolls	532	102.20	17.40	4.04	15.08
Parsley	⅛ tsp	—	—	0.01	—	—
Basil	⅛ tsp	—	0.11	0.03	0.01	—
Loose-leaf lettuce	2 leaves	4	0.70	0.26	0.06	—
Tomato	4 slices	14	3.71	0.68	0.26	0.90
TOTALS		910	107.26	73.86	17.70	15.98
<i>per serving</i>		<i>455</i>	<i>53.63</i>	<i>36.93</i>	<i>8.85</i>	<i>7.99</i>
			49%	33%	18%	

Steak Pita Pocket

YIELD: 1 SERVING

- 4 ounces beef chuck cubes
- 1/8 teaspoon salt
- 1/8 teaspoon ground cumin
- 1/4 clove garlic, diced
- 1 cup chopped tomatoes
- 1 ounce shredded low-fat cheddar cheese
- 1/2 cup shredded loose-leaf lettuce
- 5 thin slices onion
- 3 jalapeño peppers, chopped
- 1 cup sliced sweet red pepper
- 1/8 cup prepared (commercial) salsa
- 1 large whole-wheat pita

1. Preheat a skillet over high heat.
2. Slice each of the beef chuck cubes in half.
3. Coat the skillet with nonstick cooking spray and immediately add the beef chuck halves.
4. Stir-fry for about 3–5 minutes, or until the beef chuck halves brown on all sides.
5. While the beef chuck is cooking, mix the salt with the cumin and place in a bowl with the remaining ingredients except for the pita. Mix together if desired.
6. When the beef chuck is fully cooked, place it on a dish or a cutting board. Slice the beef chuck halves into smaller slices. Then place in the bowl with the other ingredients and stir until well mixed.
7. Cut a pocket into the whole-wheat pita. Place the shredded beef mixture into the pita pocket with a large spoon. Serve.

Note: If there is any extra, place on a side dish and serve with the sandwich.

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Beef chuck	4 oz	181	—	29.32	6.26	—
Salt	1/8 tsp	—	—	—	—	—
Garlic	1/4 clove	1	0.25	0.05	—	—
Cumin	1/8 tsp	1	0.12	0.05	0.06	—
Tomato	1 cup	38	8.35	1.53	0.59	2.0
Low-fat cheddar cheese	1 oz	49	0.54	6.90	1.98	—
Loose-leaf lettuce	1/2 cup	5	0.98	0.36	0.08	—
Onions	5 slices	17	3.88	0.52	0.07	0.8
Jalapeño peppers	3	13	2.48	0.57	0.26	—
Sweet red pepper	1 cup	25	5.92	0.82	0.17	3.0
Salsa	1/8 cup	9	2.02	0.41	0.08	—
Whole-wheat pita	1 large	170	35.20	6.27	1.66	4.7
TOTALS		509	59.74	46.80	11.21	10.5
			45%	36%	19%	



DINNER

Mixed-Meat Teriyaki

YIELD: 4 SERVINGS

- 1 roast or grilled, sliced chicken breast, meat only, skin and visible fat removed
- 8 ounces grilled, sliced, lean top-round or short-loin porterhouse (1/4" fat)
- 2 cups cooked medium brown rice
- 1 1/4 cup teriyaki sauce
- 1 large onion, chopped
- 1 large sweet green pepper, chopped
- 1 small zucchini, sliced
- 1 large sweet yellow pepper, chopped
- 2 tablespoons olive oil
- 1 1/2 tablespoons soy sauce
- 2 teaspoons garlic powder
- 1 teaspoon ground black pepper

1. Place the chicken, beef, and cooked rice into a bowl with the teriyaki sauce.
2. Mix the vegetables with the olive oil, soy sauce, and garlic powder. Place mixture into the bowl with the chicken, beef, and rice. Add the black pepper.
3. Divide into four equal servings. Refrigerate the leftover servings in separate containers for the rest of the week.

Per 4 servings

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Roast chicken breast, meat only	1 breast	284	—	53.35	6.14	—
Top round, lean only, 1/4" fat	8 oz	272	—	51.96	5.67	—
Medium brown rice	2 cups	437	91.96	9.05	3.24	7.0

Teriyaki sauce	1¼ cup	302	57.42	21.53	—	—
Onion	1 large	57	12.59	1.74	0.24	2.7
Sweet green pepper	1 large	44	10.55	1.46	0.31	3.0
Zucchini	1 small	17	3.42	1.37	0.17	1.4
Sweet yellow pepper	1 large	50	11.76	1.86	0.39	1.7
Olive oil	2 tbs	239	—	—	27.00	—
Soy sauce, low-sodium	1½ tbs	14	2.30	1.40	—	—
Garlic powder	2 tsp	9	2.04	0.47	0.02	—
Black pepper	1 tsp	5	1.36	0.23	0.07	—
	TOTALS	1,730	193.40	144.42	43.25	15.8
	<i>per serving</i>	<i>433</i>	<i>48.35</i>	<i>36.10</i>	<i>10.81</i>	<i>4.0</i>
			45%	33%	22%	

Lemon Sole with Broccoli

YIELD: 4 SERVINGS

- 1 pound sole or flounder fillets (4-ounces each)
- 2 tablespoons lemon juice
- Black pepper to taste
- 10 ounces broccoli spears
- 1 cup whole-wheat pasta
- 4 ounces melted low-fat cheddar cheese
- ¼ cup skim milk
- ½ teaspoon grated lemon peel
- ¼ teaspoon fresh dill
- 1½ teaspoons olive oil
- 4 slices toasted pumpernickel bread

1. Fill a large pot three-quarters with water and bring the water to a boil.
2. Sprinkle each fillet with juice and pepper. Place the broccoli spears on the narrow end of each fish fillet. Roll up the fillet starting at the end with the broccoli, so the broccoli ends up in the center of the roll. Secure the roll with toothpicks. Place the roll seam side down on a microwavable dish and cover.

3. Microwave on high for 5–7 minutes until the fish flakes easily with a fork.
4. Place whole-wheat pasta in the boiling water. (Note: individual pasta manufacturers may indicate the length of time the pasta should be cooked.)
5. Place the remaining ingredients (except the bread and the olive oil) in a microwavable bowl and mix together until blended. Microwave 2–3 minutes or until the sauce reaches a smooth consistency. (Stir the sauce after each minute to ensure even melting.)
6. Drain the water from the pasta when done and immediately mix the olive oil into the pasta.
7. Remove fillets and discard the toothpicks. Place on top of pasta (1 cup per fillet) and serve with sauce and bread.

Per 4 servings (without pasta and bread)

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Sole fillets	1 lb	530	—	109.53	6.94	—
Lemon juice	2 tbs	6	1.94	0.12	0.90	—
Broccoli spears	10 oz	70	13.38	7.75	0.28	7.5
Low-fat cheddar cheese	4 oz	196	2.17	27.61	7.94	—
Skim milk	¼ cup	23	3.07	2.19	0.15	—
Lemon peel	½ tsp	—	0.16	0.01	—	—
Fresh dill	¼ tsp	—	—	—	—	—
TOTALS		825	20.72	147.21	16.21	7.5

Per serving (with pasta and bread)

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Lemon Broccoli of Sole	1 serving	206.25	5.18	36.80	4.10	1.87
Whole-wheat pasta	1 cup	174.00	37.16	7.46	0.76	6.30
Olive oil	1½ tsp	60.00	—	—	6.75	—
Pumpernickel bread, toasted	1 slice	80.00	15.14	2.75	0.99	2.10
TOTALS		520.25	57.48	47.01	12.62	10.27
			44%	35%	21%	

Lemon Salmon with Spinach

YIELD: 1 SERVING

5 ounces farm-fresh salmon

3 tablespoons lemon juice

2 cups raw spinach, rinsed and dried

1 medium potato

1. Preheat a large skillet over medium-high heat.
2. Coat the skillet with nonfat cooking spray, and add the salmon.
3. Cook for 10 minutes for each inch of thickness.
4. Flip the fillet halfway through cooking to brown both sides.
5. When the fillet is brown on both sides, place on a dish and sprinkle the lemon juice over salmon.
6. Place the spinach on the dish with the lemon salmon, and serve with the baked potato (recipe follows).

Baked Potato

1. Preheat oven to 350°F.
2. Wash potato thoroughly with water and dry.
3. Prick the skin of the potato with a fork several times to prevent steam buildup during cooking.
4. Bake the potato for approximately 90 minutes, or until it is slightly soft (fork will pierce the potato easily) and golden brown.

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Salmon	5 oz	201	—	28.11	8.89	—
Lemon juice	3 tbs	13	3.89	0.24	0.17	—
Spinach	2 cups	14	2.10	1.72	0.22	1.6
Baked potato with skin	1 medium	145	33.63	3.06	0.16	3.8
TOTALS		380	40.67	33.99	9.55	4.6
			43%	36%	21%	

Macrobiotic Sushi Menu

YIELD: 2 ROLLS

Yellowfin Tuna Roll

- $\frac{1}{3}$ cup uncooked white short-grain rice
- 1 $\frac{1}{2}$ teaspoons cider vinegar
- $\frac{1}{4}$ medium cucumber, peeled and sliced into strips
- $\frac{1}{3}$ cup avocado, sliced into strips
- 2 $\frac{1}{2}$ sheets nori seaweed
- 6 ounces yellowfin tuna, sliced into strips

King Crab Roll

- $\frac{1}{3}$ cup uncooked white short-grain rice
- 1 $\frac{1}{2}$ tablespoons cider vinegar
- $\frac{1}{2}$ medium cucumber, peeled and sliced into strips
- $\frac{1}{3}$ cup avocado, sliced into strips
- 2 $\frac{1}{2}$ sheets nori seaweed
- 8 ounces Alaskan king crab strips

Shrimp Roll

- $\frac{1}{3}$ cup uncooked white short-grain rice
- 1 $\frac{1}{2}$ tablespoons cider vinegar
- $\frac{1}{4}$ medium cucumber, peeled and sliced into strips
- $\frac{1}{3}$ cup sliced avocado, sliced into strips
- 2 sheets nori seaweed
- 7 ounces shrimp

Rice

1. Wash the rice repeatedly under running water to remove all of the starch and until the water is clear.
2. As a rule of thumb, use twice the amount of water as rice for cooking the rice (for example, 2 cups water to 1 cup rice).
3. Place the measured water into a pot and bring to a boil.
4. Add the washed rice and return the water to a boil.
5. Add the vinegar to the rice.

6. Once the water has returned to a boil, lower the heat until it is barely simmering. Stir the rice occasionally (this makes the rice sticky, which is essential for sushi).
7. Let the pot simmer until all the water is absorbed, then take pot off heat and let it cool for an additional 10 minutes.

Sushi

1. Preheat the oven to 300°F. Heat the nori sheets on a medium baking sheet in the oven for 1 to 2 minutes, or until the sheets are warm.
2. Place the warm nori sheets on a clean dish towel. This will be used to roll the sushi.
3. Wet your hands. Then spread and press a thin layer of rice on the nori sheet.
4. Arrange and place other ingredients so that a piece of each is placed together in a single straight line, pressing gently as you go.
5. Using the dish towel, gently roll the nori sheet and press it over the ingredients.
6. Roll it forward until you make a complete roll. Repeat with the other prepared sheets.
7. Unravel the dish towel, to be reused. Cut each roll to your desired size using a wet, sharp knife.

Yellowfin Tuna Roll

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
White, short-grain rice	1/3 cup	215	47.49	3.90	0.31	1.8
Cider vinegar	1 1/2 tbs	3	0.07	—	—	—
Cucumber, peeled	1/4 medium	6	1.26	0.29	0.08	0.4
Avocado	1/3 cup	71	3.24	0.87	6.71	2.4
Nori seaweed sushi sheets	2 1/2 sheets	25	5.00	2.50	—	2.5
Tuna, yellowfin, raw	6 oz	184	—	39.75	1.67	—
TOTALS		504	57.06	47.31	8.77	7.1
			46%	38%	16%	

King Crab Roll

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
White, short-grain rice	1/3 cup	215	47.49	3.90	0.31	1.8
Cider vinegar	1 1/2 tbs	3	0.07	—	—	—
Cucumber, peeled	1/4 medium	6	1.26	0.29	0.08	0.4
Avocado, raw	1/3 cup	71	3.24	0.87	6.71	2.4
Nori seaweed sushi sheets	2.5 sheets	25	5.00	2.50	—	2.5
Alaskan king crab, raw	8 oz	190	—	41.46	1.36	—
	TOTALS	510	57.06	49.02	8.46	7.1
			46%	39%	15%	

Shrimp Roll

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
White, short-grain rice	1/3 cup	215	47.49	3.90	0.31	1.8
Cider vinegar	1 1/2 tbs	3	0.07	—	—	—
Cucumber, peeled	1/4 medium	6	1.26	0.29	0.08	0.4
Avocado, raw	1/3 cup	71	3.24	0.87	6.71	2.4
Nori seaweed sushi sheets	2.5 sheets	25	5.00	2.50	—	2.5
Shrimp, raw	7 oz	210	1.80	40.28	3.43	—
	TOTALS	530	58.86	47.84	10.53	7.1
			45%	37%	18%	

SHAKES, BARS, AND DESSERTS

Up Your MASS Shake

YIELD: 1 SERVING

4 scoops Up Your MASS
(CinnaBun or Chocolate Fudge Brownie)

15–18 ounces water

- Mix 4 scoops of Up Your MASS with 15–18 ounces of water.

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Up Your MASS Shake	4 scoops	510	58	44	11	12
Water	15–18 oz	—	—	—	—	—
TOTALS		510	58	44	11	12
			45%	35%	20%	

Up Your MASS Bar

1 BAR

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Up Your MASS bar (crunchy peanut or fudge)	1 bar	380	44	30	11	1
TOTALS		380	44	30	11	1
			46%	32%	22%	

Fruit Salad

YIELD: 1 SERVING

1/2 cup diced mandarin orange sections

1/2 cup diced strawberries

1/2 cup grapes

1/2 cup chopped apples with skin

1/2 cup peaches

1 1/2 cups low-fat cottage cheese
(not packed)

2 tablespoons flaxseed

1. Combine all of the fruit in a dish.
2. Mix in cottage cheese and sprinkle with flaxseed.

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Mandarin orange	1/2 cup	43	10.91	0.61	0.19	2.2
Strawberries	1/2 cup	25	5.83	0.51	0.31	1.9
Grapes	1/2 cup	31	7.89	0.29	0.16	0.5
Apples, raw, with skin	1/2 cup	37	9.53	0.12	0.22	1.7
Peaches	1/2 cup	37	9.43	0.59	0.08	1.7
Low-fat cottage cheese	1 1/2 cup	244	9.22	42.00	3.46	—
Flaxseed	2 tbs	118	8.22	4.68	8.16	6.7
TOTALS		535	61.03	48.80	12.58	14.7
			44%	35%	21%	

=====**Macrobiotic Instant Chocolate Pudding**=====

YIELD: 3 SERVINGS

4½ cups cold skim milk

2 packets Jell-O Brand
Instant Chocolate Pudding

(or other instant chocolate pudding)

4 scoops Up Your MASS: Fudge Brownie

1. In a bowl, combine milk and pudding mix.
2. Add 4 scoops of Up Your MASS: Fudge Brownie.
3. Using a handheld blender, begin blending mix at a low setting.
4. As the pudding gets thicker, increase the blender speed and blend for about 2 to 3 minutes, or until the pudding mix reaches a smooth consistency.
5. Cover the bowl with plastic wrap and refrigerate until firm.

Note: This dessert can also be prepared with vanilla pudding instead of chocolate, or Up Your MASS CinnaBun instead of Fudge Brownie.

Per 3 servings

Food Item	Quantity	Energy (kcal)	Carbs (g)	Protein (g)	Fat (g)	Fiber (g)
Jell-O Brand Instant Chocolate Pudding	2 packets	70	8	—	—	—
Up Your MASS: Fudge Brownie	4 scoops	510	58	46	11.0	12
Skim Milk	4½ cups	531	63	45	13.5	—
	TOTALS	1,111	129	91	24.5	12
	<i>per serving</i>	<i>370</i>	<i>43</i>	<i>30</i>	<i>8.2</i>	<i>4</i>
			45%	33%	22%	



Photo by Per Bernal.

CONCLUSION

Progress Is the Great Motivator

AS I MENTIONED EARLY IN THIS BOOK, seeing people just like you working hard in the gym and frustrated because of their lack of progress is what inspired me to write about Macrobiotic Nutrition. Like anything in life, when you feel like you are rowing against the tide, it is easy to lose interest, throw in the oars, and quit. Yet, when you are rowing with the tide and you experience progress and results, it motivates you to row even harder and faster to go further! Continual progress and motivation are residual benefits of Macrobiotic Nutrition.

What I describe as “Macrobiotic Momentum” is the net effect of Macrobiotic Nutrition’s 45/35/20 lean-mass equation. To sum it up, Macrobiotic Nutrition creates the ideal hormonal and anabolic environment to build muscle and burn body fat. Over time, as your body composition shifts to increased muscle mass and less body fat, your body’s metabolism becomes even more efficient at building muscle and burning body fat. So, you will continue to get bigger and leaner as you follow Macrobiotic Nutrition. These continual gains will keep you motivated not only to eat properly, but also to push your body to its limits when you train.

Macrobiotic Nutrition has taught you the powerful effects food has on your body. And now you understand how Macrobiotic Nutrition’s 45/35/20 lean-mass equation will optimize key hormones and supply the nutrients you need and is customized so you can reach your goals and achieve your full genetic potential.

The days of frustration and lack of progress are over. From this day forward, you can continue to make gains in lean body mass. I am confident that, if you follow the Macrobiotic Nutrition program, you will break through plateaus and reach levels of size and performance you never imagined possible.

APPENDIX A

Macrobolic Caloric Requirements Tables

In Chapter 7, you learned how to use the Macrobolic caloric equation:

$$\begin{aligned} &(\text{Bodyweight} \times \text{MGV}^*) + (\text{Bodyweight} \times \text{Lifestyle}) \\ &+ (\text{Exercise Expenditure}) = \text{Total Caloric Requirements} \end{aligned}$$

**Macrobolic Goal Variables*

If you're pressed for time or if you simply don't want to do the calculations yourself, this appendix provides you with the exact amount of calories, carbohydrates, protein, and fat you need to maximize muscle mass and target your goals based on your current body weight, goal, lifestyle, and workout schedules. First, determine your goal:

Gain, Maintain, Lose

- **Gain Body Fat/Increase Muscle**
- **Maintain Body Fat/Increase Muscle**
- **Lose Body Fat/Increase Muscle**

Then, simply locate the table that best matches your profile. For example, if your workout is 90 minutes, you weigh 215 pounds, your body fat percentage is over 15%, and you lead an active lifestyle, you'll find your target nutrient and caloric intake for workout days in the table on page 178 and for non-workout days in the table on page 179.

GOAL: LOSE BODY FAT/INCREASE MUSCLE (BODY FAT OVER 15%)

60 Minute Workouts														
BW	Sed	Carbs	Protein	Fat	BW	Mod	Carbs	Protein	Fat	BW	Act	Carbs	Protein	Fat
135	1770	199	155	39	135	2030	228	178	45	135	2680	302	235	60
140	1815	204	159	40	140	2085	235	182	46	140	2760	311	242	61
145	1860	209	163	41	145	2140	241	187	48	145	2840	320	249	63
150	1905	214	167	42	150	2195	247	192	49	150	2920	329	256	65
155	1950	219	171	43	155	2250	253	197	50	155	3000	338	263	67
160	1995	224	175	44	160	2305	259	202	51	160	3080	347	270	68
165	2040	230	179	45	165	2360	266	207	52	165	3160	356	277	70
170	2085	235	182	46	170	2415	272	211	54	170	3240	365	284	72
175	2130	240	186	47	175	2470	278	216	55	175	3320	374	291	74
180	2175	245	190	48	180	2525	284	221	56	180	3400	383	298	76
185	2220	250	194	49	185	2580	290	226	57	185	3480	392	305	77
190	2265	255	198	50	190	2635	296	231	59	190	3560	401	312	79
195	2310	260	202	51	195	2690	303	235	60	195	3640	410	319	81
200	2355	265	206	52	200	2745	309	240	61	200	3720	419	326	83
205	2400	270	210	53	205	2800	315	245	62	205	3800	428	333	84
210	2445	275	214	54	210	2855	321	250	63	210	3880	437	340	86
215	2490	280	218	55	215	2910	327	255	65	215	3960	446	347	88
220	2535	285	222	56	220	2965	334	259	66	220	4040	455	354	90
225	2580	290	226	57	225	3020	340	264	67	225	4120	464	361	92
230	2625	295	230	58	230	3075	346	269	68	230	4200	473	368	93
235	2670	300	234	59	235	3130	352	274	70	235	4280	482	375	95
240	2715	305	238	60	240	3185	358	279	71	240	4360	491	382	97
245	2760	311	242	61	245	3240	365	284	72	245	4440	500	389	99
250	2805	316	245	62	250	3295	371	288	73	250	4520	509	396	100
255	2850	321	249	63	255	3350	377	293	74	255	4600	518	403	102
260	2895	326	253	64	260	3405	383	298	76	260	4680	527	410	104
265	2940	331	257	65	265	3460	389	303	77	265	4760	536	417	106
270	2985	336	261	66	270	3515	395	308	78	270	4840	545	424	108

BW = Body Weight • SED = Sedentary • MOD = Moderately Active • ACT = Active

GOAL: LOSE BODY FAT/INCREASE MUSCLE (BODY FAT OVER 15%)**90 Minute Workouts**

BW	Sed	Carbs	Protein	Fat	BW	Mod	Carbs	Protein	Fat	BW	Act	Carbs	Protein	Fat
135	2070	233	181	46	135	2330	262	204	52	135	2980	335	261	66
140	2115	238	185	47	140	2385	268	209	53	140	3060	344	268	68
145	2160	243	189	48	145	2440	275	214	54	145	3140	353	275	70
150	2205	248	193	49	150	2495	281	218	55	150	3220	362	282	72
155	2250	253	197	50	155	2550	287	223	57	155	3300	371	289	73
160	2295	258	201	51	160	2605	293	228	58	160	3380	380	296	75
165	2340	263	205	52	165	2660	299	233	59	165	3460	389	303	77
170	2385	268	209	53	170	2715	305	238	60	170	3540	398	310	79
175	2430	273	213	54	175	2770	312	242	62	175	3620	407	317	80
180	2475	278	217	55	180	2825	318	247	63	180	3700	416	324	82
185	2520	284	221	56	185	2880	324	252	64	185	3780	425	331	84
190	2565	289	224	57	190	2935	330	257	65	190	3860	434	338	86
195	2610	294	228	58	195	2990	336	262	66	195	3940	443	345	88
200	2655	299	232	59	200	3045	343	266	68	200	4020	452	352	89
205	2700	304	236	60	205	3100	349	271	69	205	4100	461	359	91
210	2745	309	240	61	210	3155	355	276	70	210	4180	470	366	93
215	2790	314	244	62	215	3210	361	281	71	215	4260	479	373	95
220	2835	319	248	63	220	3265	367	286	73	220	4340	488	380	96
225	2880	324	252	64	225	3320	374	291	74	225	4420	497	387	98
230	2925	329	256	65	230	3375	380	295	75	230	4500	506	394	100
235	2970	334	260	66	235	3430	386	300	76	235	4580	515	401	102
240	3015	339	264	67	240	3485	392	305	77	240	4660	524	408	104
245	3060	344	268	68	245	3540	398	310	79	245	4740	533	415	105
250	3105	349	272	69	250	3595	404	315	80	250	4820	542	422	107
255	3150	354	276	70	255	3650	411	319	81	255	4900	551	429	109
260	3195	359	280	71	260	3705	417	324	82	260	4980	560	436	111
265	3240	365	284	72	265	3760	423	329	84	265	5060	569	443	112
270	3285	370	287	73	270	3815	429	334	85	270	5140	578	450	114

GOAL: LOSE BODY FAT/INCREASE MUSCLE (BODY FAT OVER 15%)

Non-Workout														
BW	Sed	Carbs	Protein	Fat	BW	Mod	Carbs	Protein	Fat	BW	Act	Carbs	Protein	Fat
135	1170	132	102	26	135	1430	161	125	32	135	2080	234	182	46
140	1215	137	106	27	140	1485	167	130	33	140	2160	243	189	48
145	1260	142	110	28	145	1540	173	135	34	145	2240	252	196	50
150	1305	147	114	29	150	1595	179	140	35	150	2320	261	203	52
155	1350	152	118	30	155	1650	186	144	37	155	2400	270	210	53
160	1395	157	122	31	160	1705	192	149	38	160	2480	279	217	55
165	1440	162	126	32	165	1760	198	154	39	165	2560	288	224	57
170	1485	167	130	33	170	1815	204	159	40	170	2640	297	231	59
175	1530	172	134	34	175	1870	210	164	42	175	2720	306	238	60
180	1575	177	138	35	180	1925	217	168	43	180	2800	315	245	62
185	1620	182	142	36	185	1980	223	173	44	185	2880	324	252	64
190	1665	187	146	37	190	2035	229	178	45	190	2960	333	259	66
195	1710	192	150	38	195	2090	235	183	46	195	3040	342	266	68
200	1755	197	154	39	200	2145	241	188	48	200	3120	351	273	69
205	1800	203	158	40	205	2200	248	193	49	205	3200	360	280	71
210	1845	208	161	41	210	2255	254	197	50	210	3280	369	287	73
215	1890	213	165	42	215	2310	260	202	51	215	3360	378	294	75
220	1935	218	169	43	220	2365	266	207	53	220	3440	387	301	76
225	1980	223	173	44	225	2420	272	212	54	225	3520	396	308	78
230	2025	228	177	45	230	2475	278	217	55	230	3600	405	315	80
235	2070	233	181	46	235	2530	285	221	56	235	3680	414	322	82
240	2115	238	185	47	240	2585	291	226	57	240	3760	423	329	84
245	2160	243	189	48	245	2640	297	231	59	245	3840	432	336	85
250	2205	248	193	49	250	2695	303	236	60	250	3920	441	343	87
255	2250	253	197	50	255	2750	309	241	61	255	4000	450	350	89
260	2295	258	201	51	260	2805	316	245	62	260	4080	459	357	91
265	2340	263	205	52	265	2860	322	250	64	265	4160	468	364	92
270	2385	268	209	53	270	2915	328	255	65	270	4240	477	371	94

BW = Body Weight • SED = Sedentary • MOD = Moderately Active • ACT = Active

GOAL: LOSE BODY FAT/INCREASE MUSCLE (BODY FAT UNDER 15%)**60 Minute Workouts**

BW	Sed	Carbs	Protein	Fat	BW	Mod	Carbs	Protein	Fat	BW	Act	Carbs	Protein	Fat
130	1900	214	166	42	130	2160	243	189	48	130	2810	316	246	62
135	1950	219	171	43	135	2220	250	194	49	135	2895	326	253	64
140	2000	225	175	44	140	2280	257	200	51	140	2980	335	261	66
145	2050	231	179	46	145	2340	263	205	52	145	3065	345	268	68
150	2100	236	184	47	150	2400	270	210	53	150	3150	354	276	70
155	2150	242	188	48	155	2460	277	215	55	155	3235	364	283	72
160	2200	248	193	49	160	2520	284	221	56	160	3320	374	291	74
165	2250	253	197	50	165	2580	290	226	57	165	3405	383	298	76
170	2300	259	201	51	170	2640	297	231	59	170	3490	393	305	78
175	2350	264	206	52	175	2700	304	236	60	175	3575	402	313	79
180	2400	270	210	53	180	2760	311	242	61	180	3660	412	320	81
185	2450	276	214	54	185	2820	317	247	63	185	3745	421	328	83
190	2500	281	219	56	190	2880	324	252	64	190	3830	431	335	85
195	2550	287	223	57	195	2940	331	257	65	195	3915	440	343	87
200	2600	293	228	58	200	3000	338	263	67	200	4000	450	350	89
205	2650	298	232	59	205	3060	344	268	68	205	4085	460	357	91
210	2700	304	236	60	210	3120	351	273	69	210	4170	469	365	93
215	2750	309	241	61	215	3180	358	278	71	215	4255	479	372	95
220	2800	315	245	62	220	3240	365	284	72	220	4340	488	380	96
225	2850	321	249	63	225	3300	371	289	73	225	4425	498	387	98
230	2900	326	254	64	230	3360	378	294	75	230	4510	507	395	100
235	2950	332	258	66	235	3420	385	299	76	235	4595	517	402	102
240	3000	338	263	67	240	3480	392	305	77	240	4680	527	410	104
245	3050	343	267	68	245	3540	398	310	79	245	4765	536	417	106
250	3100	349	271	69	250	3600	405	315	80	250	4850	546	424	108
255	3150	354	276	70	255	3660	412	320	81	255	4935	555	432	110
260	3200	360	280	71	260	3720	419	326	83	260	5020	565	439	112
265	3250	366	284	72	265	3780	425	331	84	265	5105	574	447	113
270	3300	371	289	73	270	3840	432	336	85	270	5190	584	454	115

GOAL: LOSE BODY FAT/INCREASE MUSCLE (BODY FAT UNDER 15%)

90 Minute Workouts														
BW	Sed	Carbs	Protein	Fat	BW	Mod	Carbs	Protein	Fat	BW	Act	Carbs	Protein	Fat
130	2200	248	193	49	130	2460	277	215	55	130	3110	350	272	69
135	2250	253	197	50	135	2520	284	221	56	135	3195	359	280	71
140	2300	259	201	51	140	2580	290	226	57	140	3280	369	287	73
145	2350	264	206	52	145	2640	297	231	59	145	3365	379	294	75
150	2400	270	210	53	150	2700	304	236	60	150	3450	388	302	77
155	2450	276	214	54	155	2760	311	242	61	155	3535	398	309	79
160	2500	281	219	56	160	2820	317	247	63	160	3620	407	317	80
165	2550	287	223	57	165	2880	324	252	64	165	3705	417	324	82
170	2600	293	228	58	170	2940	331	257	65	170	3790	426	332	84
175	2650	298	232	59	175	3000	338	263	67	175	3875	436	339	86
180	2700	304	236	60	180	3060	344	268	68	180	3960	446	347	88
185	2750	309	241	61	185	3120	351	273	69	185	4045	455	354	90
190	2800	315	245	62	190	3180	358	278	71	190	4130	465	361	92
195	2850	321	249	63	195	3240	365	284	72	195	4215	474	369	94
200	2900	326	254	64	200	3300	371	289	73	200	4300	484	376	96
205	2950	332	258	66	205	3360	378	294	75	205	4385	493	384	97
210	3000	338	263	67	210	3420	385	299	76	210	4470	503	391	99
215	3050	343	267	68	215	3480	392	305	77	215	4555	512	399	101
220	3100	349	271	69	220	3540	398	310	79	220	4640	522	406	103
225	3150	354	276	70	225	3600	405	315	80	225	4725	532	413	105
230	3200	360	280	71	230	3660	412	320	81	230	4810	541	421	107
235	3250	366	284	72	235	3720	419	326	83	235	4895	551	428	109
240	3300	371	289	73	240	3780	425	331	84	240	4980	560	436	111
245	3350	377	293	74	245	3840	432	336	85	245	5065	570	443	113
250	3400	383	298	76	250	3900	439	341	87	250	5150	579	451	114
255	3450	388	302	77	255	3960	446	347	88	255	5235	589	458	116
260	3500	394	306	78	260	4020	452	352	89	260	5320	599	466	118
265	3550	399	311	79	265	4080	459	357	91	265	5405	608	473	120
270	3600	405	315	80	270	4140	466	362	92	270	5490	618	480	122

BW = Body Weight • SED = Sedentary • MOD = Moderately Active • ACT = Active

GOAL: LOSE BODY FAT/INCREASE MUSCLE (BODY FAT UNDER 15%)

Non-Workout														
BW	Sed	Carbs	Protein	Fat	BW	Mod	Carbs	Protein	Fat	BW	Act	Carbs	Protein	Fat
130	1300	146	114	29	130	1560	176	137	35	130	2210	249	193	49
135	1350	152	118	30	135	1620	182	142	36	135	2295	258	201	51
140	1400	158	123	31	140	1680	189	147	37	140	2380	268	208	53
145	1450	163	127	32	145	1740	196	152	39	145	2465	277	216	55
150	1500	169	131	33	150	1800	203	158	40	150	2550	287	223	57
155	1550	174	136	34	155	1860	209	163	41	155	2635	296	231	59
160	1600	180	140	36	160	1920	216	168	43	160	2720	306	238	60
165	1650	186	144	37	165	1980	223	173	44	165	2805	316	245	62
170	1700	191	149	38	170	2040	230	179	45	170	2890	325	253	64
175	1750	197	153	39	175	2100	236	184	47	175	2975	335	260	66
180	1800	203	158	40	180	2160	243	189	48	180	3060	344	268	68
185	1850	208	162	41	185	2220	250	194	49	185	3145	354	275	70
190	1900	214	166	42	190	2280	257	200	51	190	3230	363	283	72
195	1950	219	171	43	195	2340	263	205	52	195	3315	373	290	74
200	2000	225	175	44	200	2400	270	210	53	200	3400	383	298	76
205	2050	231	179	46	205	2460	277	215	55	205	3485	392	305	77
210	2100	236	184	47	210	2520	284	221	56	210	3570	402	312	79
215	2150	242	188	48	215	2580	290	226	57	215	3655	411	320	81
220	2200	248	193	49	220	2640	297	231	59	220	3740	421	327	83
225	2250	253	197	50	225	2700	304	236	60	225	3825	430	335	85
230	2300	259	201	51	230	2760	311	242	61	230	3910	440	342	87
235	2350	264	206	52	235	2820	317	247	63	235	3995	449	350	89
240	2400	270	210	53	240	2880	324	252	64	240	4080	459	357	91
245	2450	276	214	54	245	2940	331	257	65	245	4165	469	364	93
250	2500	281	219	56	250	3000	338	263	67	250	4250	478	372	94
255	2550	287	223	57	255	3060	344	268	68	255	4335	488	379	96
260	2600	293	228	58	260	3120	351	273	69	260	4420	497	387	98
265	2650	298	232	59	265	3180	358	278	71	265	4505	507	394	100
270	2700	304	236	60	270	3240	365	284	72	270	4590	516	402	102

GOAL: MAINTAIN BODY FAT/INCREASE MUSCLE

60 Minute Workouts														
BW	Sed	Carbs	Protein	Fat	BW	Mod	Carbs	Protein	Fat	BW	Act	Carbs	Protein	Fat
130	2290	258	200	51	130	2550	287	223	57	130	3200	370	280	71
135	2355	265	206	52	135	2625	295	230	58	135	3300	371	289	73
140	2420	272	212	54	140	2700	304	236	60	140	3400	383	298	76
145	2485	280	217	55	145	2775	312	243	62	145	3500	394	306	78
150	2550	287	223	57	150	2850	321	249	63	150	3600	405	315	80
155	2615	294	229	58	155	2925	329	256	65	155	3700	416	324	82
160	2680	302	235	60	160	3000	338	263	67	160	3800	428	333	84
165	2745	309	240	61	165	3075	346	269	68	165	3900	439	341	87
170	2810	316	246	62	170	3150	354	276	70	170	4000	450	350	89
175	2875	323	252	64	175	3225	363	282	72	175	4100	461	359	91
180	2940	331	257	65	180	3300	371	289	73	180	4200	473	368	93
185	3005	338	263	67	185	3375	380	295	75	185	4300	484	376	96
190	3070	345	269	68	190	3450	388	302	77	190	4400	495	385	98
195	3135	353	274	70	195	3525	397	308	78	195	4500	506	394	100
200	3200	360	280	71	200	3600	405	315	80	200	4600	518	403	102
205	3265	367	286	73	205	3675	413	322	82	205	4700	529	411	104
210	3330	375	291	74	210	3750	422	328	83	210	4800	540	420	107
215	3395	382	297	75	215	3825	430	335	85	215	4900	551	429	109
220	3460	389	303	77	220	3900	439	341	87	220	5000	563	438	111
225	3525	397	308	78	225	3975	447	348	88	225	5100	574	446	113
230	3590	404	314	80	230	4050	456	354	90	230	5200	585	455	116
235	3655	411	320	81	235	4125	464	361	92	235	5300	596	464	118
240	3720	419	326	83	240	4200	473	368	93	240	5400	608	473	120
245	3785	426	331	84	245	4275	481	374	95	245	5500	619	481	122
250	3850	433	337	86	250	4350	489	381	97	250	5600	630	490	124
255	3915	440	343	87	255	4425	498	387	98	255	5700	641	499	127
260	3980	448	348	88	260	4500	506	394	100	260	5800	653	508	129
265	4045	455	354	90	265	4575	515	400	102	265	5900	664	516	131
270	4110	462	360	91	270	4650	523	407	103	270	6000	675	525	133

BW = Body Weight • SED = Sedentary • MOD = Moderately Active • ACT = Active

GOAL: MAINTAIN BODY FAT/INCREASE MUSCLE**90 Minute Workouts**

BW	Sed	Carbs	Protein	Fat	BW	Mod	Carbs	Protein	Fat	BW	Act	Carbs	Protein	Fat
130	2590	291	227	58	130	2850	321	249	63	130	3500	394	306	78
135	2655	299	232	59	135	2925	329	256	65	135	3600	405	315	80
140	2720	306	238	60	140	3000	338	263	67	140	3700	416	324	82
145	2785	313	244	62	145	3075	346	269	68	145	3800	428	333	84
150	2850	321	249	63	150	3150	354	276	70	150	3900	439	341	87
155	2915	328	255	65	155	3225	363	282	72	155	4000	450	350	89
160	2980	335	261	66	160	3300	371	289	73	160	4100	461	359	91
165	3045	343	266	68	165	3375	380	295	75	165	4200	473	368	93
170	3110	350	272	69	170	3450	388	302	77	170	4300	484	376	96
175	3175	357	278	71	175	3525	397	308	78	175	4400	495	385	98
180	3240	365	284	72	180	3600	405	315	80	180	4500	506	394	100
185	3305	372	289	73	185	3675	413	322	82	185	4600	518	403	102
190	3370	379	295	75	190	3750	422	328	83	190	4700	529	411	104
195	3435	386	301	76	195	3825	430	335	85	195	4800	540	420	107
200	3500	394	306	78	200	3900	439	341	87	200	4900	551	429	109
205	3565	401	312	79	205	3975	447	348	88	205	5000	563	438	111
210	3630	408	318	81	210	4050	456	354	90	210	5100	574	446	113
215	3695	416	323	82	215	4125	464	361	92	215	5200	585	455	116
220	3760	423	329	84	220	4200	473	368	93	220	5300	596	464	118
225	3825	430	335	85	225	4275	481	374	95	225	5400	608	473	120
230	3890	438	340	86	230	4350	489	381	97	230	5500	619	481	122
235	3955	445	346	88	235	4425	498	387	98	235	5600	630	490	124
240	4020	452	352	89	240	4500	506	394	100	240	5700	641	499	127
245	4085	460	357	91	245	4575	515	400	102	245	5800	653	508	129
250	4150	467	363	92	250	4650	523	407	103	250	5900	664	516	131
255	4215	474	369	94	255	4725	532	413	105	255	6000	675	525	133
260	4280	482	375	95	260	4800	540	420	107	260	6100	686	534	136
265	4345	489	380	97	265	4875	548	427	108	265	6200	698	543	138
270	4410	496	386	98	270	4950	557	433	110	270	6300	709	551	140

GOAL: MAINTAIN BODY FAT/INCREASE MUSCLE

Non-Workout														
BW	Sed	Carbs	Protein	Fat	BW	Mod	Carbs	Protein	Fat	BW	Act	Carbs	Protein	Fat
130	1690	190	148	38	130	1950	219	171	43	130	2600	293	228	58
135	1755	197	154	39	135	2025	228	177	45	135	2700	304	236	60
140	1820	205	159	40	140	2100	236	184	47	140	2800	315	245	62
145	1885	212	165	42	145	2175	245	190	48	145	2900	326	254	64
150	1950	219	171	43	150	2250	253	197	50	150	3000	338	263	67
155	2015	227	176	45	155	2325	262	203	52	155	3100	349	271	69
160	2080	234	182	46	160	2400	270	210	53	160	3200	360	280	71
165	2145	241	188	48	165	2475	278	217	55	165	3300	371	289	73
170	2210	249	193	49	170	2550	287	223	57	170	3400	383	298	76
175	2275	256	199	51	175	2625	295	230	58	175	3500	394	306	78
180	2340	263	205	52	180	2700	304	236	60	180	3600	405	315	80
185	2405	271	210	53	185	2775	312	243	62	185	3700	416	324	82
190	2470	278	216	55	190	2850	321	249	63	190	3800	428	333	84
195	2535	285	222	56	195	2925	329	256	65	195	3900	439	341	87
200	2600	293	228	58	200	3000	338	263	67	200	4000	450	350	89
205	2665	300	233	59	205	3075	346	269	68	205	4100	461	359	91
210	2730	307	239	61	210	3150	354	276	70	210	4200	473	368	93
215	2795	314	245	62	215	3225	363	282	72	215	4300	484	376	96
220	2860	322	250	64	220	3300	371	289	73	220	4400	495	385	98
225	2925	329	256	65	225	3375	380	295	75	225	4500	506	394	100
230	2990	336	262	66	230	3450	388	302	77	230	4600	518	403	102
235	3055	344	267	68	235	3525	397	308	78	235	4700	529	411	104
240	3120	351	273	69	240	3600	405	315	80	240	4800	540	420	107
245	3185	358	279	71	245	3675	413	322	82	245	4900	551	429	109
250	3250	366	284	72	250	3750	422	328	83	250	5000	563	438	111
255	3315	373	290	74	255	3825	430	335	85	255	5100	574	446	113
260	3380	380	296	75	260	3900	439	341	87	260	5200	585	455	116
265	3445	388	301	77	265	3975	447	348	88	265	5300	596	464	118
270	3510	395	307	78	270	4050	456	354	90	270	5400	608	473	120

BW = Body Weight • SED = Sedentary • MOD = Moderately Active • ACT = Active

GOAL: GAIN BODY FAT/INCREASE MUSCLE**60 Minute Workouts**

BW	Sed	Carbs	Protein	Fat	BW	Mod	Carbs	Protein	Fat	BW	Act	Carbs	Protein	Fat
130	2680	302	235	60	130	2940	331	257	65	130	3590	404	314	80
135	2760	311	242	61	135	3030	341	265	67	135	3705	417	324	82
140	2840	320	249	63	140	3120	351	273	69	140	3820	430	334	85
145	2920	329	256	65	145	3210	361	281	71	145	3935	443	344	87
150	3000	338	263	67	150	3300	371	289	73	150	4050	456	354	90
155	3080	347	270	68	155	3390	381	297	75	155	4165	469	364	93
160	3160	356	277	70	160	3480	392	305	77	160	4280	482	375	95
165	3240	365	284	72	165	3570	402	312	79	165	4395	494	385	98
170	3320	374	291	74	170	3660	412	320	81	170	4510	507	395	100
175	3400	383	298	76	175	3750	422	328	83	175	4625	520	405	103
180	3480	392	305	77	180	3840	432	336	85	180	4740	533	415	105
185	3560	401	312	79	185	3930	442	344	87	185	4855	546	425	108
190	3640	410	319	81	190	4020	452	352	89	190	4970	559	435	110
195	3720	419	326	83	195	4110	462	360	91	195	5085	572	445	113
200	3800	428	333	84	200	4200	473	368	93	200	5200	585	455	116
205	3880	437	340	86	205	4290	483	375	95	205	5315	598	465	118
210	3960	446	347	88	210	4380	493	383	97	210	5430	611	475	121
215	4040	455	354	90	215	4470	503	391	99	215	5545	624	485	123
220	4120	464	361	92	220	4560	513	399	101	220	5660	637	495	126
225	4200	473	368	93	225	4650	523	407	103	225	5775	650	505	128
230	4280	482	375	95	230	4740	533	415	105	230	5890	663	515	131
235	4360	491	382	97	235	4830	543	423	107	235	6005	676	525	133
240	4440	500	389	99	240	4920	554	431	109	240	6120	689	536	136
245	4520	509	396	100	245	5010	564	438	111	245	6235	701	546	139
250	4600	518	403	102	250	5100	574	446	113	250	6350	714	556	141
255	4680	527	410	104	255	5190	584	454	115	255	6465	727	566	144
260	4760	536	417	106	260	5280	594	462	117	260	6580	740	576	146
265	4840	545	424	108	265	5370	604	470	119	265	6695	753	586	149
270	4920	554	431	109	270	5460	614	478	121	270	6810	766	596	151

GOAL: GAIN BODY FAT/INCREASE MUSCLE

90 Minute Workouts														
BW	Sed	Carbs	Prot	Fat	BW	Mod	Carbs	Prot	Fat	BW	Act	Carbs	Prot	Fat
130	2980	335	261	66	130	3240	365	284	72	130	3890	438	340	86
135	3060	344	268	68	135	3330	375	291	74	135	4005	451	350	89
140	3140	353	275	70	140	3420	385	299	76	140	4120	464	361	92
145	3220	362	282	72	145	3510	395	307	78	145	4235	476	371	94
150	3300	371	289	73	150	3600	405	315	80	150	4350	489	381	97
155	3380	380	296	75	155	3690	415	323	82	155	4465	502	391	99
160	3460	389	303	77	160	3780	425	331	84	160	4580	515	401	102
165	3540	398	310	79	165	3870	435	339	86	165	4695	528	411	104
170	3620	407	317	80	170	3960	446	347	88	170	4810	541	421	107
175	3700	416	324	82	175	4050	456	354	90	175	4925	554	431	109
180	3780	425	331	84	180	4140	466	362	92	180	5040	567	441	112
185	3860	434	338	86	185	4230	476	370	94	185	5155	580	451	115
190	3940	443	345	88	190	4320	486	378	96	190	5270	593	461	117
195	4020	452	352	89	195	4410	496	386	98	195	5385	606	471	120
200	4100	461	359	91	200	4500	506	394	100	200	5500	619	481	122
205	4180	470	366	93	205	4590	516	402	102	205	5615	632	491	125
210	4260	479	373	95	210	4680	527	410	104	210	5730	645	501	127
215	4340	488	380	96	215	4770	537	417	106	215	5845	658	511	130
220	4420	497	387	98	220	4860	547	425	108	220	5960	671	522	132
225	4500	506	394	100	225	4950	557	433	110	225	6075	683	532	135
230	4580	515	401	102	230	5040	567	441	112	230	6190	696	542	138
235	4660	524	408	104	235	5130	577	449	114	235	6305	709	552	140
240	4740	533	415	105	240	5220	587	457	116	240	6420	722	562	143
245	4820	542	422	107	245	5310	597	465	118	245	6535	735	572	145
250	4900	551	429	109	250	5400	608	473	120	250	6650	748	582	148
255	4980	560	436	111	255	5490	618	480	122	255	6765	761	592	150
260	5060	569	443	112	260	5580	628	488	124	260	6880	774	602	153
265	5140	578	450	114	265	5670	638	496	126	265	6995	787	612	155
270	5220	587	457	116	270	5760	648	504	128	270	7110	800	622	158

BW = Body Weight • SED = Sedentary • MOD = Moderately Active • ACT = Active

GOAL: GAIN BODY FAT/INCREASE MUSCLE


Non-Workout														
BW	Sed	Carbs	Prot	Fat	BW	Mod	Carbs	Prot	Fat	BW	Act	Carbs	Prot	Fat
130	2080	234	182	46	130	2340	263	205	52	130	2990	336	262	66
135	2160	243	199	48	135	2430	273	213	54	135	3105	349	272	69
140	2240	252	196	50	140	2520	284	221	56	140	3220	362	282	72
145	2320	261	203	52	145	2610	294	228	58	145	3335	375	292	74
150	2400	270	210	53	150	2700	304	236	60	150	3450	388	302	77
155	2480	279	217	55	155	2790	314	244	62	155	3565	401	312	79
160	2560	288	224	57	160	2880	324	252	64	160	3680	414	322	82
165	2640	297	231	59	165	2970	334	260	66	165	3795	427	332	84
170	2720	306	238	60	170	3060	344	268	68	170	3910	440	342	87
175	2800	315	245	62	175	3150	354	276	70	175	4025	453	352	89
180	2880	324	252	64	180	3240	365	284	72	180	4140	466	362	92
185	2960	333	259	66	185	3330	375	291	74	185	4255	479	372	95
190	3040	342	266	68	190	3420	385	299	76	190	4370	492	382	97
195	3120	351	273	69	195	3510	395	307	78	195	4485	505	392	100
200	3200	360	280	71	200	3600	405	315	80	200	4600	518	403	102
205	3280	369	287	73	205	3690	415	323	82	205	4715	530	413	105
210	3360	378	294	75	210	3780	425	331	84	210	4830	543	423	107
215	3440	387	301	76	215	3870	435	339	86	215	4945	556	433	110
220	3520	396	308	78	220	3960	446	347	88	220	5060	569	443	112
225	3600	405	315	80	225	4050	456	354	90	225	5175	582	453	115
230	3680	414	322	82	230	4140	466	362	92	230	5290	595	463	118
235	3760	423	329	84	235	4230	476	370	94	235	5405	608	473	120
240	3840	432	336	85	240	4320	486	378	96	240	5520	621	483	123
245	3920	441	343	87	245	4410	496	386	98	245	5635	634	493	125
250	4000	450	350	89	250	4500	506	394	100	250	5750	647	503	128
255	4080	459	357	91	255	4590	516	402	102	255	5865	660	513	130
260	4160	468	364	92	260	4680	527	410	104	260	5980	673	523	133
265	4240	477	371	94	265	4770	537	417	106	265	6095	686	533	135
270	4320	486	378	96	270	4860	547	425	108	270	6210	699	543	138



BW = Body Weight • SED = Sedentary • MOD = Moderately Active • ACT = Active

APPENDIX B

Macrobolic Food Guide Pyramids

The following Macroboolic Food Guide Pyramids allow you to allot the proper amounts of food to meet the Macroboolic Lean-Mass Equation. Athletes who are on the lower end of the calorie range would choose the lower servings. Athletes who are on the higher end of the calorie range would choose the higher servings. The Macroboolic Food Guide Pyramids are to be used in conjunction with the Macroboolic Exchange Lists (see Appendix C). The proper serving sizes are predetermined in the Macroboolic Exchange Lists along with the different foods that are suggested for consumption.





APPENDIX C

Macrobiotic Exchange Lists

The Macrobiotic Exchange Lists allows you to utilize the Macrobiotic Food Guide Pyramids (see Appendix B) in meal planning. The Macrobiotic Exchange Lists will enable you to effectively “exchange” one serving from one list for another food item on the same list. For example, 1 cup of raw mushrooms has the same nutritional value as a half cup of cooked broccoli. Everything is spelled out for you, so plan, eat, and GROW!

LOW-GLYCEMIC VEGETABLES

**Macrobiotic Exchange: 1 Vegetable =
5 g carbohydrate • 2 g protein • 0 g fat • 25 calories**

All vegetables 1 cup raw or 1/2 cup cooked

Alfalfa sprouts*	Cucumbers (free)*	Radish*
Asparagus*	Eggplant*	Raw spinach (free)*
Bamboo shoots*	Green beans*	Sauerkraut*
Broccoli*	Green peppers*	Summer squash*
Brussels sprouts*	Lettuce, all varieties (free)*	Tomato*
Cabbage*	Mushrooms*	Turnip*
Cauliflower*	Onions *	Zucchini*
Celery*		

**GI values not established*

LOW-GLYCEMIC ROOT VEGETABLES

**Macrobiotic Exchange: 1 Starch =
15 g carbohydrate • 3 g protein • 1 g fat • 80 calories**

Carrots, raw (1/2 cup)	Sweet potatoes (3 oz)
New potatoes (3 oz)	White potatoes, boiled) (3 oz)
Pontiac potatoes, boiled (3 oz)	Yams, baked (2 oz)
Russet potatoes, baked w/o skin (3 oz)	

LOW-GLYCEMIC BEANS

**Macrobiotic Exchange: 1 Very Lean Starch =
15–20 g carbohydrate • 7–10 g protein • 1 g fat • 115 calories**

All beans/peas cooked w/o salt (1/2 cup)

Baked beans	Lentils	Peas
Butter beans	Lima beans	Pinto beans
Kidney beans	Marrowfat beans	Soya bean
Kidney beans (white)	Mung beans	Split pea

LOW-GLYCEMIC FRUITS

**Macrobiotic Exchange: 1 Fruit =
15 g carbohydrate • 60 calories**

Apple, small (4 oz)	Grapefruit, 1/2 large (11 oz)	Peach, medium (6 oz)
Apples, dried (4 rings)	Grapes, small 17 (3 oz)	Pear, medium
Apricots, dried (8 halves)	Kiwi (3.5 oz)	Pineapple (3/4 cup)
Banana, small (4 oz)	Mandarin orange (1/2 cup)	Strawberries (1 1/4 cup whole berries)
Cherries (1/3 cup)	Mango, 1/2 small (8 oz)	

LOW-GLYCEMIC GRAINS

**Macrobiotic Exchange: 1 Starch =
15 g carbohydrate • 3 g protein • 1 g fat • 80 calories**

Amaranth, uncooked (2 tbs)	Pasta (seminola), cooked 1/2 cup
Barley, cooked (1/2 cup)	Pita, whole-wheat, six inch (1/2)
Brown rice, steamed (3/8 cup)	Pumpnickel bread (1 slice)
Buckwheat, uncooked (2 tbs)	Rye bread (1 slice)
Bulgur, cooked (1/3 cup)	Semolina, uncooked (2 tbs)
Corn tortilla, six inch	Wheat tortilla, six inch

Durum, uncooked (2 tbs)	Whole-wheat bagel, small (1/2)
Oatmeal (1/2 cup)	Whole-wheat bread (1 slice)

LOW-GLYCEMIC/LOW-FAT DAIRY/MILK

**Macrobiotic Exchange: 1 Dairy =
12 g carbohydrate • 8 g protein • 0–3 g fat • 90 calories**

Skim Milk (1 cup)

MEATS AND MEAT SUBSTITUTES

**Macrobiotic Exchange: 1 Very Lean Meat =
0 g carbohydrate • 7 g protein • 0–1 g fat • 35 calories**

Meats, No Beef (all 1 ounce)

Chicken or turkey (white meat only, no skin)	Clams, crab, lobster, scallops, shrimp
Buffalo, ostrich	Duck, pheasant (no skin)
Fresh cod, flounder, haddock, halibut, trout, tuna (canned in water)	

Meat Substitutes

Egg whites (2 large)	Fat-free cheese (1 oz)
Nonfat/low-fat cottage cheese (1/4 cup)	

**Macrobiotic Exchange: 1 Lean Meat =
0 g carbohydrate • 7 g protein • 3 g fat • 55 calories**

Meats (all 1 ounce)

Goose (no skin)	Lamb, roast, chop, or leg	Rabbit
Herring, salmon, catfish	Lean pork	Veal, lean chop, roast

USDA **Select** or **Choice** grades of lean beef **trimmed of fat**.

For example: round, sirloin, flank steak, tenderloin, roast (rib, chunk, rump) steak (T-bone, porterhouse, cubed), ground round.

Meat Substitutes

Any cheese with 3 g of fat or less per 1 oz	Grated parmesan (2 tbs)
Low-fat Cheddar cheese (1 oz)	

**Macrobiotic Exchange: 1 Medium Fat =
0 g carbohydrate • 7 g protein • 5 g fat • 75 calories**

Meats (all 1 ounce)

Most beef products fall into this category (ground beef, meatloaf, prime grades of meat trimmed of fat, for example, prime rib)

Chicken (dark meat, with skin)

Meat Substitutes

Feta, mozzarella cheese (1 oz) Ricotta cheese ($\frac{1}{4}$ cup, 2 oz) Whole egg (1 large)

**Macrobiotic Exchange: 1 High-Fat Meat =
0 g carbohydrate • 7 g protein • 8 g fat • 100 calories**

All regular cheeses such as American, Cheddar, Monterey Jack, and Swiss (1 oz)

Pork spareribs, ground pork (1 oz)

FAT

**Macrobiotic Exchange: 1 Fat =
5 g fat • 45 calories**

Monounsaturated fats

Avocado, medium (1 oz) Peanut butter (2 tsp)

Nuts: almonds, cashews (6 nuts) Peanuts (10 nuts)

Oil: canola, olive, peanut (1 tsp) Pecans (4 halves)

Olives, ripe, black (8 large) Seeds: sesame seeds (1 tbs)

Polyunsaturated fats

Flaxseeds (1 tbs) Oil: corn, safflower, soybean (1 tsp)

Mayonnaise, reduced fat (1 tsp) Seeds: pumpkin, sunflower seeds (1 tbs)

Nuts: walnuts (4 halves)

Saturated fats (use sparingly)

Butter, stick (1 tsp) Reduced-fat sour cream (3 tbs)

Reduced-fat cream cheese (1 oz)

SUPPLEMENTS

**Macrobiotic Exchange: 3 Very Lean Meat =
2–3 g carbohydrate • 20 g protein • 2–3 g fat • 120 calories**

MHP's Probiotic (any flavor): 1 scoop (approx. 30 g)

**Macrobiotic Exchange: Macrobiotic Complete Supplement =
15 g carbohydrate • 11 g protein • 3 g fat • 130 calories**

MHP's Up Your MASS (any flavor): 1 scoop (33 g)

References

- Almada, A., et al. "Effects of B-BHBM Supplementation with and without Creatine during Training on Strength and Sprint Capacity." *Federation of American Societies of Experimental Biology Journal* 11 (1997): A374.
- Anderson, Helen L., Mary Belle Heindel, and Hellen Linkswiler. "Effect on Nitrogen Balance of Adult Man of Varying Source of Nitrogen and Level of Calorie Intake." *Journal of Nutrition* (1969): 82–90.
- Anderson J.W., B. M. Johnstone, and M. E. Cook-Newell. "Meta-Analysis of the Effects of Soy Protein Intake on Serum Lipids." *New England Journal of Medicine* 333, 5 (1995): 276–82.
- Anderson, M., et al. "Pre-Exercise Meal Affects Ride Time to Fatigue in Trained Cyclists." *Journal of the American Dietetic Association* 94 (1994): 1152–1153.
- Apfelbaum, Marian, Jacques Fricker, and Lawrence Igoin-Apfelbaum. "Low and Very Low Calorie Diets." *American Journal of Clinical Nutrition* 45 (1987): 1126–1134.
- Armstrong, R.B. "Mechanisms of Exercise-Induced Delayed Onset Muscular Soreness: A Brief Review." *Medicine and Science in Sports and Exercise* 16 no. 6 (1984): 529–538.
- . "Muscle Damage and Endurance Events." *Sports Medicine* 3 (1986): 370–381.
- Ball, T., et al. "Periodic Carbohydrate Replacement During 50 Minutes of High-Intensity Cycling Improves Subsequent Sprint Performance." *International Journal of Sport Science* 5(2) (June 1995): 151–158.
- Bamman, M.M., et al. "Changes in Body Composition, Diet, and Strength of Body-builders During the 12 Weeks Prior to Competition." *Journal of Sports Medicine and Physical Fitness* 33 (1993): 383.
- Bell, D.G., et al. "Effects of Caffeine, Ephedrine and the Combination on Time to Exhaustion during High-Intensity Exercise." *European Journal of Applied Physiology* 77 (1998): 427–433.

- Bergstrom, Jonas, and Eric Hultman. "Nutrition for Maximal Sports Performance." *Journal of the American Medical Association* 221, no. 9 (1972): 999–1004.
- Berning, J.R. "The Role of Medium-Chain Triglycerides in Exercise." *International Journal of Sport Nutrition* 6, no. 3 (1996): 121–133.
- Bohmer, D., et al. "Treatment of Chondropathia Patellae in Young Athletes with Glucosamine Sulfate." In *Current Topics in Sports Medicine*, N. Bachl, L. Prokop, and R. Suckert, eds. Vienna: Urban & Schwarzenberg, 1984, pp. 799–803.
- Boire, Y., et al. "Slow and fast dietary proteins differentially modulate postprandial protein accretion." *Proceedings of the National Academy of Science* 94(26) (Dec 1997): 14930–14935.
- Bonde-Petersen, Flemming, Howard G. Knuttgen, and Jan Henriksson. "Muscle Metabolism During Exercise with Concentric and Eccentric Contractions." *Journal of Applied Physiology* 33 (1972): 792–795.
- Boozer, C.N., P.A. Daly, P. Homel, et al. "Herbal Ephedra/Caffeine for Weight Loss: A 6-Month Randomized Safety and Efficacy Trial." *International Journal of Obesity Related Metabolic Disorders* 26, no. 5 (May 2002): 593–604.
- Boyne, P.S., and H. Medhurst. "Oral Anti-inflammatory Enzyme Therapy in Injuries in Professional Footballers." *The Practitioner* 198 (April 1967): 543–546.
- Braatan, J.T., F.W. Scott, P.J. Wood, et al. "High Beta-glucan Oat Bran and Oat Grain Reduce Post-prandial Blood Glucose and Insulin in Subjects with and without Type 2 Diabetes." *Diabetic Medicine* 11 (1994): 312–318.
- Braatan, J.T., P.J. Word, F.W. Scott, et al. "Oat Grain Lowers Glucose and Insulin After an Oral Glucose Load." *American Journal of Clinical Nutrition* 53 (1991): 1425–1430.
- Brilla, L.R., and T.E. Landerholm. "Effect of Fish Oil Supplementation and Exercise on Serum Lipids and Aerobic Fitness." *Journal of Sports Medicine and Physical Fitness* 30, no. 2 (1990): 173–180.
- Brose, A., G. Parise, and M.A. Tarnopolsky. "Creatine Supplementation Enhances Isometric Strength and Body Composition Improvements Following Strength Exercise Training in Older Adults." *Journal of Gerontology Series A: Biological Sciences and Medical Sciences* 58, no. 1 (January 2003): 9–11.
- Brown, C., and Jack H. Wilmore. "The Effects of Maximal Resistance Training on the Strength and Body Composition of Women Athletes." *Medicine and Science in Sports* 6, no. 3 (1974): 174–177.
- Bucci, L. *Nutrients as Ergogenic Aids for Sports and Exercise*. Boca Raton, FL: CRC Press, 1993.
- . *Nutrition Applied to Injury Rehabilitation and Sports Medicine*. Boca Raton, FL: CRC Press, 1995.
- Burke D.G., P.D. Chilibeck, K.S. Davidson, et al. "The Effect of Whey Protein Supplementation with and without Creatine Monohydrate Combined with Resistance Train-

- ing on Lean Tissue Mass and Muscle Strength.” *International Journal of Sport Nutrition and Exercise Metabolism* 11, no. 3 (September 2001): 349–64.
- Burke, D.G., P.D. Chilibeck, G. Parise, et al. “Effect of Creatine and Weight Training on Muscle Creatine and Performance in Vegetarians.” *Medicine & Science in Sports and Exercise* 35, no. 11 (November 2003): 1946–55.
- Burke, D.G., S. Silver, L.E. Holt, et al. “The Effect of Continuous Low Dose Creatine Supplementation on Force, Power, and Total Work.” *International Journal of Sport Nutrition and Exercise Metabolism* 10, no. 3 (September 2000): 235–44.
- Burke, L.M., and S.D. Read. “Dietary Supplements in Sport.” *Sports Medicine* 15 (1993): 43–65.
- Buskirk, Elsworth R., and José Mendez. “Sports Science and Body Composition Analysis: Emphasis on Cell and Muscle Mass.” *Medicine and Science in Sports and Exercise* 16, no. 6 (1984): 584–593.
- Butterfield, G. “Ergogenic Aids: Evaluating Sport Nutrition Products.” *International Journal of Sport Nutrition* no. 3 (1996): 191–197.
- Butterfield, Gail E., and Doris H. Calloway. “Physical Activity Improves Protein Utilization in Young Men.” *British Journal of Nutrition* 51 (1984): 171–184.
- Calver, A., et al., “Dilator Actions of Arginine in Human Peripheral Vasculature,” *Clinical Science (Colch)*, 85, no. 5 (1991): 695–700.
- Casanueva, F.F., L. Villanueva, J.A. Cabranes, et al. “Cholinergic Mediation of Growth Hormone Secretion Elicited by Arginine, Clonidine, and Physical Exercise in Man.” *Journal of Clinical Endocrinology and Metabolism* 59, no. 3 (1984): 526–530.
- Celejowa, I., and M. Homa. “Food Intake, Nitrogen and Energy Balance in Polish Weight Lifters, During Training Camp.” *Nutrition and Metabolism* 12 (1970): 259–274.
- Chandalia, M., A. Garg, D. Lutjohann, et al. “Beneficial Effects of High Dietary Fiber Intake in Patients with Type 2 Diabetes Mellitus.” *New England Journal of Medicine* 342 (2000): 1392–1398.
- Chang, Tse Wen, and Alfred L. Goldberg. “The Metabolic Fates of Amino Acids and the Formation of Glutamine in Skeletal Muscle.” *Journal of Biological Chemistry* 253, no. 10 (1978): 3685–3695.
- Chin, S. “Dietary Sources of Conjugated Dienoic Isomers of Linoleic Acid, a Newly Recognized Class of Anticarcinogens.” *Journal of Food Composition and Analysis* 5 (1992): 185–195.
- Chin, S., J. Storkron, K. Albright, et al. “Conjugated Linoleic Acid is a Growth Factor for Rats as Shown by Enhanced Weight Gain and Improved Feed Efficiency.” *Journal of Nutrition* 124 (1994): 2344–2349.
- Christensen, H. “Muscle Activity and Fatigue in the Shoulder Muscles during Repetitive Work.” *European Journal of Applied Physiology* 54 (1986): 596–601.

- Clarkson, P., and E. Haymes. "Trace Mineral Requirements for Athletes." *International Journal of Sports Nutrition* 4 (1994): 104.
- Clarkson, Priscilla M., Walter Kroll, and Thomas C. McBride. "Plantar Flexion Fatigue and Muscle Fiber Type in Power and Endurance Athletes." *Medicine and Science in Sports and Exercise* 12 (1980): 262–267.
- Colker, C.M. "Immune Status of Elite Athletes: Role of Whey Protein Concentrate: A Review." *Medicine and Science in Sports and Exercise* 30 (1998): S17.
- Conzolazio, C. Frank, Herman L. Johnson, Richard A. Nelson, et al. "Protein Metabolism during Intensive Physical Training in the Young Adult." *American Journal of Clinical Nutrition* 28 (1975): 29–35.
- Cook, James D., and Elaine R. Monsen. "Vitamin C, the Common Cold, and Iron Absorption." *American Journal of Clinical Nutrition* 30(2) (Feb 1977): 235–241.
- Copinschi, Georges, Laurence C. Wegienka, Satoshi Hane, et al. "Effect of Arginine on Serum Levels of Insulin and Growth Hormone in Obese Subjects." *Metabolism* 16 (1967): 485–491.
- Cossack, Zafrallah T., Ananda Prasad. "Effect of Protein Source on the Bioavailability of Zinc in Human Subjects." *Nutrition Research* 3 (1983): 23–31.
- Costill, D.L., R. Bowers, et al. "Muscle Glycogen Utilization during Prolonged Exercise on Successive Days." *Journal of Applied Physiology* 31 (1971): 834–838.
- Costill, D.L., and M. Hargreaves. "Carbohydrate Nutrition and Fatigue." *Sports Medicine* 13(1992): 86.
- Costill, D.L., W. M. Sherman, et al. "The Role of Dietary Carbohydrate in Muscle Glycogen Synthesis after Strenuous Running." *American Journal of Clinical Nutrition* 34 (1981): 1834–1836.
- Coyle, Edward F., and Andrew R. Coggan. "Effectiveness of Carbohydrate Feeding in Delaying Fatigue during Pro-longed Exercise." *Sports Medicine* 1(6) (Nov–Dec 1984): 446–458.
- Davies, Kelvin J. A., Alexandre T. Quintanilha, et al. "Free Radicals and Tissue Damage Produced by Exercise." *Biochemical and Biophysical Research Communications* 107, no. 4 (1982): 1198–1205.
- Despres, J. P., C. Bouchard, and R. Savard, et al. "Level of Physical Fitness and Adipocyte Lipolysis in Humans." *The American Physiological Society* (1984): 1157–1161.
- "Diets Containing Less than 42% Carbohydrates Do Not Meet the Energy Demands or Provide Adequate Glycogen Stores for Bodybuilders and Their Intense Workouts." *Strength and Conditioning Journal* 24(2002):42–53
- DiPrampero, P. Enrico. "Energetics of Muscular Exercise." *Biochemical Pharmacology* 89 (1981): 143–209.
- Dohm, G. Lynis, George J. Kasperek, and Edward B. Tapscott, et al. "Effect of Exercise on Synthesis and Degradation of Muscle Protein." *Biochemical Journal* 188 (1980): 255–262.

- Dray, F. "Role of Prostaglandins in Growth Hormone Secretion." *Advanced Prostaglandin and Thromboxane Research* 8 (1980): 1321.
- "Effects of Branched Chain Amino Acid Supplementation Before and After Training." *Medicina Dello Sport* 50 (1997): 293–303.
- Ehn, Lars, Bjorn Carlmark, and Sverker Hoglund. "Iron Status in Athletes Involved in Intense Physical Activity." *Medicine and Science in Sports and Exercise* 12, no. 1 (1980): 61–64.
- Engelhardt, M., G. Neumann, A. Berbalk, et al. "Creatine Supplementation in Endurance Sports." *Medicine and Science in Sports and Exercise* 30 (1998): 1123–1129.
- Erling, T.A. "Pilot Study With the Aim of Studying the Efficacy and Tolerability of CLA (Tonalin) on the Body Composition in Humans." Lillestrom, Norway: Medstat Research Ltd, 1997.
- "Fatigue and Underperformance in Athletes." *British Journal of Sports Nutrition* 32 (1998): 107–110.
- Ferreira, M., R. Kreider, M. Wilson, et al. "Effects of Conjugated Linoleic Acid (CLA) Supplementation during Resistance Training on Body Composition and Strength." *Journal of Strength and Conditioning Research* 11 (1997): 280.
- Food and Agriculture Organization of the United Nations. *The Amino Acid Content of Foods and Biological Data on Proteins. Nutritional Study #24*. Rome Lanham, MD: UNIPUB, 1970.
- Food and Nutrition Board. Recommended Dietary Allowances*, 9th Edition. Washington, D.C.: National Academy of Sciences, 1980.
- Forbes, Gilbert B. "Growth of the Lean Body Mass in Man." *Growth* 36 (1972): 325–338.
- . "Body Composition as Affected by Physical Activity and Nutrition." *Metabolic and Nutritional Aspects of Physical Exercise: Federation Proceedings* 44, no. 2 (1985): 334–352.
- Forbes, Richard M., and John W. Erdman, Jr. "Bioavailability of Trace Mineral Elements." *Annual Reviews of Nutrition* 3 (1983): 213–231.
- Friedman, J.E., et al. "Regulation of Glycogen Resynthesis Following Exercise." *Sports Medicine* 11 (1991): 232.
- Galton, David J., and George A. Bray. "Studies on Lipolysis in Human Adipose Cells." *Journal of Clinical Investigation* 46, no. 4 (1967): 621–629.
- Gao, J.P., D.I. Costill, C.A. Horswill, et al. "Sodium Bicarbonate Ingestion Improves Performance in Interval Swimming." *European Journal of Applied Physiology* 58 (1988): 171–174.
- Garza, C., N.S. Scrimshaw, and V.R. Young. "Human Protein Requirements: The Effect of Variations in Energy Intake within the Maintenance Range." *American Journal of Clinical Nutrition* 29 (1976): 280–287.
- Gastelu, D.L. "Developing State-of-the-Art Amino Acids." *Muscle Magazine International* (May 1989): 58–64.

- . *The Complete Nutritional Supplements Buyer's Guide*. NY: Random House, 2000.
- Gastelu, Daniel, and Fred Hatfield. *Dynamic Nutrition for Maximum Performance*. Garden City Park, NY: Avery, 1997.
- Gattas, V. "Protein-Energy Requirements of Prepubertal School-age Boys Determined by Using Nitrogen-Balance Mixed-Protein Diet." *American Journal of Clinical Nutrition* 52 (1990): 1037-42.
- Gleeson, M., et al. "Effect of Low- and High-Carbohydrate Diets on the Plasma Glutamine and Circulating Leukocyte Responses to Exercise." *International Journal of Sports Nutrition* 8 (1998): 49-59.
- Goldberg, Alfred L., Joseph D. Etlinger, David F. Goldspink, et al. "Mechanism of Work-Induced Hypertrophy of Skeletal Muscle." *Medicine and Science in Sports* 7, no. 3 (1975): 185-198.
- Goldspink, David F. "The Influence of Activity on Muscle Size and Protein Turnover." *Journal of Physiology* 264 (1976): 283-296.
- Gollnick, Philip D. "Metabolism of Substrates: Energy Substrate Metabolism during Exercise and as Modified by Training." *Metabolic and Nutritional Aspects of Physical Exercise: Federation Proceedings* 44, no. 2 (1985): 353-368.
- Gontzea, I., P. Sutzescu, and S. Dumitrache. "The Influence of Muscular Activity on Nitrogen Balance and on the Need of Man for Proteins." *Nutrition Reports International* 10 (1974): 35-43.
- Gotshalk L.A., J.S. Volek, R.S. Staron, et al. "Creatine Supplementation Improves Muscular Performance in Older Men." *Medicine & Science in Sports & Exercise* 34, no. 3 (March 2002): 537-43.
- Green A.L., E. Hultman, I.A. Macdonald, et al. "Carbohydrate Ingestion Augments Skeletal Muscle Creatine Accumulation during Creatine Supplementation in Humans." *Am J Physiol.* 271, no. 5, Pt 1 (1996): E821-6.
- Green A.L., Simpson E.J., Littlewood J.J., et al. "Carbohydrate Ingestion Augments Creatine Retention during Creatine Feeding in Humans." *Acta Physiol Scand.* 158, no. 2 (1996): 195-202.
- Greenhaff, P., et al. "Effect of Oral Creatine Supplementation on Skeletal Muscle Phosphocreatine Resynthesis." *American Journal of Physiology*, 266 (1994): E725-E730.
- Haff, G.G. "Roundtable Discussion: Low Carbohydrate Diets and Anaerobic Athletes." *Strength and Conditioning Journal* 23, no. 3 (2001): 42-61.
- Hagerman, F.C., et al. "Effects of High-Intensity Resistance Training on Untrained Older Men. I. Strength, Cardiovascular, and Metabolic Responses." *Journal of Gerontology Series A Biological Sciences Medical Sciences* 55, no. 7 (2000): B₃36-346.
- Hamilton K.L., M.C. Meyers, W.A. Skelly, et al. "Oral Creatine Supplementation and Upper Extremity Anaerobic Response in Females." *International Journal of Sport Nutrition and Exercise Metabolism* 10, no. 3 (2000): 277-89.

- Haralambie, G., and A. Berg. "Serum Urea and Amino Nitrogen Changes with Exercise Duration." *European Journal of Applied Physiology* (1976): 39–48.
- Hartog, M., R.J. Havel, G. Copinschi, et al. "The Relationship Between Changes in Serum Levels of Growth Hormone and Mobilization of Fat During Exercise in Man." *Quarterly Journal of Experimental Physiology* 52 (1967): 86–96.
- Heeker, A.L., and K.B. Wheeler. "Protein: A Misunderstood Nutrient for the Athlete." *National Strength and Conditioning Association Journal* 7 (1985): 28–29.
- Helie, R., J.-M. Lavoie, and D. Cousineau. "Effects of a 24-Hour Carbohydrate-Poor Diet on Metabolic and Hormonal Responses during Glucose-Infused Leg Exercise." *European Journal of Applied Physiology* 54 (1985): 420–426.
- Henneman, Dorothy, and Philip H. Henneman. "Effects of Human Growth Hormone on Levels of Blood and Urinary Carbohydrate and Fat Metabolites in Man." *Journal of Clinical Investigation* 39 (1960): 1239–1245.
- Hermansen, Lars, Eric Hultman, and Bengt Saltin. "Muscle Glycogen during Prolonged Severe Exercise." *Acta Physiologica Scandinavica* 71 (1967): 129–139.
- Heymsfield, Steven B., Carlos Arteaga, Clifford McManus, et al. "Measurement of Muscle Mass in Humans: Validity of the 24-Hour Urinary Creatinine Method." *American Journal of Clinical Nutrition* 37 (1983): 478–494.
- Hofman, Z., et al. "Glucose and Insulin Responses After Commonly Used Sport Feedings Before and After a 1-Hour Training Session." *International Journal of Sport Nutrition* 5 (1995): 194–205.
- Holloszy, John O. "Adaptation of Skeletal Muscle to Endurance Exercise." *Medicine and Science in Sports* 7, no. 3 (1975): 155–164.
- Hubbard, R., et al. "Apparent skeletal muscle loss related to dietary trans fatty acids in a mixed group of omnivores and vegetarians." *Nutrition Research* 23 (2003): 651–658.
- Institute of Medicine. "Military Strategies for Sustainment of Nutrition and Immune Function in the Field," Robert O. Nesheim, ed. A Report of the Committee on Military Nutrition Research, Food and Nutrition Board. Washington, D.C.: National Academy Press, 1999.
- Izquierdo M., J. Ibanez, J.J. Gonzalez-Badillo, et al. "Effects of Creatine Supplementation on Muscle Power, Endurance, and Sprint Performance." *Medicine & Science in Sports & Exercise* 34, no. 2 (2002): 332–43.
- Jakeman, P., and S. Maxwell. "Effect of Antioxidant Vitamin Supplementation on Muscle Function after Eccentric Exercise." *European Journal of Applied Physiology* 67 (1993): 426.
- James, M.J., R.A. Gibson, L.G. Cleland. "Dietary Polyunsaturated Fatty Acids and Inflammatory Mediator Production." *American Journal of Clinical Nutrition* 71 (suppl) (2002): 3435, 3485.
- Jequier, E. "Thermogenesis Induced by Nutrient Administration in Man." *Infusionsther Klin Ernahr* 11: (1984): 184–189

- Jezova, D., M. Vigas, P. Tatar, et al. "Plasma Testosterone and Catecholamine Responses to Physical Exercise of Different Intensities in Men." *European Journal of Applied Physiology* 54 (1985): 62–66.
- Kanter, M. "Free Radicals, Exercise, and Antioxidant Supplementation." *International Journal of Sports Nutrition* 4 (1994): 205.
- Karagiorgos, Athanase, Joseph F. Garcia, and George A. Brooks. "Growth Hormone Response to Continuous and Intermittent Exercise." *Medicine and Science in Sports* 11, no. 3 (1979): 302–307.
- Karlsson, Jan, and Bengt Saltin. "Lactate, ATP, and CP in Working Muscles during Exhaustive Exercise in Man." *Journal of Applied Physiology* 29, no. 5 (1970): 598–602.
- Karlsson, Jan, Lars-Olof Nordesjo, and Bengt Saltin. "Muscle Glycogen Utilization during Exercise after Physical Training." *Acta Physiologica Scandinavica* 90 (1974): 210–217.
- Kasai, Kikuo, Hitoshi Suzuki, Tsutomu Nakamura, et al. "Glycine Stimulates Growth Hormone Release in Man." *Acta Endocrinologica* 90 (1980): 283–286.
- Kasai, Kikuo, Masami Kobayashi, and Shin-Ichi Shimoda. "Stimulatory Effect of Glycine on Human Growth Hormone Secretion." *Metabolism* 27 (1978): 201–208.
- Kasperek, George J., and Rebecca D. Snider. "Increased Protein Degradation after Eccentric Exercise." *European Journal of Applied Physiology* 54 (1985): 30–34.
- Katch, Victor L., Frank I. Katch, Robert Moffatt, et al. "Muscular Development and Lean Body Weight in Body Builders and Weight Lifters." *Medicine and Science in Sports and Exercise* 12, no. 5 (1980): 340–344.
- Kellis, J.T., and L.E. Vickery. "Inhibition of Estrogen Synthetase (Aromatase) by Flavones." *Science* 225 (1984): 1032–1033.
- Kelly, V.G., and D.G. Jenkins. "Effect of Oral Creatine Supplementation on Near-Maximal Strength and Repeated Sets of High-Intensity Bench Press Exercise." *Journal of Strength and Conditioning Research* 12 (1998): 109–115.
- Kies, C.V., and J.A. Driskell. *Sports Nutrition: Minerals and Electrolytes*. Boca Raton, FL: CRC, 1995.
- Klatz, Ronald, and Robert Goldman. *The New Anti-Aging Revolution*. North Bergen, NJ: Basci Health Publications, Inc., 2003.
- Knopf, R.F., J.W. Conn, S. S. Fajans, et al. "Plasma Growth Hormone Response to Intravenous Administration of Amino Acids." *Journal of Clinical Endocrinology* 25 (1965): 1140–1144.
- Koeslag, J.H. "Post-Exercise Ketosis and the Hormone Response to Exercise: A Review." *Medicine and Science in Sports and Exercise* 14, no. 5 (1982): 327–334.
- Kreider, R. "Which Protein Is Best for Sports Performance?" *Functional Foods and Nutraceuticals* (Sept. 2002).
- Kreider, R.B., "Effects of Creatine Supplementation on Performance and Training Adaptations." *Molecular and Cellular Biochemistry* 244 (1–2) (Feb 2003): 89–94.

- Kreider, R.B., et al. "Effects of Creatine Supplementation on Body Composition, Strength, and Sprint Performance." *Medicine and Science in Sports and Exercise* 30 (1998): 73–82.
- Krieger, D.R., P.A. Daly, A.G. Dulloo, et al. "Ephedrine, Caffeine and Aspirin Promote Weight Loss in Obese Subjects." *Trans. Assoc. Am. Physicians* 103 (1990): 307–312.
- Lemon, P.W. "Effects of Exercise on Dietary Protein Requirements." *International Journal of Sport Nutrition* 8, no. 4 (1998): 426–447.
- . "Beyond the Zone: Protein Needs of Active Individuals." *Journal of the American College Nutrition* 19, no. 5, Suppl. (2000): 513S–521S.
- Lemon, P.W.R., et al. "Protein Requirements and Muscle Mass/Strength Changes During Intensive Training in Novice Bodybuilders." *Journal of Applied Physiology* 73 (1992): 767–775.
- Lemon, P.W., J.M. Berardi, and E.E. Noreen. "The Role of Protein and Amino Acid Supplements in the Athlete's Diet: Does Type or Timing of Ingestion Matter?" *Current Sports Medicine Reports* 1, no. 4 (2002): 214–21.
- Lemon, P.W.R., and J. P. Mullin. "Effect of Initial Muscle Glycogen Levels on Protein Catabolism during Exercise." *The American Physiological Society* (1980): 624–629.
- Lemon, P.W.R., and F.J. Nagle. "Effects of Exercise on Protein and Amino Acid Metabolism." *Medicine and Science in Sports and Exercise* 13, no. 3 (1981): 141–149.
- Lemon, P.W.R., and D. Proctor. "Protein Intake and Athletic Performance." *Sports Medicine* 12, no. 5 (1991): 313.
- Linderman, J., and T.D. Fahey. "Sodium Bicarbonate Ingestion and Exercise Performance." *Sports Medicine* 11, no. 9 (1991): 71.
- Lucke, Christoph, and Seymour Glick. "Experimental Modification of the Sleep-Induced Peak of Growth Hormone Secretion." *Journal of Clinical Endocrinology and Metabolism* 32 (1971): 729–736.
- MacLean, William C., Jr., and George G. Graham. "The Effect of Level of Protein Intake in Isoenergetic Diets on Energy Utilization." *American Journal of Clinical Nutrition* 32(7) (Jul 1979): 1381–1387.
- Manore, M. "Vitamin B₆ and Exercise." *International Journal of Sports Nutrition* 4 (1994): 89.
- Marable, N.L., J.F. Hickson Jr., M.K. Korslund, et al. "Urinary Nitrogen Excretion as Influenced by a Muscle-Building Exercise Program and Protein Intake Variation." *Nutrition Reports International* 19, no. 6 (1979): 795–805.
- Marriott, B. *Food Components to Enhance Performance*. Washington, D.C.: National Academy Press, 1994.
- Maughan, Ronald. "Creatine Supplementation and Exercise Performance." *International Journal of Sport Nutrition* 5(2) (Jun 1995): 94–101.
- Mayer, Jean, Roy Purnima, and Kamakhya Prasad Mitra. "Relation Between Caloric Intake, Body Weight, and Physical Work: Studies in an Industrial Male Population in West Bengal." *American Journal of Clinical Nutrition* 4, no. 2 (1956): 169–175.

- McBride, J.M., et al. "Effect of Resistance Exercise on Free Radical Production." *Medicine and Science in Sports and Exercise* 30 (1998): 67–72.
- Mercola, Joseph. "Reduce Grains and Sugar to Lose Weight and Improve Health." *Cancer Comfort* 18 (July 30, 2003).
- Merimee, T.J., D. Rabinowitz, and S.E. Fineberg. "Arginine-Initiated Release of Human Growth Hormone." *New England Journal of Medicine* 26(26) (Jun 1969): 1434–1438.
- Merimee, Thomas J., David Rabinowitz, Lamar Riggs, et al. "Plasma Growth Hormone after Arginine Infusion." *New England Journal of Medicine* 23 (1967): 434–438.
- Mitchell, J.B., D.L. Costill, J.A. Houmard, et al. "Effects of Carbohydrate Ingestion on Gastric Emptying and Exercise Performance." *Medicine and Science in Sports and Exercise* 20, no. 2 (1988): 110–115.
- Mittleman, K.D., M.R. Ricci, and S.P. Bailey. "Branched-Chain Amino Acids Prolong Exercise During Heat Stress in Men and Women." *Medicine and Science in Sports and Exercise* 30 (1998): 83–91.
- Mosoni L., and P.P. Mirand. "Type and Timing of Protein Feeding to Optimize Anabolism." *Current Opinions in Clinical Nutrition and Metabolic Care* 6, no. 3 (2003): 301–306.
- Murphy, T., et al. "Performance Enhancing Ration Components Project: U.S. Army." Abstract. Presented at the 11th Annual Symposium of Sports and Cardiovascular Nutritionists. Atlanta, Georgia (April 1994): 22–24.
- Pavlou, Konstantin N., William P. Steffee, Robert H. Lerman, et al. "Effects of Dieting and Exercise on Lean Body Mass, Oxygen Uptake, and Strength." *Medicine and Science in Sports and Exercise* 17, no. 4 (1974): 466–471.
- Piehl, Karin. "Time Course for Refilling of Glycogen Stores in Human Muscle Fibres Following Exercise-Induced Glycogen Depletion." *Acta Physiologica Scandinavica* 90 (1974): 297–302.
- Pizza, F., et al. "A Carbohydrate Loading Regimen Improves High Intensity, Short Duration Exercise Performance." *International Journal of Sport Science* (1995): 110–116.
- Prasad, Ananda S. "Role of Trace Elements in Growth and Development." *Nutrition Research* (1985): 295–299.
- Preuss H.G., D. DiFerdinando, M. Bagchi, D., et al. "Citrus Aurantium as a Thermogenic, Weight-Reduction Replacement for Ephedra: An Overview." *Journal of Medicine* 33, nos. 1–4 (2002): 247–64.
- Rasmussen, B.B., K.D. Tipton. "An Oral Essential Amino Acid-Carbohydrate Supplement Enhances Muscle Protein Anabolism After Exercise." *Journal of Applied Physiology* 88, No. 2 (Feb 2000): 386–392.
- Rawson, E.S., and J.S. Volek. "Effects of Creatine Supplementation and Resistance Training on Muscle Strength and Weightlifting Performance." *Journal of Strength and Conditioning Research* 17, no. 4 (2003): 822–31.

- Rossi, A., R.A. DiSilvestro, A. Blostein-Fujii. "Effects of Soy Consumption on Exercise-Induced Acute Muscle Damage and Oxidative Stress in Young Adult Males." *FASEB Journal* (1998): A653.
- Rudman, D., et al. "Effects of Human Growth Hormone in Men over 60 Years Old." *New England Journal of Medicine* 323 (July 5, 1990): 1–6.
- Satabin, Pascale, Pierre Portero, Gilles Defer, et al. "Metabolic and Hormonal Responses to Lipid and Carbohydrate Diets during Exercise in Man." *Medicine and Science in Sports and Exercise* 19, no. 3 (1987): 218–223.
- Schalch, Don S. "The Influence of Physical Stress and Exercise on Growth Hormone and Insulin Secretion in Man." *Journal of Laboratory and Clinical Medicine* 69, no. 2 (1967): 256–267.
- Schwartz, J.M., Y. Schulz, V. Piolino, et al. "Thermogenesis in Men and Women Induced by Fructose vs. Glucose Added to a Meal." *American Journal of Clinical Nutrition* 49 (1989): 667–674.
- Spiller, G.A., C.D. Jensen, T.S. Pattison, et al. "Effect of Protein Dose on Serum Glucose and Insulin Response to Sugars." *American Journal of Clinical Nutrition* 46 (1987): 474–480.
- Tarnopolsky, M.A., and D.P. MacLennan. "Creatine Monohydrate Supplementation Enhances High-Intensity Exercise Performance in Males and Females." *International Journal of Sport Nutrition and Exercise Metabolism* 10, no. 4 (2000): 452–63.
- Tesch, Per, et al. "Skeletal Muscle Glycogen Loss Evoked by Resistance Exercise." *Journal of Strength and Conditioning Research* 12 (1998): 67–73.
- Thomas, D., et al. "Plasma Glucose Levels after Prolonged Strenuous Exercise Correlate Inversely with Glycemic Response to Food Consumed before Exercise." *International Journal of Sport Nutrition* 4 (1994): 361.
- Tipton, K.D., and Wolfe, R.R. "Exercise, Protein Metabolism, and Muscle Growth." *International Journal of Sport Nutrition and Exercise Metabolism* 11 (2001): 109–132.
- Todd, Karen S., Gail E. Butterfield, and Doris Howes Calloway. "Nitrogen Balance in Men with Adequate and Deficient Energy Intake at Three Levels of Work." *Journal of Nutrition* 114 (1984): 2107–2118.
- Tokmakidis, S.P. "The Effects of Ibuprofen on Delayed Muscle Soreness and Muscle Performance after Eccentric Exercise." *Journal of Strength and Conditioning Research*, 17 (February 2003): 53–59.
- Tsintzas, K. and C. Williams. "Human Muscle Glycogen Metabolism during Exercise. Effect of Carbohydrate Supplementation." *Sports Medicine* 25 (1998): 7–23.
- Valeriani, A. "The Need for Carbohydrate Intake During Endurance Exercise." *Sports Medicine* 12, no. 6 (1991): 349.
- Walberg, Janet L., V. Karina Ruiz, Sandra L. Tarlton, et al. "Exercise Capacity and Nitrogen Loss during a High or Low Carbohydrate Diet." *Medicine and Science in Sports and Exercise* 20 (1986): 34–43.

- Warber, J.P., W.J. Tharion, J.F. Patton, et al. "The Effect of Creatine Monohydrate Supplementation on Obstacle Course and Multiple Bench Press Performance." *Journal of Strength and Conditioning Research* 16, no. 4 (2002): 500–8.
- Ward, P.S., and D.C.L. Savage. "Growth Hormone Responses to Sleep, Insulin Hypoglycemia and Arginine Infusion." *Hormone Research* 22 (1985): 7–11.
- Welbourne, I.C. "Increases in Plasma Bicarbonate and Growth Hormone After an Oral Glutamin Load." *American Journal of Clinication Nutrition* 61 (1995): 1058–1061.
- Weltman, Arthur, Sharleen Matter, and Bryant A. Stamford. "Caloric Restriction and/or Mild Exercise: Effects on Serum Lipids and Body Composition." *American Journal of Clinical Nutrition* 33 (1980): 1002–1009.
- West, D. "Reduced Body Fat With Conjugated Linoleic Acid Feeding in the Mouse." *Federation of American Societies of Experimental Biology Journal* 11 (1997): A599.
- Wilcox, Anthony R. "The Effects of Caffeine and Exercise on Body Weight, Fat-Pad Weight, and Fat-Cell Size." *Medicine and Science in Sports and Exercise* 14 (1981): 317–321.
- Williams, M.H. "Vitamin Supplementation and Athletic Performance." *International Journal of Vitamin and Nutrition Research* 30 (1989): 163.
- Wolfe, R.R. "Protein Supplements and Exercise." *American Journal of Clinical Nutrition* 72 Suppl. (2000): 551S–557S.
- Wolinsky, I., and J. Hickson. *Nutrition in Exercise and Sport*. 2nd ed. Boca Raton, FL: CRC Press, 1994.
- Wright, J. "Tribulus: A Natural Wonder." *Muscle and Fitness* 224 (September 1996): 140–142.
- Yan, W., et al. "Steroidal Saponins from Fruits of Tribulus Terrestris." *Phytochemistry* 42, no. 5 (1996): 1417–1422.
- Young, K., and C.T.M. Davies. "Effect of Diet on Human Muscle Weakness Following Prolonged Exercise." *European Journal of Applied Physiology* 53 (1984): 81–85.
- Young, Vernon R., and Peter L. Pellett. "Protein Intake and Requirements with Reference to Diet and Health." *American Journal of Clinical Nutrition* 45 (1987): 1323–1343.
- Zawadzki, K.M., B.B. Yaspelkis, and J.L. Ivy. "Carbohydrate-Protein Complex Increases the Rate of Muscle Glycogen Storage after Exercise." *Journal of Applied Physiology* 72 (1992): 1854–1859.

Macrobiotic Nutrition Chart References

- Y. Boirie, M. Dangin, P. Gachon, et al., "Slow and Fast Dietary Proteins Differently Modulate Postprandial Protein Accretion," *Proceedings of the National Academy of Science U.S.A.* 94 (1997): 14930–14935.
- Céline Morens, Cécile Bos, Maria E. Pueyo, et al., "Increasing Habitual Protein Intake Accentuates Differences in Postprandial Dietary Nitrogen Utilization between Protein Sources in Humans," *Journal of Nutrition* 133 (2003): 2733–2740.

Index

- AAKG, 37
Adaptogen, 146
Adenosine triphosphate. *See* ATP.
ADNO, 124
Adrenal glands, 146
Aerobic exercise, 80–81
Alanine, 13
Alpha linolenic acid, 62–63
American Academy of Anti-Aging
 Medicine, 133
American Journal of Clinical Nutrition, 21,
 35, 76
Amino acids, 13, 22, 28, 31–37, 52–57,
 59, 72–73, 74, 80
 branched-chain, 37–38, 43, 46, 48,
 141
 essential, 78
 profiles of protein sources, 51,
 52–53, 55, 59
Ammonia, 34, 37
Amylase, 71
Anabolic hormones, 35
Anabolic phase, 30, 52–54, 78
Anabolic steroid alternatives, 128–131
Anaerobic exercise. *See* Weight training.
Anderson, J.W., 50
Angle, Kurt, 143
Antioxidants, 34, 97–98, 115, 148
Arginine, 33, 36–38, 124–126, 141
Arginine alpha ketoglutarate. *See*
 AAKG.
Arginine-derived nitric oxide. *See*
 ADNO.
Aromatase, 129
Ascorbic acid. *See* Vitamin C.
Aspirin, 137–138, 144–145
Atkins, Robert, 6
Atkins Diet, 6, 66
ATP, 36, 122, 125–126

B cells, 65
Banaba, 143
Bars, 171
Basal metabolic rate. *See* BMR.
Baylor University, 122
BCAAs. *See* Amino acids, branched-
 chain.
Bedtime, 148
Beef, 41–42, 58
 protein/nutritional value of, 42
Beef mass burger, 161
Bennett, Chris “Big Guns,” 4
Bile, 72
Biological value. *See* BV.
Biotin, 94–95
Bitter orange, 145
Blood assays, 137–138
Blood flow, 36
Blood sugar, 6, 15, 17, 22–26, 72, 147
Bloomsburg University, Department of
 Exercise Science and Athletics, 121
BMR, 75–79, 83–84

- Body fat, 3, 80
Body water weight. *See* BWW.
Body weight. *See* BW.
Bodybuilding, 1–2, 3–11, 12–27, 28–60,
61–67, 68–70, 71–82, 83–91,
92–118, 119–149, 150–173, 175,
176–188, 189–190, 191–194
Bohmer, D., 138
Boirie, Yves, 48, 57
Bone growth, 105–108, 117
Boron, 118
Braatan, J.T., 21
Brand-Miller, Jennie, 15
Breakfast, 9–11, 151–157
British Journal of Nutrition, 64
BV, 31–32, 43, 47–48
BW, 84–87
BWW, 70

Caffeine, 144–145
Calcium, 106–108, 143, 146
Calcium caseinate. *See* Casein.
Caloric requirements table, 8, 29,
86–90, 176–188
Calories, 2, 4–6, 9–11, 14–15, 22, 61,
75, 79, 80, 83–91, 150–173,
176–188, 189–190
converting into grams, 89
distribution, 88
Calver, A., 36
Carbohydrates, 2, 3–4, 6–8, 12–27, 71,
74–78, 92–118, 150–173
disaccharides, 15
low, 13–14
low-glycemic, 22, 54
monosaccharides, 15
polysaccharides, 15
recommended sources, 16
Cardio exercise, 3, 79–81, 86–87
Cardiovascular health, 79
Casein, 40, 48–49, 55, 57–58
product analysis, 49
Caseinate. *See* Casein.
Catabolic hormone, 34
Catabolic state, 31, 80
Cations, 68

CCK, 66, 73
Celebrex, 135
Cell volumization, 35
Chandalia, M., 21
Chemical score, 31
Chicken breast, 42
protein/nutritional value of, 42
Chicken parm pasta salad, 158
Chloride, 104–105
Cholecystokinin, 66
Cholesterol, 65
Chromium, 117–118, 143
Citrus aurantium. *See* Bitter orange.
Claridge, Travis, 30
Clinical Science, 36
Coagulants, 101
Cobalt, 118
Coenzyme nicotinamide adenine
dinucleotide. *See* NADH.
Collagen, 139
Copper, 115–116
Cortisol, 21, 25–27, 34
COX-1 enzyme, 135–136
COX-2 enzyme, 63, 135–138
COX-2 inhibitors, 135–138
CP, 36, 122
Creatine, 121–126
supplementation results, 121
Time Release Technology, 123–126
Time Released Arginine/Creatine,
124–126
transport, 123
Creatine monohydrate, 123
Creatine phosphate. *See* CP.
Creatine phosphokinase. *See* CPK.
Creatinine, 123
Curcumin, 137
Cyanocobalamin. *See* Vitamin B₁₂.
Cyclooxygenases. *See* COX-2.

Dairy, 107
low-glycemic, 193
Delayed-onset muscle soreness. *See*
DOMS.
Dehydration, 70
Demoritus University, Greece, 135

- Desserts, 172–173
DHA, 63
DHT, 129–130
Diabetic Medicine, 21
Diet goals, 8–9
Digestion, 39–40, 71–73
Digestive enzymes, 72
Dillingham, B.L., 51
Dinner, 164–170
Docosahexaenoic acid. *See* DHA.
DOMS, 134
Dullo, A.G., 146
DuPont Protein Technologies, 50
- ECA, 144–145
EFAs, 61–62, 74
Egg protein powder, 49
Egg white omelet with macrobiotic granola, 152
Eggs, 43–44, 49
 protein/nutritional value of, 43
 whites, 43
Eicosanoids, 65
Eicosapentaenoic acid. *See* EPA.
Electrolytes, 68
Endocrine system, 134
Endocrinology and Metabolism, 76
Endurance conditioning, 80
Energy, 141
EPA, 63
Ephedrine/ephedra, 144–147
Ephedrine, caffeine and aspirin combination. *See* ECA.
Esophagus, 71
Essential fatty acids. *See* EFAs.
Estrogen, 50, 129–130
European Journal of Clinical Nutrition, 76
Exercise expenditure, 86–87
Fat, 2, 5–8, 61–67, 71–72, 74–77, 92–118, 150–173, 194
 facts on, 66–67
 hydrogenated, 66
 monounsaturated, 61, 67
 polyunsaturated, 61
 saturated, 61–62, 64–65, 66
 trans, 62, 66
 unsaturated, 61–62, 64, 66
Fat loss, 142–147
Fat storage, 20, 22
Fatty acids, 33
FDA, 32, 50, 145
Fiber, 21, 72
 insoluble, 21
 soluble, 21
Fish, 44–45
 oil, 63, 65
 protein/nutritional value of, 44
5-dihydrotestosterone. *See* DHT.
Fluoride, 117
Folic acid, 96
Food, 1–2
 nutrient content, 41
 thermic effects, 39–40, 75–77
Food and Agriculture Organization, 32
Food pyramids, 189–190
45/35/20 lean mass equation, 2, 4–8, 10–11, 18–20, 25, 27, 52, 54, 59, 71, 73, 83, 87–88, 141, 142, 147, 150, 175
Free radicals, 50, 148
Fruit salad, 172
Fruits, low-glycemic, 192
Functional Foods and Nutraceuticals, 49
- GABA, 126
GAGs, 139
Gastrin, 73
Gatas, V., 49
Georgetown University, 146
GH. *See* Growth hormone.
GI. *See* Glycemic index.
Giampapa, Vincent, 131–134
Ginkgo, 147
Ginseng, 146–147
Glucagon, 17–18, 25–27, 54, 57, 73, 74, 78
Gluconeogenesis, 6, 13
Glucosamine, 138–140
Glucose, 12, 17, 22, 33, 72, 123
Glutamine, 13, 33–36, 38, 48, 126–127, 141
 effervescent, 38, 127

- Glycemic index, 15, 17–20
 insulin and, 17
 table of foods, 19–20
- Glycemix LGI, 141
- Glycogen, 6–7, 12, 18, 37, 63, 80
 replenishment, 12–13, 22
- Glycosaminoglycans. *See* GAGs.
- Goldman, Bob, 133
- Grains, low-glycemic, 192
- Green, A.L., 123
- Green tea, 145–146
- Grilled chicken-bean salad, 159
- Growth hormone, 21, 25–27, 35–36,
 54, 57, 74, 127
- Guarana, 146
- Guide to estimating calories, 2
- Hawk, Dave, 14
- HCL acid, 127
- Heart rate, target, 81
- Herbs, 145
- HGH. *See* Growth hormone.
- Histidine, 33
- Hubbard, R., 62
- Human growth hormone. *See* Growth hormone.
- Hydration, 68–70
- Hypertension, 104
- Hypoglycemia, 17–18, 22–24
- Ibuprofen, 135–138
- IGE, 66
- IGF-1, 125, 132–133
- Immune globulin E. *See* IGE.
- Immune system, 34, 37, 65–66
- Institute of Medicine, 126
- Insulin, 17–18, 20, 22–27, 54, 57, 63,
 72–73, 74, 117
 anabolic actions of, 20, 22–23, 25
 controlling, 22–23
 hyperlipidemic actions of, 20,
 22–23, 25
 negative effects of, 23–25
 resistance, 20, 22, 24–25, 63
- Intestine, small, 72
- Interstitial fluid, 69
- Intracellular fluid, 69
- Iodine, 114–115
- Iron, 110–112
- IRS. *See* Insulin, resistance.
- Isoflavones, 50–51
- Isoleucine, 33, 37
- IsoOxygene, 136–138
- James, M.J., 63
- Jequier, E., 77
- Joint pain and inflammation, 63, 134–140
- Journal of Applied Physiology*, 78
- Journal of Biological Chemistry*, 63
- Journal of Medicine*, 145
- Journal of Strength and Conditioning Research*, 121, 135
- Ketone bodies, 33
- Ketosis, 6, 13
- Kirit, Steve, 8
- Klatz, Ron, 133
- Kreider, R., 49, 122
- Krieger, Diane R., 144
- Lactic acid, 34, 69
- L-arginine, 124–126
 dose response curve for oral, 124
- Lean-mass equation, 2, 3, 9–11, 52, 83
- Lemon salmon with spinach, 167
- Lemon sole with broccoli, 165–166
- Leptin, 45
- Leucine, 33, 37
- L-glutamine, 127
- Lifestyle, 85–87
- Linoleic acid, 62
- Lipid Complex, 142
- Lipids, 142
- Liver, 6, 37, 72
- Low-carb diets. *See* Carbohydrates, low.
- L-tyrosine, 147
- Lunch, 158–163
- Lysine, 33
- Macrobiotic caloric equation, 84–87
- Macrobiotic caloric requirements tables,
 176–188

- Macrobiotic exchange lists, 191–194
- Macrobiotic food guide pyramids, 189–190, 191
- Macrobiotic granola, 155–157
- Macrobiotic granola mix with side of eggs, 152–153
- Macrobiotic instant chocolate pudding, 173
- Macrobiotic morning mix, 153–154
- Macrobiotic Nutrition, 2, 3–11, 12–27, 28–60, 61–67, 68–70, 71–82, 83–91, 92–118, 119–149, 150–173, 175, 176–188, 189–190, 191–194
- enhancing the effect, 119–149
- hormonal effects on body composition, 27
- meals, 150–174
- Macrobiotic pancakes, 154–155
- Macrobiotic sushi menu, 168–170
- Macrobiotic vs high-glycemic meals, 23
- Macrominerals, 103–109
- Macrophages, 65
- Magnesium, 108–109, 143
- Manganese, 116–117
- Maximum Human Performance, Inc.
 See MHP.
- Meal planning, 191–194
- Meal timing, 52, 59, 88
- Meal-replacement powders. *See* MRPs.
- Meals, 71–82, 150–174
- Meat, 193
- Meat substitutes, 193
- Medicina dello Sport*, 38
- Mercola, Joseph, 24
- Metabolic goal variable. *See* MGV.
- Metabolism, 79
- Methionine, 33
- Mexican omelet with macrobiotic granola, 151
- MGV, 84–87
- MHP, 58, 120, 123–124, 127–129, 131, 134, 136, 141, 145–146, 148–149
- Microminerals, 109–118
- Micronutrients, 92–118
- Minerals, 101–118
- Mixed-meat teriyaki, 164–165
- Molybdenum, 118
- Montclair State University, 150
- Morris, Mike, 5
- MRPs, 140–142
- Muscle growth, 20, 22, 30–38, 62, 79–80
- Muscle loss, 6
- Muscle mass, 71–82
- NADH, 125–126
- Naproxen, 137
- Net protein utilization. *See* NPU.
- New England Journal of Medicine*, 21, 50, 132
- Nexrutine, 137
- Niacin. *See* Vitamin B₃.
- Nickel, 118
- Nighttime lean muscle-building requirements, 147–148
- Nitric oxide. *See* NO.
- Nitro-loading, 124
- Nitrogen, 30–31, 55, 57
- positive balance, 32–33, 39–40, 52–53, 57
- NO, 124
- No-carb diets, 6
- Nolvaldex, 50
- Nonsteroidal anti-inflammatory drugs. *See* NSAIDs.
- NPU, 31
- NSAIDs, 135–140
- Nutrition Research, 62
- Ohio State University, 50
- Omega-3 fatty acids, 49, 62–63, 65, 67
- Omega-6 fatty acids, 62, 67
- 1-Testosterone, 128
- Pancreas, 25, 72
- Pantothenic acid, 95
- PDCAAS, 31–32, 43, 47, 50
- Pepsin, 39
- Peptides, 39
- PER, 31–32
- Performance enhancement, 4, 69
- Peristalsis, 71

- PGE2, 137
PGs, 139
Phenylalanine, 33
Philippi, Mark, 9
Physique, critiquing, 90–91
Phosphate, 69
Phosphorus, 108
Piers, L.S., 64
Pituitary gland, 134
Pork tenderloin, 45
 protein/nutritional value of, 45
Potassium, 68–69, 105
Preuss, H.G., 145
Probiotic protein blend, 51, 54, 57–58,
 60, 141, 148
Probiotic SR, 148
*Proceedings of the National Academy of
 Sciences*, 48
Progress as motivation, 175
Prostaglandin, 137
Protein digestibility corrected amino
 acid score. *See* PDCAAs.
Protein efficiency ratio. *See* PER.
Protein, 2, 4, 6–8, 12–14, 28–60, 71–72,
 75–77, 92–118, 150–173
 absorption rates of, 54–56
 complex, 39
 impact on training, 59–60
 probiotic vs leading proteins, 55
 release rates of sources, 54–57
 sources, 33, 38–60, 61
 sources grading, 31–32
 supplements, 40, 46–52, 57–59
Proteoglycans. *See* PGs.
Pyridoxine. *See* Vitamin B₆.

Rasmussen, B.B., 78
Rawson, E.S., 121
Receptor cells, 130
References, 195–206
RELEVE, 139–140
Resistance training. *See* Weight training.
Resveratrol, 137
Riboflavin. *See* Vitamin B₂.
Rofecoxib, 138
Rossi, A., 50

Royal London School of Medicine, 137
Rudman, Daniel, 132–133
Rychlak, Gene, 24

Salon, 44
Schwartz, J.M., 75–76
Scrambled eggs with oatmeal and
 strawberries, 157
Sears, Barry, 6
Second messengers, 130–131
Secretin, 73
Secretagogue-One, 134
Selenium, 115
Sex-hormone-binding globulin. *See*
 SHBG.
Shakes, 171
SHBG, 129–130
Signal transduction, 130–131
Silicon, 118
Skeletal muscle, 37
Sodium, 68, 103–104
Solae Company, 49–50
Somostatin, 36, 54, 57
Soy, 40, 49–52, 55, 58
Soy protein powder, 49–54
Steak pita pocket, 162
Steroid hormones, 65
Stomach, 71–72
Strength and Conditioning Journal, 12
Strydom, Gary, 13
Sulfur, 109
Supplement sources, 46–54
Supplements, 194
Supplements vs whole foods, 38–39
Supro powder, 49–50
Syndrome X. *See* Insulin, resistance.
Sydney University Human Nutrition
 Unit, 15

T cells, 65
TakeOFF, 146–147
T-BOMB, 128–129
T-BOMB II, 129–130
TEF, 74–77, 142–144
Testimonials, 4–5, 8–9, 13–14, 24, 30,
 143

- Testosterone, 51, 65, 127–131
 optimizers, 128–131
 T-BOMB, 128–129
 T-BOMB II, 129–131
- Thermic effects of food. *See* TEF.
- Thermogenic fat-loss aids, 142–147
- Thiamine. *See* Vitamin B₁.
- Threonine, 33
- Time Release Technology. *See* TRT.
- Time Released Arginine/Creatine.
 See TRAC.
- Tocopherols, 100
- Tokmakidis, S.P., 135
- Tomoxifin citrate, 50
- TRAC, 124–126
 sustained release profile, 125
- Transactions of the Association of
 American Physicians, 144
- TRT, 123–126
- Tryptophan, 33
- Tuna, 44
- Tuna salad sandwich, 160
- University of Guelph, 51
- Up Your MASS bar, 171
- Up Your MASS nutrition products,
 119, 140–143, 148
- Up Your MASS shake, 171
- U.S. Food and Drug Administration.
 See FDA.
- U.S. Olympic Committee, 6
- Valine, 33, 37
- Vanadium, 118
- Vegetables, low-glycemic, 191–192
- Vegetables, green leafy, 63
- Viox, 135
- Vitamin A, 61, 98–99
- Vitamin B complex, 92–97
 anabolic, 95–96
- Vitamin B₁, 93
- Vitamin B₂, 93–94
- Vitamin B₃, 94
- Vitamin B₆, 95–96
- Vitamin B₁₂, 96–97
- Vitamin C, 97–98, 148
- Vitamin D, 61, 65, 99–100
- Vitamin E, 61, 100–101
- Vitamin K, 61, 101
- Vitamins, 92–101
 fat-soluble, 98–101
 water-soluble, 92–98
- Volek, J.S., 121
- Warn, Roger, 150
- Water, 68–70
 dosages, 70
 from food, 70
 impact on performance, 69
- Weight gain, 5
- Weight loss MRPs, 140–141
- Weight training, 2, 3–11, 12–27, 28–60,
 61–67, 68–70, 71–82, 83–91,
 92–118, 119–149, 150–173, 175,
 176–188, 189–190, 191–194
- Welbourne, Thomas, 35
- Whey, 40, 46–48, 55, 57–58
 grades, 46–47
 product analysis, 47
- Whey protein concentrate, 46–48
- Whey protein isolate, 46–48
- WHMA, 137
- WHO, 32
- Whole blood assays, 137–138
- Whole foods vs supplements, 38–39
- World Health Organization. *See*
 WHO.
- Wound healing, 37
- WPC. *See* Whey protein concentrate.
- WPI. *See* Whey protein isolate.
- Zinc, 113–114
- Zone diet, 6–7

About the Authors



Gerard Dente has over fifteen years of experience in bodybuilding and nutrition. He is a former top-ranked national bodybuilder who has now dedicated his life to sports enhancement through nutrition and supplementation. He is the founder and president of one of the leading sports nutrition companies, Maximum Human Performance. He also serves as a nutritional consultant to many professional athletes and bodybuilders.



Kevin Hopkins attended Rutgers University College of Pharmacy and took a position with one of the nation's largest compounding pharmacies. Shortly thereafter, he started a business that integrated the use of nutritional supplements with prescription medications and nutrition. He also works with MHP's research and development team in designing cutting-edge supplements. He owns Gibson's Gym in Washington, New Jersey, and is continuing to compete recreationally as a bodybuilder in the NPC.