

**NASA S-NPP VIIRS Snow Cover
Products
Collection 1 (C1)
User Guide**

Release 1.1

**George A. Riggs
Dorothy K. Hall
Miguel O. Román**

April 2019

Table of Contents

1.0 Overview	4
2.0 NASA VIIRS Snow Cover Data Products	5
3.0 VNP10	6
3.1 Geolocation Data	7
3.2 SnowData Group	7
3.2.1 Variables	8
3.2.1.1 NDSI_Snow_Cover	8
3.2.1.2 Algorithm_bit_flags_QA	10
3.2.1.3 Basic_QA	11
3.2.1.4 NDSI	12
3.3 Snow Cover Detection Algorithm Synopsis	12
3.3.1 Data Screens	14
3.3.1.1 Low VIS reflectance screen	14
3.3.1.2 Low NDSI screen	15
3.3.1.3 Estimated surface temperature and surface height screen.	15
3.3.1.4 High SWIR reflectance screen.	15
3.3.1.5 Solar zenith screen.	15
3.3.2 Lake Ice Algorithm	15
3.3.3 Cloud Masking	16
3.3.4. Quality Assessment (QA)	16
3.4 Interpretation of Snow Cover Detection Accuracy, Uncertainty and Errors	16
3.4.1 Warm surfaces	17
3.4.2 Low reflectance	19
3.4.3 Low NDSI	20
3.4.4 High SWIR reflectance	20
3.4.5 Cloud and snow confusion	20
3.4.6 Lake ice	22
3.4.7 Bright surface features	22
3.4.8 Land/water mask	22
3.4.9 Geolocation accuracy	23
3.4.10 Antarctica	23
4.0 VNP10A1	23
4.1 Variables	24
4.1.1 NDSI_Snow_Cover	24
4.1.2 Basic_QA	25
4.1.3 Algorithm_bit_flags_QA	25
4.1.4 NDSI	25
4.1.5 granule_pnt	26
4.1.6 Projection	26
4.2 Interpretation of Snow Cover Detection Accuracy, Uncertainty and Errors	26
5.0 VNP10A1F	26
5.1 Algorithm Description	27

5.2 Variables	28
5.2.1 CGF_NDSI_Snow_Cover	28
5.2.2 Basic_QA	29
5.2.3 Algorithm_bit_flags_QA	29
5.2.4 Projection	29
5.2.5 VNP10A1_NDSI_Snow_Cover	29
5.3 Interpretation of Snow Cover Accuracy, Uncertainty and Errors	29
6.0 Related Web Sites	31
7.0 References	32
8.0 List of Acronyms	33
9.0 Appendix A VNP10 Global Attributes	34
10.0 Appendix B VNP10A1 Contents	36
11.0 Appendix C VNP10A1F Contents	40

1.0 Overview

The NASA Suomi-National Polar-orbiting Partnership (S-NPP) Visible Infrared Imaging Radiometer Suite (VIIRS) snow cover algorithm and data product is developed synergistically with the Moderate Resolution Imaging Spectroradiometer (MODIS) Collection 6.1 (C6.1) snow cover algorithms, leveraging analysis and evaluation from both to make nearly-identical algorithms and similar data products. The overall objective for VIIRS Collection 1 (C1) is to make the NASA VIIRS snow cover mapping algorithms compatible with the C6.1 MODIS snow cover algorithms for eventual development of a climate-data record (CDR) using products from the two sensors. Differences between the MODIS C6.1 and the NASA VIIRS algorithms originate from the physical differences between the MODIS and VIIRS sensors, including spatial resolution and band location and width, and the respective cloud masks that are input to the algorithms. The NASA VIIRS snow cover algorithms and data products in C1 have been significantly revised and data content has been increased compared to the original NOAA-Interface Data Processing Segment (IDPS) snow cover products that were developed based on the MODIS Collection 5 (C5) snow cover algorithms.

Snow cover is detected using the Normalized Difference Snow Index (NDSI) snow cover. As in MODIS C6.1 snow products, both a NDSI snow cover map with masks of clouds, night and oceans are applied and the NDSI calculated for all underlying land pixels in the products. The estimate of Fractional Snow Cover (FSC) that was done in MODIS C5 is not made in the algorithm. The NDSI is related to the presence of snow in a pixel and is a more accurate description of the snow detection as compared to estimating FSC based on empirical relationships and allows a user greater flexibility in interpreting data. A detailed explanation for the change from FSC to NDSI snow cover is given in the NASA VIIRS snow cover ATBD [\[http://npp.gsfc.nasa.gov/documents.html\]](http://npp.gsfc.nasa.gov/documents.html). Should a user want to estimate FSC they could develop their own relationship between NDSI and FSC for a given study area.

The VIIRS snow cover data products are produced in different file formats depending on product processing level. The swath Level-2 (L2) VNP10 product is in HDF5 and uses netCDF Climate Forecast (CF-1.6) conventions for some global and local attributes and for georeference of variables. The Level-3 (L3) products are gridded and projected to the sinusoidal projection, which is the same grid and projection as MODIS products but at the VIIRS nominal spatial resolution of 375 m, and in HDF-EOS5 format with the addition of CF-1.6 conventions for global and local attributes and for georeference of variables. Information on file formats can be found at: netCDF <https://www.unidata.ucar.edu/software/netcdf/docs/index.html>, CF-1.6 <http://cfconventions.org>, Hierarchical Data Format 5 (HDF5) <https://www.hdfgroup.org/HDF5/> and HDF-EOS5 <https://earthdata.nasa.gov/user-resources/standards-and-references/hdf-eos5>. Two changes from the MODIS snow cover products are that the VNP10 Level-2 products consist of 6-minute swaths, MODIS

swaths are 5 minutes, and the products are in different formats; the MODIS product are in HDF-EOS4.

This User Guide describes each of the NASA VIIRS C1 snow cover products in sequence from Level-2 to Level-3. The VIIRS snow products are referenced by their Earth Science Data Type (ESDT) name, e.g., VNP10* in this guide (the asterisk refers to all of the NASA VNP snow products, not a specific product). The ESDTs are produced as a series of products in which the snow detection algorithm results in the L2 product are propagated to the higher level products where gridding, projection, compositing and other algorithms are applied to produce the L3 products. Summaries of the algorithms, data products content, and commentary on evaluation and interpretation of data are given for each product. The reader is referred to the VIIRS Algorithm Theoretical Basis Document (ATBD) [<https://npp.gsfc.nasa.gov/documents.html>] (Riggs et al., 2016) for further details.

Note: The User Guide is developed in increments for each product as they are scheduled to be released so check that you have the latest version of the guide.

2.0 NASA VIIRS Snow Cover Data Products

The NASA VIIRS land snow cover data products are listed in Table 1. Snow cover data products are produced in sequence beginning with a swath at a nominal pixel spatial resolution of 375 m with nominal swath coverage of 6400 pixels (across track) by 6464 pixels (along track), consisting of 6 minutes of VIIRS instrument scans. Products in EOSDIS are labeled with ESDT name, e.g. VNP10*, in which the asterisk refers to all three of the NASA VIIRS snow cover products. The ESDT name is used to identify the snow data products. The ESDT also indicates the spatial and temporal processing that has been applied to the data product. Data product levels briefly described are: Level-1B (L1B) is a swath (scene) of VIIRS data geolocated to latitude and longitude. A Level-2 (L2) product is a geophysical product that remains in the latitude and longitude orientation of L1B. A Level-2 gridded (L2G) product is in a gridded format of the sinusoidal projection for VIIRS snow products. At L2G the data products are referred to as tiles, each tile being 10° x 10° area of the global map projection. L2 data products are gridded into L2G tiles by mapping the L2 pixels into cells of a tile in the map projection grid. The L2G algorithm creates a gridded product necessary for the Level-3 (L3) products. An L3 product is a geophysical product that has been temporally and or spatially manipulated, and is in a gridded map projection format and comes as a tile of the global grid. The VIIRS L3 snow products (VNP10A1 and VNP10A1F) are in the sinusoidal projection at 375 m spatial resolution.

The VNP10 snow cover product is in HDF5 format and netCDF Climate and Forecast Metadata Conventions, Version 1.6, 5 December 2011 (CF-1.6) have been applied to relevant attributes and variables.

The series of NASA VIIRS snow cover products to be produced in C1 is listed in Table 1. Description of each product, synopsis of the algorithm and commentary on snow cover detection, QA, accuracy and errors is given in following sections.

Global attributes describing the time of acquisition of the swath, geographic location of swath, production of the data product, provenance and DOI of the product are attached to the root group; those attributes are listed in Appendix A.

Table 1: Summary of land snow cover products produced at the Land Science Investigator-led Processing System (LSIPS).

Products	ESDT	Description
Snow Cover (L2 Daily Swath product)	VNP10	VIIRS/NPP Snow Cover 6-Min Swath 375 m (HDF5)
Snow Cover (L3 Daily Tiled product)	VNP10A1	VIIRS/NPP Snow Cover Map Daily L3 Global 375 m SIN Grid Day (HDF-EOS5 with CF-1.6 conventions)
Snow Cover (L3 Daily CGF Product)	VNP10A1F	VIIRS/NPP CGF Snow Cover Map Daily L3 Global 375 m SIN Grid Day (HDF-EOS5 with CF-1.6 conventions)

3.0 VNP10

The NASA VIIRS snow cover swath product, VNP10, contains dimensions, a SnowData group of variables and a GeolocationData group of variables. A file level description is given in List 1 and the data groups, variables and attributes are described in following sections.

List 1. File level description of the contents of the VNP10 product.

dimensions:

number_of_lines = 6464 ;

number_of_pixels = 6400 ;

global attributes:

group: GeolocationData

group: SnowData

3.1 Geolocation Data

The latitude and longitude data for each pixel in a swath are stored as auxiliary coordinate variables in the GeolocationData group in the VNP10. The coordinate variables, attributes and datasets use the netCDF CF-1.6 conventions for georeference. Software tools that work with the netCDF or HDF5 data formats should be able to work with the VNP10 product. Description of the GeolocationData group is given in List 2.

List 2. Description of the GeolocationData group and attributes in VNP10.

```
group: GeolocationData {
  variables:
    float latitude(number_of_lines, number_of_pixels) ;
      latitude:long_name = "Latitude data" ;
      latitude:units = "degrees_north" ;
      latitude:standard_name = "latitude" ;
      latitude:_FillValue = -999.f ;
      latitude:valid_range = -90.f, 90.f ;
    float longitude(number_of_lines, number_of_pixels) ;
      longitude:long_name = "Longitude data" ;
      longitude:units = "degrees_east" ;
      longitude:standard_name = "longitude" ;
      longitude:_FillValue = -999.f ;
      longitude:valid_range = -180.f, 180.f ;
} // group GeolocationData
```

3.2 SnowData Group

Descriptions of the SnowData group variables and attributes are given in List 3 and in Section 3.2.1. A few of the attributes are descriptive summary statistics compiled during a run of the algorithm that provide information on overall viewing conditions, e.g. cloud cover, extent of snow cover, basic QA, and threshold settings of some data screens. The purpose of these attributes is to provide an overall view of what is observed in the scene.

List 3. Description of SnowData group datasets and attributes in VNP10.

```
group: SnowData {
  variables:
    ubyte Algorithm_bit_flags_QA(number_of_lines, number_of_pixels) ;
      Algorithm_bit_flags_QA:coordinates = "latitude longitude" ;
      Algorithm_bit_flags_QA:long_name = "Algorithm bit flags" ;
      Algorithm_bit_flags_QA:flag_masks = "1b, 2b, 4b, 8b, 16b, 32b, 64b,
128b" ;
      Algorithm_bit_flags_QA:flag_meanings = "inland_water_flag
low_visible_screen low_NDSI_screen
combined_surface_temperature_and_height_screen/flag spare high_SWIR_screen/flag
spare solar_zenith_flag" ;
```

Algorithm_bit_flags_QA:comment = "Bit flags are set for select conditions detected by data screens in the algorithm, multiple flags may be set for a pixel.Default is all bits off" ;

```
    ubyte Basic_QA(number_of_lines, number_of_pixels) ;
        Basic_QA:coordinates = "latitude longitude" ;
        Basic_QA:long_name = "Basic QA value" ;
        Basic_QA:valid_range = 0UB, 3UB ;
        Basic_QA:mask_values = 211UB, 239UB, 250UB, 252UB, 253UB ;
        Basic_QA:mask_meanings = "211=night 239=ocean 250=cloud
252=no_decision 253=bowtie_trim" ;
        Basic_QA:key = "0=good, 1=poor, 2=bad, 3=other" ;
        Basic_QA:_FillValue = 255UB ;
    short NDSI(number_of_lines, number_of_pixels) ;
        NDSI:coordinates = "latitude longitude" ;
NDSI:long_name = "NDSI for land/inland water pixels" ;
        NDSI:valid_range = -1000s, 1000s ;
        NDSI:scale_factor = 0.001f ;
        NDSI:mask_values = 21100s, 23900s, 25100s, 25200s, 25300s, 25400s ;
        NDSI:mask_meanings = "21100=night, 23900=ocean,
25100=L1B_missing, 25200=L1B_unusable, 25300=bowtie_trim, 25400=L1B_fill" ;
        NDSI:_FillValue = 32767s ;
    ubyte NDSI_Snow_Cover(number_of_lines, number_of_pixels) ;
        NDSI_Snow_Cover:mask_meanings = "201=no decision, 211=night,
237=lake, 239=ocean, 250=cloud, 251=missing data, 252=L1B_unusable, 253=bowtie
trim, 254=L1B fill" ;
        NDSI_Snow_Cover:_FillValue = 255UB ;
        NDSI_Snow_Cover:coordinates = "latitude longitude" ;
        NDSI_Snow_Cover:long_name = "Snow cover by NDSI" ;
        NDSI_Snow_Cover:valid_range = 0UB, 100UB ;
        NDSI_Snow_Cover:mask_values = 201UB, 211UB, 237UB, 239UB,
250UB, 251UB, 252UB, 253UB, 254UB ;

// group attributes:
    :Surface_temperature_screen_threshold = "281.0 K" ;
    :Surface_height_screen_threshold = "1300 m" ;
    :Land_in_clear_view = "59.0%" ;
    :Cloud_cover = "41.0%" ;
    :Snow_Cover_Extent = "8.8%" ;
} // group SnowData
}
```

3.2.1 Variables

The VNP10 product has the following variables: NDSI_Snow_Cover, Basic_QA, Algorithm_bit_flags_QA and NDSI, each with attributes describing the data.

3.2.1.1 NDSI_Snow_Cover

The NDSI_Snow_Cover variable is the snow cover extent map generated by the algorithm. Snow cover is represented by NDSI values in the range of 0 – 100, from “no snow cover” to “total snow cover” in a pixel. To give a complete view of conditions in the scene the cloud mask, ocean mask, and night mask are overlaid on the NDSI snow cover data. Onboard VIIRS bowtie trim lines are retained in this swath product. An example of the NDSI_Snow_Cover dataset, with colored ranges of NDSI_Snow_Cover is shown in Figure 1.




Figure 1. VNP10.A2017105.2012.001.*.nc. NDSI_Snow_Cover. The western United States is imaged on 15 April 2017. Orientation is with north at bottom of image. Snow covered Rockies in the Colorado region is top center of swath, Uinta Mountain Range and Great Salt Lake are to the north and the Wind River Range, Yellowstone region and Big Horn Range, with cloud cover are to the north of that region.

3.2.1.2 Algorithm_bit_flags_QA

Algorithm-specific bit flags are set in this variable for the data screens that are applied in the algorithm. Multiple bit flags may be set for a pixel. For all pixels that were detected as snow the data screens were applied and the snow detection may have been reversed to “not snow” or flagged as “uncertain snow detection.” Algorithm bit flags are set if a snow detection was reversed or flagged as uncertain by one or more data screens applied in the algorithm. Some of the bit flags have a dual purpose to either reverse a snow detection or to flag an uncertain pixel result. Some screens are also applied to all land pixels in clear view. The Algorithm_bit_flags_QA variable for the swath shown in Fig.1, with selected bit flags displayed, is shown in Fig. 2. See Section 3.3.1 for description of the bit flags. Local attributes describing each bit flag are included.




Figure 2. VNP10.A2017105.2012.001.*.nc. Algorithm_bit_flags_QA variable that corresponds to the NDSI_Snow_Cover image in Fig.1. Only a few bit flags and bit flag combinations are shown to demonstrate how the bit flags can be used in evaluation/analysis of the snow cover. These bit flags indicate situations where there is increased uncertainty in snow cover detection along the edges of snow cover, cloud cover and shadowed landscape. At the scale of this image it is difficult to evaluate the results: zooming in on the maps is required for detailed analysis. An example of a detailed analysis is given in Section 3.4.

3.2.1.3 Basic_QA

A general quality value is given for pixels processed in the algorithm. Masked features, e.g. oceans, are set to mask values. This is a basic quality value used to indicate quality ranging from best quality to poor quality to provide a user with a value for initial quality assessment of the data. Local attributes describing the data are included.

3.2.1.4 NDSI

The calculated NDSI values for all land and inland water pixels without the cloud mask are in this variable. The NDSI data is packed (scaled) with an NDSI valid range of -1.0 to 1.0 (unpacked); the cloud mask is not applied in this variable but the ocean, night and other masks are. Local attributes describing the data are included.

3.3 Snow Cover Detection Algorithm Synopsis

A brief description of the algorithm approach is provided to explain the flow of the algorithm and the basic technique used to detect snow cover. A detailed description of the algorithm can be found in the VIIRS Snow Cover ATBD (Riggs et al., 2015).

The basis of the NASA VIIRS snow-mapping algorithms is the NDSI. Snow typically has very high visible (VIS) reflectance and very low reflectance in the shortwave infrared (SWIR), a characteristic commonly used to detect snow and to distinguish snow from most cloud types. The ability to detect snow cover is related to the difference in reflectance of snow cover in the VIS and SWIR in which the greater the VIS-SWIR difference, the higher the NDSI. The NDSI for VIIRS is:

$$\text{NDSI} = (I1 - I3) / (I1 + I3),$$

Where I1 is VIIRS band I1 (0.64 μm), and I3 is VIIRS band I3 (1.61 μm). The NDSI indicates the presence of snow cover on the surface based on the snow characteristics of high VIS reflectance and very low SWIR. If snow is present and viewable by a satellite then the NDSI will be in the theoretical range of -1.0 to +1.0, with a value of 0.0 or less indicating no snow. If snow is present and viewable by the sensor the NDSI will be > 0 . Snow cover is reported over the NDSI > 0.0 range.

A binary snow cover area (SCA) map is not output. Users may be familiar with the SCA map that was part of the MODIS C5 snow cover product suite but that SCA is not output in the MODIS C6 or C6.1 nor in the NASA VIIRS C1 snow cover products. Users may want a SCA map. A commonly accepted global NDSI threshold value of 0.4 has been used to make SCA maps, although many researchers have shown that better SCA maps can be made in specific situations for local or regional snow mapping if, for example, the NDSI is set for that situation with methods of threshold selection based on validation with meteorological station data, visual inspection/interpretation, empirical relationships or automated selection. In such cases the NDSI threshold setting may be as low as 0.1 for SCA identification. Thus an NDSI threshold selection is left to the user and is no longer static at a threshold of 0.4. In short, a user can make a SCA map using the NDSI_Snow_Cover or NDSI variables in the product by setting an NDSI threshold that is appropriate for a particular study area.

The NDSI is calculated in the algorithm for all land and inland water bodies in daylight, and then the data screens are applied. The data screens that are applied to each pixel depend on the NDSI value so different series of data screens can be applied to a pixel.

Applying data screens to a pixel can result in more than one data screen to be set for a non-snow detection, snow commission error or uncertain snow detection. A snow pixel that fails any single data screen will be reversed to 'not snow' and since multiple data screens are applied, more than a single Algorithm_bit_flag_QA flag may be set. The same data screens are applied to land and inland water pixels. Inland water bodies are mapped to bit 0 of the Algorithm_bit_flag_QA flags. The cloud mask, ocean mask, and night mask are laid on the NDSI_Snow_Cover to make a contextural map of snow cover. The NDSI value is output for all land and inland water pixels.

Data product inputs to the NASA VIIRS snow detection algorithm are listed in Table 2 and basic processing flow is depicted in Figure 3. The processing flow for a pixel is determined based on the land/water mask. Land and inland water bodies in daylight are processed for snow detection or ice/snow on water detection. VIIRS top-of-atmosphere (TOA) reflectance data are checked for missing or uncalibrated values; pixels with those values are set to a mask value and not processed for snow cover detection.

Table 2. VIIRS data product inputs to the VNP10 algorithm.

ESDT	Dataset	Center wavelength	Nominal spatial resolution
NPP_VIAES_L1	Reflectance_I1	0.640 μm	375 m
	Reflectance_I2	0.865 μm	375 m
	Reflectance_I3	1.61 μm	375 m
	BrightnessTemperature_I5	11.450 μm	375 m
NPP_VMAES_L1	Reflectance_M4	0.555 μm	750 m
NPP_IMFTS_L1	Latitude, Longitude, solar zenith angle		375m
VNP35_L2	Cloud confidence flag, land/water mask		750m




Figure 3. Snow Cover Environmental Data Record (EDR) processing architecture.

3.3.1 Data Screens

If a pixel has been determined to have some snow present based on the NDSI value, it is subjected to the following series of screens to alleviate snow commission errors and flag uncertain snow detections. Though snow typically has high VIS reflectance and low SWIR reflectance, the amount of reflectance in any band and the difference in reflectance between bands varies with viewing conditions and surface features. Screens are used to detect reflectance relationships that are atypical of snow and are applied to either reverse a snow detection to a ‘no snow’ or ‘other’ decision, or to flag the snow as ‘possibly not snow’ by Algorithm_bit_flags_QA flags. Bounding conditions of ‘too low reflectance’ or ‘too high reflectance’ are also set by screens. Each screen has a bit flag in the Algorithm_bit_flags_QA variable that is set to ‘on’ if a screen was failed. Users can extract specific bit flags for analysis.

3.3.1.1 Low VIS reflectance screen.

If the VIS reflectance from VIIRS band I1 is ≤ 0.10 or band M4 is ≤ 0.11 then a pixel fails to pass this screen. If a pixel is failed a “no decision” is the result. This screen is tracked in bit 1 of the Algorithm_bit_flags_QA.

3.3.1.2 Low NDSI screen

Pixels detected with snow cover in the $0.0 < \text{NDSI} < 0.10$ range are reversed to a ‘no snow’ result and bit 2 of the Algorithm_bit_flags_QA is set. That bit flag can be used to find where a snow cover detection was reversed to “not snow”.

3.3.1.3 Estimated surface temperature and surface height screen.

There is a dual purpose for this ‘estimated surface temperature linked with surface height screen’. It is used to alleviate snow commission errors on low elevations that appear spectrally to be similar to snow but are too warm to be snow. It is also used to flag snow detections on high elevations that are warmer than expected. If snow is detected in a pixel at elevations < 1300 m and that pixel has an estimated brightness temperature (BT) ≥ 281 K (using VIIRS band I5), that snow detection decision is reversed to “not snow” and bit 3 is set in the Algorithm_bit_flags_QA. If snow is detected in a pixel at elevations ≥ 1300 m and with estimated BT ≥ 281 K that snow detection is flagged as unusually warm by setting bit 3 in the Algorithm_bit_flags_QA.

3.3.1.4 High SWIR reflectance screen.

The purpose of this screen is to prevent non-snow features that are spectrally similar to snow from being detected as snow but also to allow snow detection in situations where snow cover SWIR reflectance is anomalously high. This screen has two threshold settings for different situations. While snow typically has SWIR reflectance less than about 0.20, in some situations, e.g., low sun angle, snow can have a higher reflectance in the SWIR. If a snow pixel has a SWIR reflectance in range of $0.25 < \text{SWIR} \leq 0.45$, it is flagged as unusually high for snow and bit 4 of Algorithm_bit_flags_QA is set. If a snow pixel has SWIR reflectance > 0.45 it is reversed to “not snow” and bit 4 of Algorithm_bit_flags_QA is set.

3.3.1.5 Solar zenith screen.

Low illumination conditions exist at SZAs $> 70^\circ$ which represents a challenging situation for snow cover detection. A SZA mask of $> 70^\circ$ is made by setting bit 7 of the Algorithm_bit_flags_QA. This mask is set across the entire swath. Night is defined as the SZA $\geq 85^\circ$ and pixels with SZA $\geq 85^\circ$ are masked as night.

3.3.2 Lake Ice Algorithm

The lake ice / snow covered ice detection algorithm is the same as the NDSI snow cover algorithm. Inland water bodies are tracked by setting bit 0 of Algorithm_bit_flags_QA. Users can extract or mask inland water bodies in the NDSI_Snow_Cover variable using this inland water bit flag. This algorithm uses the basic assumption that a water body is deep and clear and therefore absorbs all solar radiation incident upon it. Water bodies with high turbidity or algal blooms or other

conditions of relatively high reflectance from the water may be erroneously detected as snow/ice covered.

3.3.3 Cloud Masking

The cloud confidence flag from VNP35_L2 is used to mask clouds. The cloud confidence flag gives four levels of confidence: confident cloudy, probably cloudy, confident clear, and probably clear. If the cloud mask flags confident cloudy then the pixel is masked as cloud. If the cloud mask flag is set confident clear, probably clear or probably cloudy, it is interpreted as clear in the algorithm. The 750 m cloud mask is applied to the four corresponding 375 m pixels.

3.3.4. Quality Assessment (QA)

Two QA variables are output; the Basic_QA which gives a simple value score, and the Algorithm_bit_flags_QA which reports results of data screens as bit flags. The basic QA value is a qualitative estimate of the algorithm result for a pixel. The basic QA value is initialized to the good value and is adjusted based on the quality of the L1B input data and the solar zenith data screen. If the reflectance data is outside the range of 5-100% it is usable, but the QA value is set to poor. If the SZA is in the range of $70^\circ \leq \text{SZA} < 85^\circ$, the QA is set to poor which indicates increased uncertainty in results because of low illumination. If input data is unusable the QA value is set to 'other'. Conditions for a bad result are not defined. For features that are masked, e.g. ocean and night, the mask values are applied.

The Algorithm_bit_flags_QA dataset contains bit flags of data screen results applied in the algorithm. The data screens serve to provide more information on quality of an observation and the result, they indicate why a snow detection was reversed to "not snow," and indicate an uncertain snow detection, or challenging viewing conditions. More than one bit flag may be set because multiple data screens can be applied to a pixel. By examining the bit flags a user can determine if a snow cover result was changed to a "not snow" result by a screen or screens, or if a snow covered pixel has certain screens set to "on" indicative of an uncertain snow detection. The screens and bit flags have a dual purpose; some flag pixels where snow detection was reversed or flag snow detection as uncertain. More than one data screen can be on for a snow detection reversal or for uncertain snow detection, or for a non-snow detection.

Bits for the data screens are set to "on" if the screen was failed. Many combinations of bit flags may be set. A user can investigate any bit flag or combinations of bit flags. The inland water mask is also set as a bit flag (bit 0) to support analysis of inland waters for snow/ice cover.

3.4 Interpretation of Snow Cover Detection Accuracy, Uncertainty and Errors

The NASA VIIRS snow cover detection algorithm was designed to detect snow globally in all situations. The NDSI technique for snow detection has proven to be a robust

indicator of snow around the globe. Numerous investigators have used the MODIS snow products and reported accuracy statistics under cloud-free conditions in the range of 88-93%. (See listing of publications at <https://modis-snow-ice.gsfc.nasa.gov/?c=publications>). The MODIS and NASA VIIRS snow cover algorithms both use the same basic NDSI snow-detection algorithm, albeit adjusted for sensor and input data product differences.

In VIIRS C1 (and MODIS C6.1) snow cover data is output in two ways: 1) the NDSI based snow cover map that reports the NDSI value for snow over the 0.0 to 1.0 range with data screens applied to reduce error in snow cover detections and with masks of clouds and other features overlaid and 2) the NDSI data is output without data screens or cloud mask applied, but is masked for oceans and other features.

Snow cover is detectable with good accuracy when illumination conditions are near ideal, skies are clear, and several centimeters or more of snow are present on the landscape. Snow cover can occur on many different landscapes, including forests, plains and mountains, and under all types of viewing conditions. Viewing conditions change from day to day and across the landscape. The diversity of situations where snow may be found makes it challenging to develop a globally-applicable snow cover detection algorithm.

Analysis of MODIS C5 snow cover maps, with emphasis on snow cover omission and commission errors observed and reported in the literature prompted changes in the snow cover detection algorithm for MODIS C6.1 and VIIRS C1. The algorithm logic is as follows: snow cover always has an NDSI > 0 but not all features with NDSI > 0 are snow. Snow detection is applied to all land pixels in a swath then snow detections are screened to prevent possible snow commission errors, flag uncertain snow detections and set algorithm flags. Results of the data screens are set as bit flags in the `Algorithm_bit_flags_QA`. Multiple data screens are applied so it is possible that more than one bit flag is set for a pixel. Some situations associated with snow commission errors and possible ways to interpret the algorithm bit flags are discussed in following subsections.

3.4.1 Warm surfaces

Snow commission errors on warm surfaces with positive NDSI values can be reduced by screening based on estimated surface temperature. A surface temperature screen was applied in the MODIS C5 snow-mapping algorithm to reverse all snow detections that were thought to be too warm to be snow. A decision on any pixel detected as snow cover and having an estimated surface temperature > 283 K was reversed to “no snow.” That temperature screen was shown to dramatically reduce the occurrence of erroneous snow cover in warm regions of the world and along warm coastal regions. However, it was discovered that the temperature screen also caused significant snow omission errors in spring and summer on snow covered mountain ranges. These errors could be very large as the average surface temperature within a pixel increased above 283 K. The effect of the temperature screen on mapping of snow cover on the Sierra Nevada from 1 May to 1 August 2010 is shown at [https://modis-snow-](https://modis-snow-ice.gsfc.nasa.gov/?c=publications)

ice.gsfc.gov/?c=collection6. Snow omission errors were around 10% at start of that time period then rose to near 90% by August.

In MODIS C6, C6.1 and NASA VIIRS C1 the surface temperature screen is combined with surface elevation and used in two ways. This combined screen reverses snow cover detection on low elevation < 1300 m surfaces that are too warm for snow and the Algorithm_bit_flags_QA bit 3 is set. Snow cover detection at ≥ 1300 m on a surface that is too warm for snow is not reversed but that snow cover detection is flagged as “too warm” by also setting that bit flag.

A possible effect of this screen may be observed along the edge of mountain snow where snow cover detection is changed to “no snow”. An example of the situation is shown in Fig. 4 on the Uinta Mountain Range, Utah, subimage from Fig. 1, where snow cover detection was changed to “not snow” by only the combined high elevation and surface temperature screen (bit 3 in the Algorithm_bit_flags_QA variable) shown by the red pixels in the right image. Detecting snow along the edge of the mountain snow cover, i.e. defining exactly where snow cover ends, is a challenge. Visual evaluation of visible and temperature data and NDSI_Snow_Cover find that this screen does have an effect along the edge of snow cover and that it can be interpreted as showing an edge or band of one to few pixels of uncertainty in snow cover. It is possible to change those “no snow” pixels to “snow” pixels by using the unique combination of NDSI_Snow_Cover = 0 and bit 3 on (Algorithm_bit_flags_QA = 8) and the corresponding NDSI data value to restore snow to the pixel. This demonstrates that a user has options on how to interpret and utilize the variables for their research or application.




Figure 4. Uinta Mountain Range, Utah, with only the combined high elevation and surface temperature screen showing were snow cover detection was changed to a “no snow” cover result in the NDSI_Snow_Cover variable. The left image is a false color of a VIIRS bands I1, I2, I3 subimage from NPP_VIAES_L1.A2017105.2012.001.*.nc. The right image is the corresponding NDSI_Snow_Cover

from VNP10.A2017105.2012.001*.nc (Fig. 1) with where only the Algorithm_bit_flags_QA combined high elevation and surface temperature screen (bit 3) changed a snow detection to “no snow” shown in red.

The effectiveness of the surface temperature and height screen varies as the surface changes over seasons. It is effective at reversing snow commission errors of some surface features, and cloud contaminated pixels over some landscapes when the surface is warm. However when the surface is below the threshold temperature, or cloud contamination lowers the estimated surface temperature, this screen is not effective. A surface feature that is spectrally similar to snow, for example the Bonneville Salt Flats, may have snow detection reversed by this screen when the surface is warm but may not be reversed when the surface is cold and snow-free in the winter.

3.4.2 Low reflectance

Situations of low reflectance from the surface for various reasons pose challenges to snow detection and may cause snow commission errors. Several data screens and bit flags are set for low reflectance conditions.

Low solar illumination conditions occurring when the SZA is $> 70.0^\circ$ and near to the day/night terminator are a challenge to snow detection. That situation is identified when the low solar zenith flag (bit 7) in the Algorithms_bit_flags_QA variable set to on.

Low reflectance situations in which reflectance is $\sim 30\%$ across the visible bands is also a challenge for snow detection. Low reflectance across the VIS and SWIR bands can result in relatively small differences between the VIS and SWIR bands and can give an NDSI > 0 for some non-snow covered surfaces. Investigation and discussion with some users of the MODIS C5 products who encountered errors associated with low reflectance surface conditions resulted in setting a low reflectance limit in the algorithm. If VIS reflectance is too low, a pixel is set to “no decision” and the low VIS data screen bit flag is set in the Algorithm_bit_flags_QA variable. This is considered a low limit to accurate detection of snow cover on the landscape. The NDSI is calculated for those “no decision” results so a user can see the NDSI value by using the low visible Algorithm_bit_flags_QA and NDSI variables.

Low reflectance associated with low illumination, landscape shadowed by clouds or terrain, and unmapped water bodies or inundated landscape can exhibit reflectance characteristics similar to snow and thus be erroneously detected as snow by the algorithm. Very low visible reflectance is a cause for increased uncertainty in detection of snow cover. Though the data screens applied can prevent snow commission error, some snow commission errors can go undetected, notably on cloud shadowed snow-free landscape as shown in Fig. 5. A subimage is shown from a region of scattered clouds over snow free land near the east side of nadir track north of the Big Horn Range in Wyoming from VNP10 (Fig. 1); a few scattered clouds and their shadows on the surface may be seen, left image Fig. 5, and the NDSI_Snow_Cover, center image, with snow commission errors associated with the shadowed surface. In this situation the cloud mask detects the clouds, and several of the data screens prevent snow commission errors as shown by the colored pixels, right image, that are bit flags set to

prevent snow commission errors. However snow commission errors, blue pixels center image, occur that are associated with the periphery of cloud and shadowed land.



Figure 5. Example of snow commission associated with cloud periphery and shadowed surfaces. On the left image (50 x 50 pixels) is a false color of VIIRS bands I1, I2, I3 subimage from NPP_VIAES_L1.A2017105.2012.001*.nc. The center image is the corresponding NDSI_Snow_Cover from VNP10.A2017105.2012.001*.nc (Fig. 1). Algorithm_bit_flags_QA set to on to prevent snow commission error are shown as colored pixels in the right image.

3.4.3 Low NDSI

Low VIS reflectance situations, snow covered or snow free surfaces, where the difference between VIS and SWIR is very small can have very low positive NDSI values. Those low positive NDSI results can occur where visible reflectance is low or high and where the associated SWIR is low or high but slightly lower than the VIS so that the NDSI is a very low positive value. In our analysis of many such situations we found that very uncertain snow detections or snow commission errors were common when the $0.0 \leq \text{NDSI} < 0.1$. Based on that analysis a low NDSI screen is applied. If NDSI is < 0.1 a snow detection is reversed to “not snow,” and the low NDSI bit 2 flag is set in the Algorithm_bit_flags_QA. To determine if these situations were found in a swath, the bit 2 flag can be used to find them and the corresponding NDSI value where snow detections were reversed.

3.4.4 High SWIR reflectance

Unusually high SWIR reflectance may be observed for some snow cover situations, from some types of clouds not masked as confident cloudy or from non-snow surface features. A SWIR screen is applied at two thresholds to either reverse a possible snow commission error or flag snow detection with unusually high SWIR. A user can check this bit flag to find where uncertain snow cover detections occurred or where snow detection was reversed to “not snow.”

3.4.5 Cloud and snow confusion

Cloud and snow confusion in the VIIRS C1 snow cover is similar to the snow and cloud confusion seen in the MODIS C6 snow cover product. Both cloud masks are cloud conservative which tend to favor cloud detection over cloud clearing. Two common

sources of cloud/snow confusion are that the cloud mask does not correctly flag cloudy or clear conditions, and where subpixel clouds (cloud mask is at 750 m resolution) escape detection.

The cloud mask algorithm uses many tests to detect cloud and the combination of tests applied to a pixel, the processing path, depends on whether or not the surface is snow covered. (Details of the cloud mask algorithm and product can be found in the cloud mask Operational Algorithm Document (OAD)

[<https://jointmission.gsfc.nasa.gov/documents.html>, Document # 474-0062 VIIRS Cloud Mask (VCM) Intermediate Product (IP) Software – OAD Revision I]. An external snow/ice background map and an internal check for snow cover is made in the cloud mask algorithm; if that initial determination for snow is incorrect then the wrong processing path is followed and a possible erroneous cloud determination is made, e.g. flagging snow as confident cloudy. An example of that can be seen in Fig. 4 where there are clouds along the periphery of the mountain snow cover but there is no visible evidence of clouds in the corresponding visible color image. This situation can be seen associated with swaths of snow cover from storms crossing the Great Plains where snow on the periphery of the snow covered region is flagged as confident cloudy by the cloud mask. We have investigated this cloud/snow confusion situation and found that the snow was detected as cloud by only a single visible cloud test of the several cloud spectral tests applied in the processing path. We found that by examining the cloud mask algorithm processing path and results of all cloud spectral tests applied, that the cloud mask could be reinterpreted as clear in that specific situation and the snow could then be correctly detected. That reinterpretation test was partially effective at resolving this specific cloud/snow confusion situation however in a global application of that test, inconsistencies in results were found, and so further investigation is required.

Subpixel size clouds that escape detection as confident cloudy by the cloud mask algorithm may be detected as snow in the snow algorithm because the cloud reflectance can cause an underlying snow free surface to have one or more reflectance features similar to snow. This situation frequently results in snow commission errors associated with the periphery of clouds, especially with cloud formations of scattered, popcorn-like cloud formations over vegetated landscapes. Multilayer cloud formations where there are different types of clouds, warm and cold, and where cloud shadows fall on clouds may have some regions of the cloud cover not detected as confident cloudy which may then be detected as snow in the snow cover algorithm. In those types of cloud cover conditions the subpixel contaminated clouds and self-shadowed clouds are spectrally indistinct from snow in the algorithm. Use of the cloud mask algorithm processing path flags and individual cloud spectral tests flags possibly in combination with other screens for snow reflectance holds promise for resolving some snow/cloud confusion situations and is being investigated. Cloud conditions are typically transient. Such transient cloud/snow commission errors can possibly be filtered temporally or spatially or by a combination of filters developed by users.

3.4.6 Lake ice

A lake ice detection algorithm is included in the VIIRS C1 to map ice or snow and ice covered lakes and rivers. The lake ice detection algorithm is similar to the snow cover detection algorithm with lake ice cover included in the NDSI_Snow_Cover variable. The lake ice algorithm is the same as the NDSI snow detection algorithm. Inland water bodies are mapped in bit 0 of the Algorithm_bit_flags_QA variable for use in analysis of lake ice.

Lake ice is included in the NDSI_Snow_Cover variable so that a spatially coherent image of a snow covered landscape can be seen. A user can extract the inland water mask from bit 0 of the Algorithm_bit_flags_QA variable for use in analysis or to apply as a static water mask.

Visual analysis of VNP10 swaths and experience with the MOD10_L2 products acquired during the boreal winter when lakes are frozen finds that snow/ice covered lakes are detected with 90-100% accuracy. Disappearance of lake ice also appears to be detected with high accuracy. During the ice free season, changes in physical characteristics of a lake can greatly affect the accuracy of the algorithm. Sediment loads, high turbidity, aquatic vegetation and algae blooms change the reflectance characteristics and may cause erroneous lake or river ice detection in the spring or summer. A lake-ice-specific algorithm should be developed in a future version of the algorithm.

3.4.7 Bright surface features

Surface features such as salt flats, bright sands, or sandy beaches that have VIS and SWIR reflectance characteristics similar to snow may be detected as snow cover based solely on the NDSI value, thus resulting in errors of commission. The data screens applied in the algorithm can reduce the occurrence of snow commission errors in some situations, e.g., a low elevation; too-warm surface can be blocked by the combined surface temperature and height screen, but may not be effective in other situations. These types of surface features are static so a user could mask or flag these surfaces relevant to a specific research or application.

3.4.8 Land/water mask

In this C1 processing of VNP10 the land/water mask is read from the VNP35 product. That land /water mask (flag) is determined using the Quarterly Surface Type/Land Water Mask specifically for the cloud mask algorithm. That land/water mask was used because it was available in the production stream of input products. That land/water mask is not the same as the MODIS land/water mask used in MOD10_L2 C6 so it is possible that differences in locations of coastlines and water bodies may be observed when comparing VNP10 to MOD10_L2.

When the NASA VIIRS geolocation data product VNP03IMG that contains the MODIS C6.1 land/water mask becomes available, along with the NASA L1B products, the

VNP10 algorithm will be revised to use those inputs in C2. So this section will need to be revised.

3.4.9 Geolocation accuracy

Geolocation accuracy is good which provides consistent high accuracy in mapping of the VIIRS data products (<https://viirsland.gsfc.nasa.gov/index.html>). The small errors in geolocation are negligible in the swath level products, however geolocation error may be observed in the daily gridded products as a shifting of features, e.g. lake shore line location, in cells from day to day. *A comment on geolocation accuracy in gridded products will be provided later when those products are added to future versions of this User Guide.*

3.4.10 Antarctica

The Antarctic continent is nearly completely ice and snow covered year 'round, with very little annual variation, though some changes are observable on the Antarctic Peninsula. The NASA VIIRS C1 snow cover detection algorithm is run for Antarctica without any Antarctica-specific processing paths. The resulting snow cover map may show some snow-free areas which is an obvious error. That error is related primarily to the great difficulty in detecting clouds over the Antarctic continent. The similarity in reflectance and lack of thermal contrast between clouds and ice/snow cover, sometimes related to thermal inversions, are major challenges to accurate snow/cloud discrimination over the Antarctic continent. In situations where the cloud mask fails to identify confident cloudy the snow algorithm assumes a cloud-free view and either identifies the surface as "not snow covered" or identifies the cloud as snow. In either case the result is wrong. Though the VNP10 is generated for Antarctica, it must be carefully scrutinized for accuracy and quality.

4.0 VNP10A1

The daily, gridded and projected, snow cover product is produced by algorithms that select the 'best' observation of the day from the single to multiple VIIRS observations that were acquired on a day. The VNP10A1 is in HDF5-EOS format but also has additional variables and attributes that follow netCDF CF-1.6 conventions for local and global attributes and for georeference, that allow for more tools to work with the product.

The VNP10A1 product contains the same datasets as in the VNP10 product, but in a different file format. There is no snow detection done in the processing of the VNP10 swath products, the algorithms grid and project the swath data products on to the sinusoidal projection. The VIIRS bowtie striping present in the VNP10 is removed in the gridding and reprojection processing. When there is more than one observation in a grid cell the observations are stacked, in no particular order, to produce an intermediate product that has multiple observations stored for grid cells. (These intermediate products are not archived at the NSIDC DAAC.) Then a selection algorithm is run with the intermediate products, and viewing geometry products, as input and the 'best'

observation based on solar zenith angle, distance from nadir and observation coverage in a grid cells. The 'best' observation for each product is based only on those criteria so that the observation selected is nearest local solar noon time, nearest the orbit nadir track and with most coverage in a grid cell, which is considered the best sensor view of the surface on a day relevant to snow cover detection. This strategy results in a contiguous mapping of swaths with a weave or checkerboard pattern along stitched-together swath edges within a tile. That weave pattern is sometimes apparent where cloud cover changed between acquisition times of overlapping swaths.

Included in VNP10A1 is a pointer variable that points to the granule (VNP10 swath) from which each observation was selected. That pointer can be linked to the beginning and/or ending times of the individual input swaths stored as global attributes to determine the date and time of acquisition of each observation.

4.1 Variables

The VNP10A1 product has these variables: NDSI_Snow_Cover, Basic_QA, Algorithm_bit_flags_QA, NDSI, granule_pnt and Projection, local attributes are attached to the variables. The local attributes follow netCDF CF-1.6 conventions. The variable Projection is for CF-1.6 georeference of the variables. A brief description of each variable is give in following sections and a listing of the complete file contents is given in Appendix B.

4.1.1 NDSI_Snow_Cover

The NDSI_Snow_Cover variable is the snow cover extent made with the selected 'best' observations from the VNP10 product where the snow detection algorithm was run. Snow cover is represented by NDSI values in the range of 0 – 100, from “no snow cover” to “total snow cover” in a pixel. To give a contextual view of snow cover in the tile on a day, clouds, oceans, inland water and other flag values are included with the snow cover data. An example of the NDSI_Snow_Cover dataset, with colorized ranges of NDSI is shown in Figure 6. See Appendix B for listing of the local attributes.




Figure 6. NDSI_Snow_Cover in VNP10A1.A2018007.h10v04.*.h5. Un-projected image of tile h10v04 that covers from southern Nebraska (bottom) to northern Montana (top), snow cover on the Rockies and High Plains.

4.1.2 Basic_QA

The Basic_QA variable is a general quality value assigned to observations in the VNP10 algorithm. This is a basic quality value used to indicate quality ranging from highest to poor to provide a user with a convenient value for initial quality assessment of the data. Features, e.g. oceans, are set to flag values. See Appendix B for listing of the local attributes.

4.1.3 Algorithm_bit_flags_QA

Algorithm-specific flag masks in this variable are the result of data screens that were applied in the VNP10 algorithm. Algorithm flag masks were set for certain conditions observed in the VNP10 algorithm. These flag masks provide QA information regarding an observation and can be used to determine if a snow detection was flagged as uncertain or reversed to no snow by one or more data screens applied in the algorithm. Some of the flag masks have a dual purpose to either reverse a snow detection or to flag an uncertain observation result. Multiple flag masks may be set for an observation. See Section 3.3.1 for description of flag masks. See Appendix B for listing of the local attributes.

4.1.4 NDSI

The NDSI values for all land and inland water pixels without the cloud mask applied are given in this variable. These are the NDSI values calculated in VNP10 and correspond to the 'best' observation selected. The NDSI is packed data that can be unpacked using the `scale_factor` local attribute. NDSI valid range is -1.0 to 1.0, when unpacked and has flag values for ocean, night and other conditions. See Appendix B for listing of the local attributes.

4.1.5 granule_pnt

The `granule_pnt` variable data is a pointer, a numeric value that points to the index of values stored in the global attributes `GranulePointerArray`, `GranuleBeginningDateTime` and `GranuleEndingDateTime`. This pointer points to the index of the value in those arrays from which the 'best' observation was selected. Non-negative values in `GranulePointerArray` correspond by index to the `GranuleBeginningDateTime` and `GranuleEndingDateTime` arrays. The time of an observation can be determined using the pointer variable and those data arrays. See Appendix B for listing of the local and global attributes.

4.1.6 Projection

Projection is an empty variable, it has no data. The Projection variable serves as a container with local attributes that provide information on the projection. These local attributes follow CF-1.6 convention for georeference and are used by tools to project or re-project from the native sinusoidal projection. See Appendix B for listing of the local attributes.

4.2 Interpretation of Snow Cover Detection Accuracy, Uncertainty and Errors

The interpretation for accuracy, uncertainty and errors for snow cover detection is the same as for the VNP10 product. Refer to Section 3.4 for discussion of accuracy and errors.

Geolocation error may be seen in the product due to uncertainty or error that may occur in L2 geolocation, and gridding and projecting the swath data to the sinusoidal projection from swath latitude and longitude reference system. That type of geolocation error is commonly observed in the location of freshwater bodies over time. In a composite of a tile over the course of several consecutive days the position of a lake shoreline may shift by one or more cells in the horizontal or vertical directions each day resulting in a blurred outline of the lake composited over time.

5.0 VNP10A1F

The purpose of the daily cloud-gap-filled (CGF) snow cover product is to provide a daily "cloud-free" map of snow cover extent. The CGF daily map is generated by retaining a previous day non-cloud observation when the current day has a cloud observation. The data product includes the CGF snow cover map, a cloud persistence map, the basic QA and algorithm QA flags for observations, and the NDSI snow cover map from the

corresponding daily VNP10A1 product. The snow cover map from VNP10A1 is included to facilitate comparison with the CGF snow cover map. An example of VNP10A1F is shown in Figure 7. The VNP10A1F is in HDF5-EOS format but also has additional variables and attributes that follow netCDF CF-1.6 conventions for local and global attributes and for georeference, that allow for more tools to work with the product. A brief description of the algorithm and each variable is given in following sections and a listing of the complete file contents is given in Appendix C.




Figure 7. CGF_NDSI_Snow_Cover in VNP10A1F.A2018007.h10v04.*.h5. Un-projected image of tile h10v04 that covers from southern Nebraska (bottom) to northern Montana (top), snow cover on the Rockies and High Plains. A virtually ‘cloud free’ image of snow cover is attained in this image on the seventh consecutive day of a test run of the algorithm. Compare this image to the daily NDSI_Snow_Cover in Figure 6 to see the effect of the CGF algorithm in producing a ‘cloud free’ image.

5.1 Algorithm Description

The VIIRS CGF algorithm is the same as the MODIS CGF algorithm. The MODIS CGF algorithm was developed for the daily global M*10CMG and evaluated in a data assimilation experiment with the Noah land surface model in the NASA Land Information System (LIS) (Hall et al., 2010) was adapted for use with the M*D10A1 daily snow cover product. Since the M*D10A1 product is the most often used snow cover product the CGF product was developed at this level to provide the opportunity for maximum usage of the product.

Inputs to the CGF algorithm are the current day VNP10A1 and the previous day VNP10A1F. The CGF current day snow cover map is generated by replacing the current day cloud observations in VNP10A1 with a non-cloud observation from previous day CGF snow cover map. Cloud persistence is tracked by incrementing or resetting the count of consecutive days of cloud observed for a cell in the Cloud_Persistence variable. Cloud tracking is done by incrementing the count of consecutive cloud days in VNP10A1F. If the current day is a cloud observation, then the count is incremented by one day. If the current day is a non-cloud observation then the cloud persistence count is reset to 0. The Basic_QA and the Algorithm_Bit_Flags_QA datasets in VNP10A1F are also set to the current day non-cloud observation corresponding QA data value of VNP10A1 or replaced with previous day VNP10A1F values if current day observation is cloud. The VNP10A1F also contains a copy of the current day VNP10A1 NDSI_Snow_Cover variable to facilitate comparison with the CGF snow cover.

The CGF product will be produced as a 12 month sequence corresponding to the United States Geological Service (USGS) “water year” beginning on 1 October and ending on 30 September of each year. The exception is that for the first year of S-NPP production will begin on 19 January 2012 the start date of data collection. On the first day of the “water year” (or first day of temporal coverage) the VNP10A1F is produced as a copy of the VNP10A1 variables, with the cloud persistence data set to one for cells that are cloudy.

In situations where there is fill data in orbit gaps or missing parts of swaths, the fill data is replaced with a non-fill data value from the previous day CGF and the cloud persistence count is incremented by one.

There are some missing days of VNP10A1 tiles in the data record. When a missing tile is encountered, the previous day VNP10A1F becomes the current day CGF and the cloud persistence data is incremented by one for all the cells. In this situation the missing data is treated as a cloud observation and the cloud persistence count of days is incremented by one. The global attribute “MissingDaysOfVNP10A1” reports the number of missing day(s). There are some gaps in the data record that are longer than a single day.

5.2 Variables

The VNP10A1F product has these variables: CGF_NDSI_Snow_Cover, Basic_QA, Algorithm_Bit_Flags_QA, Cloud_Persistence, VNP10A1_NDSI_Snow_Cover and Projection. Local attributes are attached to the variables follow netCDF CF-1.6 conventions. The variable Projection is for CF-1.6 georeference of the variables. Brief description of each variable is give in following sections and a listing of the complete file contents is given in Appendix C.

5.2.1 CGF_NDSI_Snow_Cover

The CGF_NDSI_Snow_Cover variable is the cloud-gap-filled snow cover extent resulting from the algorithm. Basically the current day cloud observations are replaced

with non-cloud observations from a previous day. See Section 5.2 for a description of the algorithm. Snow cover is represented by NDSI values in the range of 0 – 100, from “no snow cover” to “total snow cover” in a pixel. To give a contextual view of snow cover in the tile clouds, oceans, inland water and other flag values are included with the snow cover data. See Appendix C for listing of the local attributes.

5.2.2 Basic_QA

The Basic_QA variable is a general quality value assigned to observations in the VNP10 algorithm. This is a basic quality value use to indicate quality ranging from highest to poor to provide a user with a convenient value for initial quality assessment of the data. Features, e.g. oceans, are set to flag values. The observation selected corresponds to the observation selected for the CGF_NDSI_Snow_Cover variable. See Appendix C for listing of the local attributes.

5.2.3 Algorithm_bit_flags_QA

Algorithm specific flag masks in this variable are the result of data screens that were applied in the VNP10 algorithm. Algorithm flag masks were set for certain conditions observed in the VNP10 algorithm. These flag masks provide QA information regarding an observation and can be used to determine if a snow detection was flagged as uncertain or reversed to no snow by one or more data screens applied in the algorithm. Some of the flag masks have dual purpose to either reverse a snow detection or to flag uncertain observation result. Multiple flag masks may be set for an observation. See Section 3.3.1 for description of flag masks. The observation selected corresponds to the observation selected for the CGF_NDSI_Snow_Cover variable. See Appendix C for listing of the local attributes.

5.2.4 Projection

Projection is an empty variable, it has no data. The Projection variable serves as a container with local attributes that provide information on the projection. These local attributes follow CF-1.6 convention for georeference and are used by tools to project or re-project from the native sinusoidal projection. See Appendix C for listing of the local attributes.

5.2.5 VNP10A1_NDSI_Snow_Cover

The VNP10A1_NDSI_Snow_Cover variable is a copy of the NDSI_Snow_Cover variable from the VNP10A1 input. It is include to facilitate a convenient comparison to the CGF_NDSI_Snow_Cover variable. See Section 4.0 for a description of the algorithm. Snow cover is represented by NDSI values in the range of 0 – 100, from “no snow cover” to “total snow cover” in a pixel. To give a contextual view of snow cover in the tile clouds, oceans, inland water and other flag values are included with the snow cover data. See Appendix C for listing of the local attributes.

5.3 Interpretation of Snow Cover Accuracy, Uncertainty and Errors

The CGF snow cover map is an estimate of the snow cover that might exist under current cloud cover. The CGF snow cover map is made by using all the cloud-free

observations from the current day and by replacing the current day cloud observations with the non-cloud (clear view) observation from the previous day VNP10A1F. The persistence of cloud cover is tracked by incrementing the count of days of cloud cover for each consecutive day that cloud is observed in a cell. The number of days since the last non-cloud observation in a cell is tracked in the Cloud_Persistence variable. When a cloud-free observation is made the cloud persistence day count is reset to 0. If the cloud persistence is 0 for a grid cell that means that a cloud-free observation was made on the current day. A cloud persistence value of 1 means that current day was cloudy. A cloud persistence value greater than 1 is the count of previous consecutive days of cloud cover observed for a cell; it is the number of days since a non-cloud observation was observed. The Cloud_Persistence variable should be used as quality assessment (QA) data when using the CGF_NDSI_Snow_Cover variable to determine the age of an observation. The Basic_QA and Algorithm_Bit_Flag_QA variables are copied from the VNP10A1 for non-cloud observations and from previous day VNP10A1F, for cloudy observations. The snow Basic_QA and Algorithm_Bit_Flag_QA were set in the L2 processing so the user is referred to Section 3.4 for discussion regarding interpretation of these variables.

On the first day of VNP10A1F production the CGF snow map will be the same as the VNP10A1; on successive days the cloud cover in the CGF will decline, eventually to zero, as non-cloud observations replace cloud observations over time. A reasonable estimate of the number of days to reach a nearly cloud free CGF is five to seven days, but is dependent on the season and location imaged. The production plan is to produce VNP10A1F for each day of a USGS “water year” beginning on 1 October and ending on 30 September, except for the first year of S-NPP which begins 19 January 2012. A user can determine if the VNP10A1F is the initial day of a time series or a day in the series by reading the global attribute FirstDayOfSeries. FirstDayOfSeries” is set to “Y” for the first day in a time series and to “N” for all other days in the time series. The global attribute “TimeSeriesDay” is the count of days in the series since the first day.

Interpretation/evaluation of the CGF snow cover relative to the current day VNP10A1 snow cover can be done by comparing the CGF to the VNP10A1_NDSI_Snow_Cover variable which is copied from the VNP10A1 product to facilitate this comparison. The discussion of interpretation, accuracy and errors for CGF snow cover is similar to that for VNP10 in Section 3.4 where the snow detection algorithm is applied.

Cloud/snow confusion, confusing snow cover as cloud in the VNP10 algorithm, can be passed through to the VNP10A1F and may appear as persistent cloud cover over more days than would seem reasonable for cloud cover to persist in a location at a time when snow cover is expected. Analysis/evaluation of persistent cloud cover may include comparison with either the VNP02MOD and VNP02IMG reflectance products, or the VNP09A1 daily surface reflectance product. Cloud/snow confusion continues to be investigated in the VNP10 algorithm and if that confusion can be decreased, the quality of the sequence of VNP10* snow products would be enhanced.

There are some single day or multiple days missing VNP10A1 products (tiles) in the S-NPP data record. The CGF algorithm processes a missing day as a completely cloudy day; the previous day VNP10A1F becomes the current day VNP10A1F and the cloud persistence count is incremented by one. The global attribute MissingDaysOfVNP10A1 reports the number of missing day(s), it is incremented by one for each missing day then reset to 0 when VNP10A1 is again available. S-NPP VIIRS data outages are listed at https://modaps.modaps.eosdis.nasa.gov/services/production/outages_npp.html. A single day of missing data has minimal impact on the continuity of snow cover, however the impact can vary temporally and by region. The effect of multiple consecutive days of missing VNP10A1 tile inputs has not been assessed but would probably be significant, especially during periods when snow cover could be reasonably expected to occur.

Orbit gaps and missing swath data that appear in the VNP10A1 product are filled with fill data. In the CGF algorithm fill data is processed in a manner similar to how a cloud observation is processed. A fill data value is replaced with a non-fill data value from yesterday's VNP10A1F and the cloud persistence count is incremented by one. If the observation from previous day VNP10A1F is fill data then fill data is written for the cell and the cloud persistence count is incremented by one. The objective of processing fill data in this way is to provide a CGF snow map without fill data disrupting the continuity of the CGF snow cover map over time. However, situations of persistent fill data will be retained as fill data until non-fill data is available.

=====

6.0 Related Web Sites

Suomi-NPP

<https://jointmission.gsfc.nasa.gov/index.html>

VIIRS

VIIRS Land: <https://viirsland.gsfc.nasa.gov/>

VIIRS Cryosphere: <https://viirsland.gsfc.nasa.gov/Products/NASA/CryoESDR.html>

MODIS Snow/Ice Global Mapping Project:

<https://modis-snow-ice.gsfc.nasa.gov>

Imagery and Data Product Viewing

Worldview: <https://worldview.earthdata.nasa.gov>

LANCE: <https://wiki.earthdata.nasa.gov/display/GIBS/2015/12/10/VIIRS+is+Here>
<https://earthdata.nasa.gov/earth-observation-data/near-real-time/download-nrt-data/viirs-nrt>

NSIDC Data Ordering & User Services

National Snow and Ice Data Center: <http://nsidc.org/data/viirs>

LAADS DAAC

<https://ladsweb.modaps.eosdis.nasa.gov>

HDF5

The HDF Group: <https://www.hdfgroup.org/HDF5/>

NetCDF

<http://www.unidata.ucar.edu/software/netcdf/docs/index.html>

7.0 References

Riggs, G., Hall, D.K. and Román, M.O. 2015 VIIRS Snow Cover Algorithm Theoretical Basis Document (ATBD) https://modis-snow-ice.gsfc.nasa.gov/uploads/VIIRS_snow_cover_ATBD_2015.pdf

8.0 List of Acronyms

ATBD	Algorithm Theoretical Basis Document
BT	Brightness Temperature
Cx	Collection number
CDR	Climate Data Record
CMG	Climate-Modeling Grid
EDR	Environmental Data Record
EOSDIS	Earth Observing System Data Information System
ESDT	Earth Science Data Type
FSC	Fractional Snow Cover
HDF5	Hierarchical Data Format 5
IDPS	Interface Data Processing Segment
L1 / L2 / L3	Level 1, Level 2 or Level 3 data product
LSIPS	Land Science Investigator-led Processing System
MOD10	ESDT of the MODIS L2 snow cover product
MODIS	Moderate-resolution Imaging Spectroradiometer
NDSI	Normalized Difference Snow Index
QA	Quality Assessment
SCA	Snow covered Area
SIN	Sinusoidal Projection
S-NPP	Suomi National Polar-orbiting Partnership
SWIR	Short Wave Infrared
SZA	Solar Zenith Angle
TOA	Top-of-Atmosphere
VIIRS	Visible Infrared Imager Radiometer Suite
VNP10	ESDT name for the VIIRS Level-2 swath-based Snow Cover Data Product
VNP10A1	ESDT name for the VIIRS Level-3 tiled Snow Cover Data Product
VNP10A1F	ESDT name for the VIIRS Level-3 tiled Cloud-gap-filled Snow Cover Data Product
VNP10C1	ESDT name for the VIIRS Level-3 global Snow Cover Data Product
VIS	Visible

9.0 Appendix A VNP10 Global Attributes

Example of VNP10 global attributes.

```
// global attributes:
:QAPercentCloudCover = "41.0%" ;
:Snow_Cover_Extent = "8.8%" ;
:QAPercentBestQuality = "99.9%" ;
:QAPercentGoodQuality = "0.1%" ;
:QAPercentPoorQuality = "0.0%" ;
:QAPercentOtherQuality = "0.0%" ;
:RangeEndingTime = "20:18:00.000000" ;
:RangeBeginningTime = "20:12:00.000000" ;
:AlgorithmType = "OPS" ;
:processing_level = "Level 2" ;
:creator_email = "modis-ops@lists.nasa.gov" ;
:AlgorithmVersion = "NPP_PR10 1.0.3" ;
:PGEVersion = "1.0.6" ;
:ProductionTime = "2017-04-16 05:25:46.000" ;
:keywords_vocabulary = "NASA Global Change Master Directory (GCMD)
Science Keywords" ;
:title = "VIIRS Snow Cover Data" ;
:naming_authority = "gov.nasa.gsfc.VIIRSLand" ;
:publisher_name = "LAADS" ;
:creator_url = "http://ladsweb.nascom.nasa.gov" ;
:RangeBeginningDate = "2017-04-15" ;
:WestBoundingCoord = -135.0753f ;
:Conventions = "CF-1.6" ;
:ProcessingEnvironment = "Linux minion7239 3.10.0-514.10.2.el7.x86_64
#1 SMP Fri Mar 3 00:04:05 UTC 2017 x86_64 x86_64 x86_64 GNU/Linux" ;
:VersionID = "001" ;
:StartTime = "2017-04-15 20:12:00.000" ;
:SatelliteInstrument = "NPP OPS" ;
:stdname_vocabulary = "NetCDF Climate and Forecast (CF) Metadata
Convention" ;
:publisher_url = "http://ladsweb.nascom.nasa.gov" ;
:cdm_data_type = "swath" ;
:institution = "NASA Goddard Space Flight Center" ;
:GRingPointLongitude = -90.9374, -122.728, -135.075, -91.5693 ;
:LongName = "VIIRS/NPP Snow Cover 6-Min L2 Swath 375m" ;
:identifier_product_doi_authority = "http://dx.doi.org" ;
:ProcessingCenter = "MODAPS-NASA" ;
:PGE_EndTime = "2017-04-15 20:18:00.000" ;
:publisher_email = "modis-ops@lists.nasa.gov" ;
:RangeEndingDate = "2017-04-15" ;
:SouthBoundingCoord = 28.67912f ;
:LocalGranuleID = "VNP10.A2017105.2012.001.2017106052546.nc" ;
```

:PGE_StartTime = "2017-04-15 20:12:00.000" ;
:identifier_product_doi = "10.5067/VIIRS/VNP10.001" ;
:NorthBoundingCoord = 54.35787f ;
:GRingPointLatitude = 33.4794, 28.6791, 47.6999, 54.2566 ;
:DayNightFlag = "Day" ;
:license = "http://science.nasa.gov/earth-science/earth-science-data/data-information-policy/" ;
:EndTime = "2017-04-15 20:18:00.000" ;
:PGE_Name = "PGE507" ;
:EastBoundingCoord = -90.93742f ;
:project = "VIIRS Land SIPS Snow Cover Project" ;
:ShortName = "VNP10" ;
:creator_name = "VIIRS Land SIPS Processing Group" ;
:InputPointer =
"VNP35_L2.A2017105.2012.001.2017106052320.hdf,NPP_VIAES_L1.A2017105.2012.001.2017106050141.hdf,NPP_VMAES_L1.A2017105.2012.001.2017106050141.hdf,NPP_IMFTS_L1.A2017105.2012.001.2017106043639.hdf" ;


```

:InputPointer =
"/MODAPSops4/archive/f7053/running/VNP_LP_L5Sm7/1690846031/VNP10GA.A2018
008.h10v04.001.2019064202638.hdf,/MODAPSops4/archive/f7053/running/VNP_LP_L
5Sm7/1690846031/VNP10GA.A2018008.h10v04.001.2019064202638.hdf" ;
:Platform_Short_Name = "NPP" ;
:PGE_Name = "PGE543" ;
:AlgorithmVersion = "NPP_PR10A1 1.0.0" ;
:TileID = "51010004" ;
:ProcessVersion = "001" ;
:EndTime = "2018-01-08 23:59:59" ;
:ProductionTime = "2019-03-05 20:26:54.000" ;
:RangeBeginningDate = "2018-01-08" ;
:LongName = "VIIRS/NPP L3 Snow Global 375m SIN Grid" ;
:LocalGranuleID = "VNP10A1.A2018008.h10v04.001.2019064202654.h5"
;

:identifier_product_doi_authority = "http://dx.doi.org" ;
:InstrumentShortname = "VIIRS" ;
:PGE_EndTime = "2018-01-08 23:59:59.000000Z" ;
:ShortName = "VNP10A1" ;
:RangeEndingTime = "23:59:59.000" ;
:PGE_StartTime = "2018-01-08 00:00:00.000" ;
:StartTime = "2018-01-08 00:00:00" ;
:Conventions = "CF-1.6" ;
:GranuleDayNightFlag = "Day" ;
:ProcessingEnvironment = "Linux minion7053 3.10.0-957.5.1.el7.x86_64
#1 SMP Fri Feb 1 14:54:57 UTC 2019 x86_64 x86_64 x86_64 GNU/Linux" ;
:RangeBeginningTime = "00:00:00.000" ;
:DayNightFlag = "Day" ;
:PGEVersion = "1.0.3" ;
:AlgorithmType = "NPP_OPS" ;
:SatelliteInstrument = "NPP_OPS" ;
:identifier_product_doi = "10.5067/VIIRS/VNP10A1.001" ;
:GRingLatitude = 39.755674, 49.967704, 50.104591, 39.855422 ;
:HorizontalTileNumber = "10" ;

```

```
group: HDFEOS {
```

```
  group: ADDITIONAL {
```

```
    group: FILE_ATTRIBUTES {
      } // group FILE_ATTRIBUTES
    } // group ADDITIONAL
```

```
  group: GRIDS {
```

```
    group: NPP_Grid_IMG_2D {
```

```

dimensions:
  XDim = 3000 ;
  YDim = 3000 ;
variables:
  double XDim(XDim) ;
    XDim:standard_name = "projection_x_coordinate" ;
    XDim:long_name = "x coordinate of projection" ;
    XDim:units = "m" ;
  double YDim(YDim) ;
    YDim:standard_name = "projection_y_coordinate" ;
    YDim:long_name = "y coordinate of projection" ;
    YDim:units = "m" ;

group: Data\ Fields {
  dimensions:
    phony_dim_2 = 1 ;
  variables:
    ubyte Algorithm_bit_flags_QA(YDim, XDim) ;
      Algorithm_bit_flags_QA:long_name = "Algorithm bit flags QA" ;
      Algorithm_bit_flags_QA:valid_range = 0UB, 255UB ;
      Algorithm_bit_flags_QA:flag_masks = 1UB, 2UB, 4UB, 8UB, 16UB, 32UB,
64UB, 128UB ;
      Algorithm_bit_flags_QA:flag_meanings = "inland_water_flag
low_visible_screen low_NDSI_screen
combined_surface_temperature_and_height_screen_or_flag spare
high_SWIR_screen_or_flag spare solar_zenith_flag" ;
      Algorithm_bit_flags_QA:comment = "A bitfield of flags set for certain
conditions detected by data screens in the algorithm, multiple flags may be set for a
pixel." ;
      Algorithm_bit_flags_QA:grid_mapping = "Projection" ;
      Algorithm_bit_flags_QA:_FillValue = 255UB ;
    ubyte Basic_QA(YDim, XDim) ;
      Basic_QA:long_name = "Basic QA value" ;
      Basic_QA:valid_range = 0UB, 3UB ;
      Basic_QA:_FillValue = 255UB ;
      Basic_QA:key = "0=good, 1=poor, 2=bad, 3=other" ;
      Basic_QA:flag_values = 211UB, 239UB, 250UB, 252UB, 253UB ;
      Basic_QA:flag_meanings = "night ocean cloud no_decision bowtie_trim" ;
      Basic_QA:grid_mapping = "Projection" ;
    short NDSI(YDim, XDim) ;
      NDSI:long_name = "NDSI for all land and inland water pixels" ;
      NDSI:valid_range = -1000s, 1000s ;
      NDSI:_FillValue = 32767s ;
      NDSI:scale_factor = 0.001f ;
      NDSI:flag_values = 21100s, 23900s, 25100s, 25200s, 25300s, 25400s ;

```

```

        NDSI:flag_meanings = "night ocean L1B_missing L1B_unusable
bowtie_trim L1B_fill" ;
        NDSI:grid_mapping = "Projection" ;
        ubyte NDSI_Snow_Cover(YDim, XDim) ;
        NDSI_Snow_Cover:long_name = "Snow cover by NDSI" ;
        NDSI_Snow_Cover:valid_range = 0UB, 100UB ;
        NDSI_Snow_Cover:flag_values = 201UB, 211UB, 237UB, 239UB, 250UB,
251UB, 253UB, 254UB ;
        NDSI_Snow_Cover:flag_meanings = "no_decision night lake ocean cloud
missing_data bowtie_trim L1B_fill" ;
        NDSI_Snow_Cover:key = "0-100=NDSI snow, 201=no decision,
211=night, 237=inland water, 239=ocean, 250=cloud, 254=L1B fill, 255=fill" ;
        NDSI_Snow_Cover:grid_mapping = "Projection" ;
        NDSI_Snow_Cover:_FillValue = 255UB ;
        int Projection(phony_dim_2) ;
        Projection:grid_mapping_name = "sinusoidal" ;
        Projection:longitude_of_central_meridian = 0. ;
        Projection:false_easting = 0. ;
        Projection:false_northing = 0. ;
        Projection:earth_radius = 6371007.181 ;
        ubyte granule_pnt(YDim, XDim) ;
        granule_pnt:long_name = "Granule pointer" ;
        granule_pnt:valid_range = 0UB, 254UB ;
        granule_pnt:_FillValue = 255UB ;
        granule_pnt:grid_mapping = "Projection" ;
    } // group Data\ Fields
} // group NPP_Grid_IMG_2D
} // group GRIDS
} // group HDFEOS

group: HDFEOS\ INFORMATION {
    variables:
        string StructMetadata.0 ;

    // group attributes:
        :HDFEOSVersion = "HDFEOS_5.1.15" ;
} // group HDFEOS\ INFORMATION
}

```

11.0 Appendix C VNP10A1F Contents

Example of VNP10A1F file contents.

```
netcdf VNP10A1F.A2018008.h10v04.001 {  
  
// global attributes:  
    :TileID = "51010004" ;  
    :VerticalTileNumber = "04" ;  
    :HorizontalTileNumber = "10" ;  
    :GranuleDayNightFlag = "Day" ;  
    :GranuleDayOfYear = "8" ;  
    :GeoAnyAbnormal = "False" ;  
    :DataColumns = 3000s ;  
    :DataRows = 3000s ;  
    :ZonIdentifier = 0s ;  
    :GlobalGridRows = 54000 ;  
    :GlobalGridColumns = 108000 ;  
    :CharacteristicBinSize = 370.650173222222 ;  
    :CharacteristicBinAngularSize = 12. ;  
    :NorthBoundingCoord = 50. ;  
    :SouthBoundingCoord = 40. ;  
    :EastBoundingCoord = -91.369808 ;  
    :WestBoundingCoord = -124.45791 ;  
    :Snow_Cover_Extent = "80.5%" ;  
    :Cloud_Cover_Extent = "1.6%" ;  
    :Land_Day_Extent = "100.0%" ;  
    :QAPercentGoodQuality = "68.6%" ;  
    :QAPercentPoorQuality = "29.8%" ;  
    :QAPercentOtherQuality = "0.0%" ;  
    :ProductionDateTime = "2019-04-09T13:42:42.000Z" ;  
    :ShortName = "VNP10A1F" ;  
    :LongName = "VIIRS/NPP L3 Snow Global CGF 375m SIN Grid" ;  
    :VersionID = "001" ;  
    :PGEName = "PGE656" ;  
    :PGEVersion = "1.0.0" ;  
    :ProcessingEnvironment = "Linux nppdev-c7 3.10.0-693.11.6.el7.x86_64  
#1 SMP Thu Jan 4 01:06:37 UTC 2018 x86_64 x86_64 x86_64 GNU/Linux" ;  
    :SatelliteInstrument = "NPP OPS" ;  
    :SensorShortName = "VIIRS" ;  
    :LocalGranuleID =  
"VNP10A1F.A2018008.h10v04.001.2019099134242.he5" ;  
    :LocalVersionID = "1.0.0" ;  
    :InputPointer =  
"/NPPdata/griggs/AS3251/VNP10A1.A2018008.h10v04.001.2019064202654.h5,/NPPd  
ata/griggs/CM_PGE543_V1.0.1/VNP10A1F_test/VNP10A1F.A2018007.h10v04.001.he  
5" ;
```



```

:FirstDayOfSeries = "N" ;
:MissingDaysOfVNP10A1 = 0s ;
:TimeSeriesDay = 8s ;

group: HDFEOS {

group: ADDITIONAL {

group: FILE_ATTRIBUTES {
} // group FILE_ATTRIBUTES
} // group ADDITIONAL

group: GRIDS {

group: NPP_Grid_IMG_2D {
dimensions:
XDim = 3000 ;
YDim = 3000 ;
variables:
double XDim(XDim) ;
XDim:standard_name = "projection_x_coordinate" ;
XDim:long_name = "x coordinate of projection" ;
XDim:units = "m" ;
double YDim(YDim) ;
YDim:standard_name = "projection_y_coordinate" ;
YDim:long_name = "y coordinate of projection" ;
YDim:units = "m" ;

group: Data\ Fields {
dimensions:
phony_dim_2 = 1 ;
variables:
ubyte Algorithm_Bit_Flags_QA(YDim, XDim) ;
Algorithm_Bit_Flags_QA:long_name = "Algorithm bit flags QAsnow cover"
;
Algorithm_Bit_Flags_QA:comment = "A bitfield of flags set for certain
conditions detected by data screens in the algorithm, multiple flags may be set for a
pixel." ;
Algorithm_Bit_Flags_QA:flag_meanings = "inland_water_flag
low_visible_screen low_NDSI_screen
combined_surface_temperature_and_height_screen_or_flag spare
high_SWIR_screen_or_flag spare solar_zenith_flag" ;
Algorithm_Bit_Flags_QA:flag_masks = 1UB, 2UB, 4UB, 8UB, 16UB,
32UB, 64UB, 128UB ;
Algorithm_Bit_Flags_QA:grid_mapping = "Projection" ;
ubyte Basic_QA(YDim, XDim) ;

```

```

Basic_QA:long_name = "Basic QA valueed NDSI snow cover" ;
Basic_QA:valid_range = 0UB, 3UB ;
Basic_QA:_FillValue = 255UB ;
Basic_QA:key = "0=good, 1=poor, 2=bad, 3=other" ;
Basic_QA:flag_values = 211UB, 239UB, 250UB, 252UB, 253UB ;
Basic_QA:flag_meanings = "night ocean cloud no_decision bowtie_trim" ;
Basic_QA:grid_mapping = "Projection" ;
ubyte CGF_NDSI_Snow_Cover(YDim, XDim) ;
CGF_NDSI_Snow_Cover:long_name = "Cloud Gap Filled NDSI snow
cover" ;
CGF_NDSI_Snow_Cover:valid_range = 0UB, 100UB ;
CGF_NDSI_Snow_Cover:flag_values = 201UB, 211UB, 237UB, 239UB,
250UB, 251UB, 253UB, 254UB ;
CGF_NDSI_Snow_Cover:flag_meanings = "no_decision night lake ocean
cloud missing_data bowtie_trim L1B_fill" ;
CGF_NDSI_Snow_Cover:key = "0-100=NDSI snow, 201=no decision,
211=night, 237=inland water, 239=ocean, 250=cloud, 254=L1B fill, 255=fill" ;
CGF_NDSI_Snow_Cover:_FillValue = 255UB ;
CGF_NDSI_Snow_Cover:grid_mapping = "Projection" ;
ubyte Cloud_Persistence(YDim, XDim) ;
Cloud_Persistence:long_name = "consecutive days of cloud cover" ;
Cloud_Persistence:valid_range = 0UB, 254UB ;
Cloud_Persistence:_FillValue = 255UB ;
Cloud_Persistence:comment = "count of consecutive days of cloud cover"
;
Cloud_Persistence:grid_mapping = "Projection" ;
int Projection(phony_dim_2) ;
Projection:grid_mapping_name = "sinusoidal" ;
Projection:longitude_of_central_meridian = 0. ;
Projection:false_easting = 0. ;
Projection:false_northing = 0. ;
Projection:earth_radius = 6371007.181 ;
ubyte VNP10A1_NDSI_Snow_Cover(YDim, XDim) ;
VNP10A1_NDSI_Snow_Cover:long_name = "Daily VNP10A1 NDSI snow
cover for today" ;
VNP10A1_NDSI_Snow_Cover:valid_range = 0UB, 100UB ;
VNP10A1_NDSI_Snow_Cover:flag_values = 201UB, 211UB, 237UB,
239UB, 250UB, 251UB, 253UB, 254UB ;
VNP10A1_NDSI_Snow_Cover:flag_meanings = "no_decision night lake
ocean cloud missing_data bowtie_trim L1B_fill" ;
VNP10A1_NDSI_Snow_Cover:key = "0-100=NDSI snow, 201=no
decision, 211=night, 237=inland water, 239=ocean, 250=cloud, 254=L1B fill, 255=fill" ;
VNP10A1_NDSI_Snow_Cover:_FillValue = 255UB ;
VNP10A1_NDSI_Snow_Cover:grid_mapping = "Projection" ;
} // group Data\ Fields
} // group NPP_Grid_IMG_2D

```

```
    } // group GRIDS
  } // group HDFEOS

group: HDFEOS\ INFORMATION {
  variables:
    string StructMetadata.0 ;

  // group attributes:
    :HDFEOSVersion = "HDFEOS_5.1.15" ;
} // group HDFEOS\ INFORMATION
}
```